








Plate XVI







COAL FISH



Ancw. Clark
597
N53
A NEW SYSTEM

OF THE

NATURAL HISTORY

OF

FISHES AND INSECTS.

-Volucrumque genus.

BUCH. SAT. IN CAR. LOTH. CARD.

EDINBURGH:

ERINTED FOR PETER HILL, EDINBURGH 3
and
THOMAS CADELL, LONDON.

M.DCC.XCII.



a namu daya yangan

Vision Service Co. Lora, Chan.

HOSTENIGE

SHOWN KILL "YOUR MEET, I TON DRY UND

the line of the continues

TORDOLL SOIL

\$17 0 3 100 \$17 0 3 100 597 QL N53 45 N53 N53

CONTENTS.

NATURAL HISTORY OF FISHES.

	Pa	ge		Page
OF	Fishes in general,	I	Spec. 2. Sea Lamprey, -	92
		ib.	3. River Lamprey, -	96
	2. Of the external parts, and		4. Pride,	97
	motion of Fifbes,	8	Gen. 6. Ray.	99
	3. Of the respiration, food,	. 1	Spec. i. Skate,	105
		17	2. Sharp hofed Skate,	106
	4. Of the generation, fecundi-		3. Torpedo, or Electric	
		23	Ray,	107
	5. Of the growth, longevity,		4. Fire-flare, or Sting,	110
	and distical uses of Fishes,	31	5. Thornback, -	113
			Gen. 7. Shark; -	115
	Order I Cetaceous Fishes,	1	Spec. I. Angel,	IIO
			2. Basking Shark, -	IZE
	Genus I. Common Whale.		3. White Shark,	123
Spec.	1. Common Whale, -	55	4. Blue Shark, -	125
-		57	5. Long tailed Shark, or	
	Gen. 2. Narwhal, or Sea-		Sea Ape,	129
	Unicorn,	59	Gen. 8. Angler, -	129
	Gen. 3. Cachalot,		Spec. I. Common Angler, -	130
Spec.	I. Cachalot,	64	Gen. 9. Sturgeon,	132
	2. Blunt-headed Cachalot,	68	Spec. I. Common Sturgeon,	133
	3. Round-headed Cachalot,	69	2. Huso of Germany, -	136
	Gen. 4. Dolphin.		Gen. 10. Diadon, -	138
Spec.	I. Dolphin,	72	Spec. I. Globe,	139
1	2. Porposse,	77	2. Oblong Diadon, -	140
	3. Grampus, 2	80	Gen. II. Sucker, -	I42
			Spec. I. Lump,	ib.
0	rder II Cartilaginous Fiftes.		2. Sea Snail,	144
	1		Gen. 12. Soa Porcupine,	145
	Gen. 5. Lamprey, - 8	39	Gen. 13. Oftracion,	147
Spec.	I. Muræna, 5	94		

Page	t Pa	A 54
Page	Pa	
Gen. 14. Pipe Fift, 149		16
Spec. I. Longer Pipe Fish, - ib.		17
2. Hippocampus, or Sea-		ib.
Horfe, - 150	Gen. 36. Wraffe, 23	20
0 1 III 90 F.0	1 222 00	21
Order III Spinous Fishes.		22
Division I. Of Apodal Fishes.		24 ib.
		26
Gen. 15. Eel, - 157	- n: n :	27
Spec. 1. Common Eel, - 159 2. Conger Eel, - 160	2. Baffe, - = 22	28
2. Conger Eel, - 160 3. Sea Serpent, - 162		30
Gen. 16. Sea Wolf, 164	Spec. I. Three fpined Stickle-	,0
17. Sand-Eel, or	back, - 23	+
Lanco, 166	2. Fifteen fpined Stickle-).1
18. Ophidium, 167		ь.
19. Stromateus, 168	Gen. 40. Mackrel, 23	
	Spec. I. Tunny, - 23	-
20. 8:00/4-1/3, 109		38
Division II. Jugular Fishes.	Spec. I. Striped Surmullet, 23	500
Divinoit II. Jagaiar Pipess.	Gen. 42. Gurnard, - 24	
Gen. 21. Pragonet, 172	Spec. I. Gray Gurnard, - 24	
22. Uranoscopus, 175	Spec. 1. Gray Garnard,	42
-23. Weever, - 176	Division IV. Abdominal Fishes.	
24. Cod.	Division 2 v. 22000minus 21 juess	
P 0:10	Gen. 43. Loche, - 24	12
2. Haddock, 182 3. Coal Fifh, 184	-44. Amia, = 24	
_3. Coal Fifh, - = 184	45. Silurus, 24	-
	-46. Teuthis, - 24	
4. Hake, 185 5. Ling, 187	- 47. Salmon,	,0
6. Five Bearded Cod, 190		10
7. Torsk, 191	Spec. 1. Salmon, 24 2. Sea Trout, 25	
Gen. 25. Blenny, 193	3. River Trout, = 25	
Spec. 7. Gattorugine, - 194	61	
2. Viviparous Blenny, 195	4. Charr, 25 5. Gwiniad, 25	
	Gen. 48. Pike, 25	
Division III. Thoracie Fiftes.	Spec. 1. Gar, 26	
A STATE OF THE PARTY OF THE PAR	Gen. 49. Elops, 26	
Gen. 26. Cepola, 197	50. Argentine, 26	
27. Remora, - 199	51. Atherine, 26	
	52. Mullet, 26	
Spec. 1. Pompilus, - ib. Gen. 29. Goby, - 202	- 53. Flying Fift, 27	
Gen. 29. Goby, - 202	-54. Polynemus, 27	
30. Bull-bead, 203	- 55. Mormynis, 27	
Spec. I. River Goby, ib.	56. Herring, 27	
2. Father Lasher, - 205	Chec T Dilaband	
Gen. 31. Dorce, ~ 207	2. Anchovy, 28 3. Shad, - 28	2,
Spec. I. Opan, - 208	3. Shad, - 28	a
Gen. 32. Scorpion, 210	3. Shad, - 28 Gen. 57. Carp, - 28 Spec. 1. Carp 28	7
33. Flounder, - 212	Spec. I. Carp, 28	
Spec. I. Holibut, 213	2. Barbel, 29 3. Tench, 29	I
2. Turbot, 214	3. Tench, 29.	3
	Charles and the second second	

NATURAL HISTORY OF INSECTS.

P	age	Page
Or Insects in General, -	305	Gen. 9. Caffidz, - 376
General idea of the Number and	•	- 10. Coccinella, 377
Variety of Infects Induce-		- II. Chryfomela, 379
ments to the study of Entymology.		12. Hispa, - 381
-History of this Science,	ib.	13. Bruchus, 382
The Literary History of Enty-		—— 13. Bruchus, 382 —— 14. Curculio, 383
	318	15. Ettelabus, - 385
A comparative view of the Senses		16. Cerambys.
and Endowments of Infects,		The Capricorn, 586
	326	17. Leptura. The
Of the external Parts, and Classi-		Wood Beetle, - 387
	335	18. Necydalis. The
Of the Generation and Metamor-		Carrion Eater, - 388
	341	19. Lampyris. The
Of the Habitation and Food of In-		Fire-fly, - 389
feets, and their uses in the econo-	240	20. Cantharis, 390
my of Nature, -	349	Slipper
Order I Coleopterous Infects,	355	Skipper, = 391 22. Cicindela. The
Oraci i. donoprervas injecis,	222	0
Genus I. Scarabæus.		Sparkler, 392
	356	Covo Burner, - 393
Spec. I. Bull-comber, -	357	- 24. Dytifeus. The
2. Dung Beetle, -	358	Diver, - 394
3. Golden Beetle, -	360	- 25. Carabus. The
	361	
Spec. I. Stag Beetle, -	3.62	26. Tenebrio. The
2 Parallelipiped Beetle,	363	Darkling, - 397
Gen. 3. Dermestes,	365	-27. Meloe. The
Spec. 1. Domestic Dermestes,	366	Bloffom-eater, - 398
2. Dermestes Lardarius,	367	28. Mordella. The
Gen. 4. Ptinus,	369	Nilbler, - 401
- 5. Hilter	371	- 29. Staphylinus, 402
O. Gerinus,	372	30. For fivula. The
- J. Byrrous, -	374	Earwig, - 493
8. Silpha, -	275	

Page	Pag	ze.
	Gen. 4. Ichneumon, 49	6
Order II Hemiptera, 405		
1	- 6. Chrysis. The Golden Fly, 50	
Genus I. Blatta, - 406		
2. Mantis, or Soothfayer, 408		
- 3. Gryllus, 410		
4. Fulgora, or Lantborn-fly, 422	W 41 111	
5. Gicada, or Flea Locust, 424		
- 6. Notonecta, or Boat-fly, 427	Order VI Depterous Infects, 52	6
7. Nepa, or Water Scor-		
pion, 429	Gen. I. Deftrus. The Gad Fly, 52	7
8. Gimex. The Bug, 431	- 2. Tipula. The Crane Fly, 53	0
- 9. Aphis. The Plant Loufe, 436	- 3. Musca. The Fly, - 53	3
10. Chermes, 440		
II. Coccus. The Cocwineal, 443	or Flogg, - 53 5. Culex. The Gnat, 53	8
12. Thrips, 447	5. Culex. The Gnat, 53	9
	- 6. Empis, - 54	Í
Order III.—Lepidopterous Infects, 448	7. Conops, 54 	2
	8. Afilis, 54	3
Gen. I. Papilio. The Butterfly, 449	- 9. Bombilius. The Buzz	
- 2. Sphinx, - 455	Fly, - 54	4
- 3. Phalenc. The Moths, 458	Io. Hippobosca. The Horse	
0.1.777	Leech, 54.	5
Order IV.—Neuropterous Infects, 468		,
G 11111 G1 D	Order VII.—Apierous Infects, 54	6
Gen. I. Libellula. The Dragon	G = 7 436	
Fly, 469		
2. Ephemera. The Day	- 2. Podura. The Springtail, 54	-
Fly, 474	3. Termes, 55	
Fly, - 474 - 3. Phryganea, - 478 - 4. Hemerobius, - 480 - 5. Myrmeleon, - 482	4. Pediculus. The Loufe, 55	
4. Hemerobius, - 480		
5. Myrmoleon, - 482	- 6. Acarus. The Tick, 550	
6. Panorpa. The Scorpion, 485	7. Phalangium, - 55	
7. Raphidia, 487	- 8. Aranea. The Spider, 550	
Order V - Hamenotteraus I-Cas. 199	9. Scorpio. The Scorpion, 56	
Order V Hymenopterous Infects, 488	IO. Cancer. The Grab, 56	
Gen. 1. Cynips. The Gall Fly, 489		
- 2. Tenthredo. The Saw Fly, 492	12. Scolopendra, - 57.	-
3. Siren. The Tailed Wafp, 494		4
J. Diren. 2 de Latteu Wajp, 494		

DIRECTIONS FOR THE BINDER.

YOL. III.

FISHES.		Fishes.	
PLATE I. to face Pag	e 55.		233.
2.	72.	26.	249-
3.	90.	27.	257-
4.	99•	28.	259.
5.	IIC.	29.	269.
6.	119.	30.	288.
7-	127.	31.	295
8.	133.		
9.	142.	INSECTS.	
IO.	145.		
II.	149.	PLATE 1. to face Page	356.
12.	157.	2.	362.
13.	169.	3.	383-
44.	172.	4 o	396.
15.	178.	5. 6.	410.
16.	184.	6.	449.
17.	191.	7.	468.
18.	194.	8.	474-
19.	202.	9.	489.
20.	207.	IO.	508.
21.	213.	11.	547-
22.	214.	12.	558.
23.	217.	13,	569.
2.1,	270.	-	



NATURAL HISTORY

O I'

FISHES.

CHAPTER I. OF FISHES IN GENERAL.

SECT. I .- History of Ichthyology.

In the early periods of fociety, the necessities and wants of men, rather than their curiosity, prompt them to pay attention to the different objects, which the munificent Author of nature has placed around them: And, though sishes constitute a greater part of the subsistence of men, in their rude and savage state, than perhaps in any suture stage of their political existence, yet their acquaintance with this part of the animal kingdom, in the earlier ages of society, is by no means extensive. A sew species that are most common, or most easily taken in the nearest river or lake, are all that the necessity of the savage requires; and, as in him, curiosity has not yet begun to be a sleady principle of action, it seldom leads him to examine more.

Vol. III. A Even

Even after nations have attained to fome degree of knowledge and civilization, many ages clapse before they pull their inquiries far into the subject of ichthyology, or acquire any confiderable acquaintance with the inhabitants of the ocean. In the unfathomed depths of that turbulent and extensive element, probably millions refide, of which the far greater part are feeluded from human observation; and, even of the few which the industry of man has, at last, drawn from their hidden abode, we hardly know any thing, but the external figure, and the names. Their food, their longevity, their method of propagating their kind, and the whole of their manners and economy, remain fill among those numberless secrets of Nature, which human ingenuity has not hitherto been able to explore. Hence, the natural history of fishes has feldom been found interesting, because it is destitute of that information, which it is the province of history to convey. It is more imperfect and obscure, than that of quadrupeds and birds, in proportion as the element, in which fishes reside, is more extensive and inaccessible.

Before the Christian era, few writers had turned their attention to this difficult branch of Natural History. Aristotle, Theopkrasius, Strabo, and Terentius Parro, had made their observations on such as were then known; but they had scarcely any idea of treating the subject in a scientistic manner. After that period, Appian, Ovid and Columbia, Schribed the sithes of the Euxine and Adriatic seas, as far as they had access to examine them. Pliny was the last, and by far the most copious writer upon this subject, among the ancients. He is, indeed, too diffuse; while his credulity and love of the marvellous, considerably weaken the authority of his narrative. An hundred and twenty-sour species,

DSI

were

were all that the affiduity of the ancient naturalists had discovered.

During the decline and fall of the Roman empire, attention to ichthyology, as well as to every other part of literature, was entirely laid afide; nor did the pernicious effects of the Gothic invafions, allow it to be refumed, till the year 1524, when Paulus Jovias*, an Italian writer, gave an account of those fishes, that were known to the ancient Romans. After him, various local historians appeared. Belonius and Rondeletius succeeded each other in describing the fishes of the Mediterranean; while Swinfield, Marcgrave, and Catesby, gave, successively, an account of the fishes of Silesia, Brasil, and the Carollinas.

While ichthyology was thus enriched by historians of particular districts, there were other naturalists, who engaged in this science in a more general and systematic manner. The Honourable Francis Willoughby published, in 1686, a history of sishes, which is still deemed one of the most valuable upon this subject. He was afterwards followed by Artedis and Linneus, who carried the science to a greater degree of perfection, than it had ever hitherto attained.

If we take into account the vast number of fishes, of which the very names have not yet found a place in the systems of naturalists, we must necessarily conclude, that this part of science, is still in its infancy. Some very skilful ichthyologists affert, that there are to be found in the different collections of sishes about London, six hundred kinds not enumerated by Linneus; and we may perhaps, easily admit, that there are yet, in the unsathomable depths of the ocean, at least an equal num-

A 2 ber,

[&]quot; De piscibus Romanis.

ber, that have not come into the possession of the curi-

It is remarkable, that there is no fyshem of ichthyology in the English language. Goldsmith's plan did not admit of any thing more, than a general sketch of this subject. Accordingly, he has not described, perhaps, a twentieth part of those, that are already found in the systems of Artedi or Linnæus. Pennant has, indeed, given a correct and elegant history of the British sishes; but these make only a small part of the inhabitants of the ocean. We dare not promise, in the narrow limits of the following work, to offer a complete system to the public. All the genera, however, are included, and a few of the most remarkable species belonging to each, are selected for a particular description.

By the labours, however, of the different philosophers already mentioned, nearly five hundred kinds of fishes have been enumerated, and deferibed; and, in appearance at least, considerable progress has been made in explaining their history: But, unfortunately, the names and external figure of many of these are all that we know. Their food, migrations, manner of life, and every quality that can render their history interesting, still remain to be explored. While, therefore, we are guided principally by the arrangement of Linnaus, in giving the following account of fishes, we will spare our readers the trouble of perufing the tedious catalogue, as often as it prefents nothing, but the dry detail of names. We shall thus fecure ourselves an opportunity of enlarging more fully upon those particular fishes, whose history may appear most authentic, curious, or useful.

Ariffette, that great father of naturalists, first suggests od the excellent arrangement of fishes, into the cetaccous, cartilaginous:

cartilaginous, and fpinous.*; which, as far as it goes, feems impossible to be altered for the better †. Rondeletius, the first naturalist, who, after the revival of learning, turned his attention to this subject, attempted to lay afide the Aristotelian division, and to suffictute, in its room, another, founded upon the habitation of silves, or those places where they reside. He, accordingly, classed them into the sishes of the fea, rivers and lakes †. Since, however, many silves reside indiscriminately in all these situations, this method was abandoned by Willoughby and Ray, who again resumed the arrangement of Aristotle; and, the celebrated Linneus has added to it some tarther subdivisions, which render it still more useful.

The cetaceous fishes have, in the later editions of his works, been classed among the quadrudeds; the cartilaginous have been referred to the amphibious tribes, among which they constitute a particular order §; while the spinous or bony, are alone allowed to retain the name of fishes.

As we have already hinted, that we are to pass by, or but slightly mention, some species enumerated by that naturalist, whose manners are not interesting, or but inspersectly known; so, on the other hand, we shall restore to their natural station among the sinny tribes, the cetaceous and cartilaginous sisses, which he has banished from that part of the animal kingdom.

The grand chanical characters, which nature has imprinted on the cetaceous order, will, in a philosophical wiew,

^{*} Hift. Animal.

[†] The first comprehends the whole kind; the second, these whose fish is supported by extillages; and the third, those whose muscles are supported by spines or body matter.

[#] Marini, Fluviatiles, and Lacuftres.

[§] Amphibia Nantia, Lin. Syft. Nat. Ed. 12.

view, vindicate their arrangement among quadrupeds. Their internal structure agrees, in every respect, with that of the nimmalia quadrupedia of Linneus; and their external conformation also is, in some parts, similar.

The cetaceous fishes are destitute of gills: They breathe by means of lungs, and, on that account, are obliged to rife frequently to the furface of the water for respiration.* They resemble land animals too in having warm blood; in being provided with external organs of generation; and in their manner of copulating and bringing forth their young, which they suckle, and protect with parental attachment. They have the power of uttering founds. fuch as of bellowing and making other noises; a faculty denied to the other inhabitants of the deep.

But notwithstanding this striking similarity between the cetaceous sishes, and the land animals, there are many other properties belonging to the former, which must determine us to rank them among sishes, where the general voice and language of men have always placed them. The feals, manati and whales, are evidently the steps, by which Nature proceeds from the one of these her great families to the other: Here they approximate, and appear kindred tribes. In the last of these species of animals however, the great outlines and form of a sish predominate: It is entirely naked, or covered with a smooth skin; it lives wholly in the water, and has all the actions and habits of its aquatic neighbours ‡.

It is with much greater confidence and facility, that we reftore to their original flation, the cartilaginous fifthes a

^{*} Vide British Zoology.

[†] Blaffie's Anat. Animal. 288.

Vide Raii Synop. Pifc.

iffhes; because greater violence has been used in excluding them from it. They are by no means capable of living indiscriminately in the water and on shore, and, therefore, merit not the name of amphibious, which Linameus has assigned them. The single circumstance, by which that naturalist determined their rank in the animal kingdom, is the want of the bony operculum, or covering of the gills, which is common to the spinous sisses.

This last order, which alone he admits to be files, he has very properly subdivided into four different sections, the apodal, the thoracic, the jugular, and acidominal, fishes *. This arrangement is founded on the comparison of the ventral sins to the feet of land animals; and the particular situation of these fins with respect to the rest, determine the place, where each sish is to be ranked.

The apodal fishes, are such as want the ventral sins altogether, such as the fwordsish and cel. In the jugular, the ventral sins are placed before the pectoral, as is exemplified in the codsish and bienny. The thoracic are distinguished by having the ventral sins placed beneath the pectoral, as is illustrated by the mackrel, or father is no. The abdominal sishes are known, by having the ventral sins placed behind the pectoral sins, near to the abdomen, as is the case in the falmon and pike. This division of sishes, according to the situation of their sins, is exceedingly judicious and natural. It sirst occurred to Linneus, when examining a collection of prepared subjects, in the presence of Dr. Sollander, who witnessed the extasy of this indefatigable naturalist on making this discovery.

SECT.

^{*} Vide Systema Nat. p. 422.

f He used the exciamation of the Gresian mathematician Archimides. Evenue.

SECTION II.

Of the external Parts and Motion of Fishes.

THE external form of the greater part of fishes, tends greatly to the case and celerity of their motion. It is sharp at either end, and swelling towards the middle; and is modelled by nature after that shape, which we endeavour to imitate in those veriess, that are intended for the greatest despatch. Every human contrivance, however, falls far short of the rapidity of the natives of the sea. All the larger sishes can easily overtake the best constructed vessel while in sull fail; and play around it, without any apparent effort.

The principal indrument of this great velocity in fillies, is the tail, aided by the flrength and flexibility of the back-bone. The other fins are too finall, flender, and flexible, compared with the weight of the animal, to impel it through the water with fuch vast rapidity and force. Their principal use is to direct and moderate the movement, communicated by the impulse of the tail. A fish, when deprived of these, and put into a pond, darts upwards, downwards, and laterally, with all its wonted velocity, but without being able to direct its course *. The ventral and dorsal fins, serve the purpose of keeping the fishes in an erect and perpendicular polition; and, perhaps, the first contributes to raise or fipls them in the water. The pectoral fins regulate and affilt progressive motion. When swimming rapidly forward.

^{*} Berelli apud Will. de motu anim.

ward, fishes can, by extending these, stop their motion, and produce mora; and, when swimming in a streight direction, they can, by folding either, while the other continues to ply, direct their motion to that side. The size of these fins, is, in general, proportioned to that of the head of the sish to which they belong; and it is probably one use of them, to prevent the sish from being overbalanced by the weight of its head, and precipitated to the bottom. In some sishes, the pectoral sins serve the same purpose as the wings of birds; for, by their means, they are enabled to rife from their watery element, and to sly for a considerable space, till their sins be so dried by the air, that, in spite of every exertion, they again sink into the water.

A fish completely equipped for swimming, has seven sins, two pairs, and three single ones; and of the latter, two are above, and one below: But, since those which have the greatest number of sins, are not the swiftest, it has been concluded, that the tail is the principal cause of the swiftness of a sish's motion. When in pursuit of its prey, or avoiding an enemy, all the smaller sins are laid close to its body; then, by the impulse of the tail alone, it skims through the water with the incredible velocity of a dart or an arrow * The muscles, by which the tail is moved, are, by far, the thickest and strongest of the whole body; and to give direction to the great impetus which they communicate to the sish, seems to be the chief purpose of all the smaller sins.

The motion of fishes, is supposed to be affished by the swimming bladder, which, by contraction and dilatation, serves to raise or fink them in the watery element at pleasure; and, as that element is of very different degrees

Vol. III. B of

^{*} Willoubii Ichthyol. p. 4.

of weight, according to its depth, the fishes, by thus varying their specific gravity, can poile themselves in any part of it *. That this is the use of the air bladder, fome naturalists pretend to have ascertained by experiment: An incision made into this organ, by which the air is allowed to escape, obliges the fishes, after some efforts, to fink to the bottom of the water. This opinion feems to receive confirmation from the manuers of those kinds where it is wanting. All flounders and ground-fithes are destitute of a swimming bladder; and, by confequence, lie constantly at the bottom; while the cetaccons and cartilaginous orders, which often support themselves at the surface, without the aid of this infirument, are supplied with lungs, which serve the same purpose, by admitting the air, in much greater quantity, into the cavity of their bodies.

So far this matter feemed to be clear; but, upon examining this fingular organ, it appeared to be supplied with no muscles that could enable the animal to contract or dilate it, in a voluntary manner. Two other opinions have, therefore, been formed concerning the use of the swimming bladder. Dr. Necdbam observed, that in the greater part of sishes, whether this organ was single, as in some species, or double, as in others, it was furnished with a duct or canal that opened into the stomach: Hence he concluded, that its use was to collect the air secreted from the blood, and to convey it into the stomach, where it affished the process of digestion. In this manner, he endeavoured to account for the extraordinary voracity of sishes, and their uncommon powers of digestion.

No

Willoubii Ichthyol. p. 13.

No philosopher seems yet to have made observations sufficiently accurate, fully to determine this matter; for it is not yet ascertained, whether the neck of the canal be supplied with a valve to emit air secreted from the blood, or to receive it from the stomach, or with a muscle that may equally serve to admit, or eject it: And what militates against Needbam's opinion, is, that the cartilaginous sisses, which have nothing of this kind to aid their digestive powers, are as remarkable for voracity as any other kind.

The other use to which naturalists have destined this organ is, to serve as a receptacle for air, instead of the lungs in terrestrial animals, by which the necessities of these creatures may be supplied. The air bladder of sistes is confined in a triangular space at the upper part of the abdomen, and consists of one, two, and sometimes three divisions, all separated from the rest of the viscera by a thin membrane, attached, on each side, to the ribs. An apparatus of this kind, though we be not able to explain its use, no one can imagine was formed in vain. It is possible, that it may serve some purpose different from any of the above, and probably more than one even of these.

Fishes thus fitted for motion in their element, by their internal structure, as well as by their outward shape, and the fituation of their fins, scem as well furnished with the means of happiness, as either quadrupeds or birds. Like these they are furnished with an external covering, to desend them from injuries in the turbulent sluid which they inhabit. That slimy and glutinous substance, which is secreted from the pores of all sishes, not only desends their bedies from accident, but is happily contrived to give little obstruction to their progress through the water. Besides thus substance, which desends their lives

from the immediate contact of the furrounding fluid, the greater part of fishes are provided with a strong covering of scales, which still more powerfully protects them from injury; and, beneath this, they are supplied with an oily matter, which preserves the body in warmth and vigour.

When, however, we examine the fenses, and other faculties, of this part of the animal kingdom, we find it in a rank greatly below the other tribes; and that Nature having intended them for lets perfect beings, has been sparing in her endowments. The brain, the seat of fensation, is much smaller in sishes, than in other animals; and, probably, gives off a smaller number of nerves to the different parts of the body *. Those strong teguments, with which we have observed that their bodies are covered, must greatly obstruct their sense of touch, which, in all probability, is far from being delicate.

The external organs of fmell, and the nerves which fupply them, are perceptible in the greater part of fishes; and, even where the apparatus is not discernable without, the formation of the bones within, plainly indicate an apparatus for this purpose. But as air is the only medium we know for the distribution of odours, it cannot be supposed, that these animals, residing in water, can be possessed of any capacity of being assected by them. If they have any perception of smells, it must be in the same manner as we distinguish by our taste; and it is probable, that the olfactory membrane in fishes, ferves them instead of a distinguishing palate: By this they judge of substances, whose vapours, having tinctured the water, are sent to their nostrils, and, no doubt,

produce

^{*} Willisius de anat. cerebri apud Willoubium.

produce some kind of sensation. This, most probably, is the use of that organ in these animals; as, otherwise, they would be provided with the instruments of a sense, which they could not enjoy from want of an opportunity of using them *.

The fense of taste in sishes must be very imperfect, if its delicacy arise from the pliancy and softness of the organ. The whole mouth of sishes is covered with a hard bony substance, which must deprive them of almost all power of distinguishing different substances by the palate: Their voracity, accordingly, is so indiscriminate, that there is hardly any kind of food which they will not swallow: They devour the sisherman's plummet instead of the bait.

All fishes, except the cetaceous, are deprived, not only of external ears, but also of the auditory nerves and canal. Of the sense of hearing, therefore, it is probable, that they are altogether destitute †. As they are incapable of uttering founds, they could seldom have an opportunity of hearing, even though Nature had endowed them with that power. They have no voice to communicate with each other, and, consequently, have no need of that organ by which it is discerned. The whales, and reshaps the cartilaginous sishes, are found possessed in all apertures for the admission of sounds; but, even in these, this sense must, from the smallness of the organ, be very impersect.

Naturalitis, determined by these reasons, feem, in general, agreed, that all the spinous tithes are destitute of the faculty of hearing. Limitum who has probably examined a greater number of subjects, and with more accuracy than any other philosopher, contesses, that he

has

⁶ Goldsmith's Nat. Hist. vol. 6. p. 160.

[?] Raius apud Will. c. iv. lib. r. de pisc. auditu.

has not been able to discover any auditory organs in this class of the animal kingdom*. Mr. Klein, however, imagines he has found out organs of this sense; and he has allotted for that purpose, those bones which are sound in the head of some sithes †. But these are so dissimilar to the organs of hearing in other animals, that it is improbable, that Nature intended them for that purpose; and, besides, there are many species, in which they are not to be discovered.

But although Nature, in their conformation, had made a provision for the hearing of fishes, that sense must have been extremely limited and imperfect, from the nature of the element which they inhabit. Experiments have been made on the capacity of water to transmit founds; and by these it has been found, that it is capable of conveying them but a fhort way; for it quickly deadens that vibration upon which they depend. A man whose head is one foot immerfed in water, hears voices and words uttered in the air; but, when funk to the depth of twelve feet, he scarcely hears a musket shot, though discharged over his head. Hence it is probable, that Nature has made no organic apparatus to convey founds to fishes, fince fhe has forever configued them to an element, which must, in a great measure, have defeated its purpose.

Rondeletius †, and feveral of the old naturalists, who plead for the heaving of fishes, allege, in proof of it, that certain kinds of them are so assected with noise, that they become unwholesome after thunder; and that, in ponds where they are tamed, in some places of Germany, they are convened regularly by the call of a bell to their food.

^{*} Systema Naturæ. † Vide Historiam Piscium. † Rondel, de piscibus, lib. 7. cap. 14.

food. But the experiments made by Mr. Gorian*, professor of medicine at Montpelier, clearly demonstrate the fallacy of these proofs. The gold sishes, which he kept in vases, he could never disturb by the loudest noise, provided he could prevent the tremor of the air from asserting the water. It appears, therefore, that sishes are as destitute of hearing as of voice; and that, when they appear to come to their food at the call of a bell or whistle, it is either by feeling the vibrations of the sound assert the water, or by seeing the persons approach by whom they are accustomed to be fed †.

The fight of fishes is probably the most perfect of all their fenses, and yet it is far inferior to that of most other animals. They have, properly speaking, no eye-lids: Their fight is protected in the water by a nictating membrane, which is a continuation of the fame transparent skin that covers the rest of the head. The crystalline humour, which in most other animals is flat, is in them convex, and round like a ball t. In confequence of this, these animals must be near-sighted, even in water, which, like a concave glass, corrects, in some degree, this defect of the organ of vision. We have no evidence of any fishes seeing at a considerable distance; and the case with many of them, that are deceived by the different kinds of bait prepared in imitation of their food, gives room to suspect, that objects are not very distinctly perceived by them, even when near.

From this short account of the external senses of sishes, it must appear, that their faculties, in point of persection, fall greatly below those of the other tribes, which have al-

ready

Historia Pisc. † Linnæi Systema Naturæ.

de Goldsmith's Nat. Hist. vol. vi.

ready been reviewed in the preceding part of this work. In every thing refembling intelligence, their inferiority is equally striking. They are incapable of attachment, or of acquiring any new habits by domestication and intercourfe with man. Some faint traces, indeed, of memory they discover, if it be true, as is alleged, that they regularly return to the place where they have been fed *; but even this small share of recollection, can hardly be allowed them without hesitation; for they may be assembled there, merely by feeing one of their number pick up the remains of the food which they had formerly left. It forms no exception to the general conclusion, that all their powers and faculties are of a subordinate kind, suited to that humble and passive existence which Nature has affigned them. To preserve existence, and to continue it to posterity, fill up the whole circle of their purfuits and enjoyments, to which they are impelled rather by necessity than choice. While they feem mechanically excited to every fruition, their fenses are incapable of making any distinctions; and they are hurried forward in pursuit of whatever they can swallow, conquer, or enjoy +.

^{*} Willoub. Hift. Piscium, cap. iii.

[†] Goldsmith, Nat. Hist. vol. vi. p. 164.

The

SECTION III.

Of the Respiration, Food, &c. of Fishes.

In all animals, respiration, or the admission of air into the body, feems necessary to the support of life. From the experiments that have been made upon fishes, almost in the infancy of science, it appears, that they are incapable of subfishing without air for any considerable time. The cetaceous and cartilaginous fishes, are supplied with the necessary quantity of this sluid, by means of lungs like the terrestrial animals; and hence the Swedish naturalist has arranged them in the same class of beings *. In the spinous fishes, respiration is performed by branchia, or gills, without the cavity of the body; but the precife manner in which this operation is carried on, is one of those fecrets of Nature, which neither the glasses nor the knife of the anatomist have ever yet been able to develope. The manner, indeed, in which the air is transmitted from the lungs of quadrupeds into the blood, is perhaps equally mysterious, as its passage from the branchiæ of fishes into the arteries leading to the heart. This difficulty Artedi confesses; and Rondeletius, Needbam, and other philosophers, have in vain endeavoured to explain it.

Vol. III. C

^{*} Vide Systema Naturæ, last edition.

[†] Quomodo aer intret branchias piscium, dissicile est dictu.

The gills, it would appear, have a power of abforbing the air, without giving admission to the water which is received into the mouth, and ejected by the branchiæ, at each movement of the opercula which cover them; but in what manner the subtile separation is performed, no naturalist has ever ventured to declare. As the air and water pass quickly through the gills, without any appearent effort to separate them, it is probable, that only a small quantity of the former is absorbed, but that, by the frequent transition of the water, a sufficient quantity of it may be admitted to supply animals whose blood is naturally cold, and not in great abundance *.

But, however feall a quantity of air may fuffice to support the life of a fish, some portion of it is absolutely necessary to every living being. Some have been flut up in a narrow mouthed to lel, and have lived there for feveral years; but the air is no fooner excluded, by flopping up the opening, than the animals are suffocated in the course of a few minutes. When this experiment was repeated with the veffel filled half with water, and half with air, after covering the mouth, as the air below began to be exhausted, the fishes were seen struggling to rife above one another, and inhale a fmall supply at the furface of the water; and the fame confequence will follow, if you fathen down the opercula of the gills with a string. fo that respiration cannot be performed +. This probably is the cause why so many sishes are defroyed in rivers by fevere frost: When, by the congelation of the whole furface of the water, the external air is excluded, the animals below must necessarily perish t.

Aelian

^{*} Willoughby, lib. i.

[#] Willoughby, cap. v.

[†] Rondelet. lib. iv.

Actuan mentions a method of taking fishes in the river Isler, founded upon this observation. In the rigour of winter, the fishermen broke small holes through the ice. The suffocating animals instantly crouded to the aperture, in order to procure a supply of air; and so eager were they to obtain it, that rather than abandon the attempt, they suffered themselves to be caught with the hand.

Next to the necessity of breathing air, that of devouring food feems to be the most constant and urgent in the nature of fishes. Among them this appetite, both in flrength and avidity, feems to furpass those limits which Nature has prescribed to herself in the other orders of the animal kingdom. Every aquatic animal that has life, falls a victim to the indifcriminate voracity of one or other of the fishes. The smaller tribes devour insects, worms, or the spawn of the rest of the tenants of the waters; while they, in their turn, are purfued by millions larger and more rapacious than themselves. A few of them feed upon mud, the aquatic plants or grains of corn; but by far the greater numbers fubfift upon animal food alone; and of this they are fo veracious, that they spare not even those of their own kind*. Charr kept in a pond, if fcantily supplied, frequently devour their own young. Others that are larger, go in quest of a larger prey; it matters not of what kind, whether of another species, or of their own. Those with the most capacious mouths, purfue almost every thing that has life, and often meet each other in fierce opposition, when the fish which has the widest throat comes off with victory, and devours its antagonist +.

Thus

[&]quot; Willoughby, lib. i. cap. x.

Thus the rapacions fishes are different from the predatory kinds of terrestrial animals; they are neither few in number, nor folitary in their habits. Their rapacity is not confined to a few foecies, one region of the fea, or individual efforts. Almost the whole order is continually irritated by the cravings of an appetite, the strength of which excites them to encounter every danger, and which, by its excels, often destroys that life which it was intended to prolong. Innumerable floals of one species, purfue those of another through vast tracts of the ocean, from the vicinity of the pole, down to the equator. It is thus that the cod purfues the whiting, which flies before it, from the banks of Newfoundland, to the fouthern coasts of Spain. Thus too the cachalot drives whole armies of herrings from the regions of the north, devouring, at every instant, thousands in the rear. Hence, the life of every fish, from the smallest to the greatest, is but a continued fcene of rapine; and every quarter of the immense ocean, presents one uniform picture of hostility, violence and invasion.

In this unceasing conflict, occasioned by the voracity of the different kinds of sishes, the smaller tribes must have, long since, sallen victims to the avidity of the larger, had not Nature skilfully proportioned the means of their escape, their numbers, and their productive powers, to the extent and variety of the dangers, to which they are continually exposed. To supply the constant waste, occasioned by their destruction in the unequal combat, they are not only more numerous and prolific, than the larger kinds; but, by a happy instinct are directed to seek for food and protection near the shore; where, from the shallowness of the water, their destroyers are unable to pursue them. These, yeilding

to the strong impulse of hunger, become plunderers in their turn, and revenge the injuries committed on their kind, by destroying the spawn of the larger sishes, which they find floating upon the surface of the water *. Even there, however, they often meet with that violence which their own hostility merits; for the oyster, the sallop, and the muscle, lie in ambush at the bottom, with their shells open; and, whatever little sish inadvertently comes into contact, they instantly close their shells upon it, and, at their leisure, devour it in the concealed mansions of a prison, from which there is no possibility of escape.

In what manner digestion is carried on, to such an amazing extent and rapidity, in the stomachs of sishes, no inquiries of naturalists have yet been able to ascertain. If we were to judge from their substance or heat, we would be led to conclude, that the digestive powers of these animals are seeble and impersect; whereas, they appear so far to exceed every thing that can be effected, either by trituration, the operation of heat, or of a dissolving sluid, that a celebrated physician, after various experiments, has been of opinion, that none of these causes is equal to the effect; and that the power of digestion, in the cold maw of sishes, is so great, as to overturn those systems, that have attempted to account for it on these principles.

The powers of affimulation in fishes, seem to increase with the quantity of food with which they are supplied. A pike sparingly fed, can be habituated to subsist on very little nourishment; if fully supplied, it acquires the power of devouring an hundred roaches in three days. The digesting power of fishes, is as extraordinary, as their

^{*} Goldsmith, Nat. Hist. vol. vi. + Dr. Hunter.

their appetite is voracious. The cod and sturgeon will not only swallow, but dissolve crabs, muscles, lobsters, and every kind of shell sish, whose coverings are much harder than the coats of their stomachs. On dissection, the maw is found to be a fost bag silled with shells; the calcareous earth of which, undoubtedly, supplies the animal with nourishment. From this fact it would appear, that sishes are supplied with a powerful solvent, hitherto undescribed, which enables them to convert the substances they swallow into a sluid, sitted for their peculiar support.

Some exceptions to the extraordinary voracity of fishes, are produced by naturalists, which, if fully examined, will, perhaps, appear more apparent, than real. Some are faid to subsist on pure water alone; an affertion, which is supposed to be proved by numberless instances of their subsisting, for several months, in ponds constructed of hewn stone, where they had been supplied with no food *. It is alleged too, that those, which are carried from a distance to the markets of London and Paris in perforated vessels, must subsist upon water alone.

The element of water, however, is feldom found pure and unmixed. The very epithets of falt, bitter and fweet, imply a composition and mixture, perceptible even to the taste. The particles of the earth upon which it runs, necessarily enter into it, and vitiate its purity. These substances, together with myriads of animalcules with which it teems, may supply, for a while, a scanty subsistence, and support the life of the most voracious animals.

In

In the mean time, it will readily be allowed, that afhes, though for ever hungry and prowling, can endure the want of food for a long time. In them, habits feem to be formed by the circumstances in which they are placed. Want produces abilinence; from abundance they learn voracity. A pike, one of the most gluttonous of fishes, will live, and even thrive, in a pond where there is none but itself; and the gold and filver fishes, which we confine in glass vases, subsist, frequently for years, without any visible support but water. Rondeletius mentions one that was kept at his house, in this manner, for three years, which grew to fuch a fize, that the vafe could scarcely contain it, nor could it be brought out at the same passage by which it was introduced into the veffel *. It would appear, therefore, that, in certain fituations, fishes are as remarkable for abstinence, as, in others, they are distinguished for voracity; and that Nature, in compassion to the want which they must often fuffer, has indulged them with a power of accommodating their appetite to fearcity of food, as well as to abundance.

SECTION IV.

Of the Generation, Fecundity, &c. of Fishes.

The grand division of sishes into the cetaceous, cartilaginous, and spinous, was formed by Aristotle, according

to

Hoc modo fuavissima mea conjux, tres annos, piscem domi aluit; sicque educatus, in eam corporis molem accrevit, ut tandem nec vas cura capere, nec qua ingressus sucrat, eâdem exire posset. Rondelet.

to the three different modes of their generation. Among those of the first order, secundation is accomplished within the body of the semale, by means of a penis intrans, as in terrestrial animals. They are all viviparous; that is, the semale, after having been secundified by the male, and after a certain period of gentation, produces a living and persect animal.

The cartilaginous are also all, except some species of the flurgeon, viviparous; and among this order too, fecundation feems to be performed within the body of the female, who conceives two or more large eggs, distinctly containing red and white, like those of birds. In these eggs the fetus is formed; and, by the white of them, it is fed, while it is hatched within the body of the mother, without being excluded in the egg flate, as is the case in birds. In this state, the eggs have been found in the belly of a dogfish, with the young completely formed, and of a very confiderable fize *. The number of these eggs lodged at one time in the uterus, is various, according to the species to which the animal belongs. One naturalist + observed in the belly of t a dogfish, fix that were arrived at their full fize, befides many others, in which the fetus was fcarcely formed; another faw, in the cornua uteri of a torpedo, fix eggs on one fide, and eight upon the other.

The mode of generation that obtains among the viviparous or spinous fishes, is, from their situation and manner of life, involved in great obscurity. It is generally
supposed, that their eggs are not secundified, till after
their exclusion from the matrix of the semale. They
seem, indeed, amidst their evolutions in the deep, to copulate:

^{*} Raius apud Willough. Lib. I. cap. ix. † Rondeletius.

Francisc. Redi. Galei Spinacis.

pulate; but, as the male is furnished with no external organs of generation, his junction with the female is only to emit his impregnating milt upon the eggs, as they fall from her body. For this purpose, it is said, he pursues them along the stream, carefully impregnating them, one after another.

These facts, however, are controverted by Linneus, who maintains, that no fecundation can take place, except within the body of the semale *, although the generation of frogs and lizards has always been regarded as an example of the contrary. In support of his hypothesis, he afferts that he observed, at spawning season, every male pike surrounded by several semales; and that, as soon as the milt was ejected by him, it was immediately swallowed by the semales; a procedure which he had occasion to notice in several other kinds of sith.

The experiments that have lately been made at Berline by a skilful naturalist; feem totally to overthrow this doctrine of Linneus. It has been found, that of both salmon and trout, the roes, artiscially extracted from the body of the semale, were capable of being secundified by an admixture with the milt of the male. Hence it is probable, that, though both the male and the semale concur in the great work of impregnation, yet the act is performed without the body of the latter; and that the spawner ejects her eggs, while the milter sprinkles them with sperm.

The science of ichthyology is still in its infancy. Supported by sew accurate observations, and still sewer experiments, it has hitherto been regarded as a field for Vol. III.

D theory

Vide Systema Naturæ ad locum,
 M. Jacobi,

theory, speculation, and conjecture. Hence we are no fooner free of one difficulty, than we are met by another equally perplexing and infurmountable. It has long been a question, among philosophers, whether fishes are produced by fpontaneous generation *. That they are, an ancient naturalist pretends to have proved by his own experiments and observations +. There is, says he, in the vicinity of Pifa, and furrounded, on every fide, by hills, a place, into which the waters of no fountain, stream, or lake, ever flowed; a place, which no moisture, except the rain from the clouds, ever watered. A few days before my arrival at this spot, a copious shower of rain had overflowed it with water, and nature herfelf had fupplied it with fishes, among which vast numbers of carp were observed. The same thing, he adds, is feen in many pools in France, into which no flock, or original, of the many various fishes they contain, was ever placed.

Another philosopher has adopted the same opinion concerning the spontaneous production of sish; because, in ponds newly dug out, sishes have been observed during the sirst year; and, in succeeding ones, many other different kinds, although none had been originally transported thither. It is a phenomenon in Nature, for which it is indeed difficult to account, but notwithstanding an incontrovertible sact, that in stagnating pools occasioned by the rain in Bombay, which have no communication with any river, or the sea, sishes are generated, of which many persons have eaten, and which,

upon

^{*} Willoughby, lib. I. cap. ix.

[†] Rondeletius de Piscib. lib. I. cap. iv.

Albertus magnus apud Gefnerum.

upon the drying of these ponds, die, and are corrupted *. But, however dissicult these appearances may be to explain, or in whatever manner the fishes were introduced into these ponds, we must conclude, that they were originally produced from the eggs of animals of their own kind; because the idea of spontaneous generation is repugnant to every maxim of sound philosophy †.

Fishes, in general, are male and female; the former possessing the milt, and the latter the roe. Some individuals of the cod and sturgeon, are faid, indeed, to contain both. The spawn of the greater number of fishes is deposited in the fand or gravel; and, in that state, it is probable, that the roe and milt are mixed together. Summer is the most common time for the spawning of fishes; because, at that season, the water is tepisied by the beams of the fun, and therefore better fitted for quickening the eggs into life. It is probably for this reason, that the herrings frequent the shores of Britain, at the spawning season; there not being, in the unfathomable depths of the ocean, a fufficient quantity of heat to hatch their eggs. When they have deposited their burdens, they return to their former stations, and leave their infant progeny to shift for themselves.

The spawn of different fishes continues in the state of eggs, sometimes for a longer, and sometimes a shorter period. In general, however, this period is proportioned to the size of the annual. In the salmon kind, the young animal continues in the form of an egg, from the month of December till April; the young carp con-

D 2 tinues

^{*} Vide Lind. on hot climates, Part II p 52 .

[†] Linnæus, omne animal ex ovo.

tinues not, in that state, for above three weeks; and the gold sish from China, is produced in a still shorter period *.

When excluded from the egg, the young fishes all, at first, escape by their minuteness and agility. They perform all their motions with much greater celerity and ease, than grown and bulky sishes; and can make their way into shallow water, where they cannot easily be purfued. But, with all these advantages, not one, perhaps, of a thousand, survives the dangers of its youth. There is no instance, among the spinous sishes, of any thing resembling parental affection: They all abandon their eggs to be hatched by the heat of the season; and, if they ever return to the spot where their young have begun the career of life, the male and semale that gave them birth, forgetting all parental relation, become enemies as formidable as the most rapacious plunderers of the deep.

Though the fishes fall in millions by the rapacity of one another, yet they have other enemies than the animals of their own kind. Many of the quadrupeds frequent the shores, rivers, and lakes, where fish is almost their only food; and we have already had occasion to remark, that a great proportion of the fowls reside comfantly on the sea, where they either seed upon spawn, or become the merciles invaders of the small sishes. Thousands too of the human race, wander, in a savage state, around the lakes and rivers, whence they derive a considerable portion of their sustenance. And, among those

^{*} Goldsmith's Nat. Hist. Vol. VI.

[†] The collis boviæ is the only exception to this observation; it is said to fit upon its eggs at the bottom of the water.

those nations, whom arts and agriculture have rendered less dependent on this precarious support, superstition has come in the place of want, and given a new edge to their avidity for this species of sood. From the invasions, therefore, of terrestrial animals, and from their own mutual rapacity, the annual consumption of sishes is constant and immense; but the muniscent Author of Nature has made a kind provision for his creatures, by the amazing fertility which he has conferred on this class of beings.

The fecundity of fish in general far surpasses that of any other animals; in some, it exceeds belief, for there are individuals among this class of the animal kingdom capable of producing, in one year, a greater number of their species than all the inhabitants of Great Britain. Nine millions of ova have been found in the spawn of a single cod *; one million, three hundred, and sisty-seven thousand, four hundred, have been taken from the belly of a slounder: The mackerel, carp, tench, and a variety of other species, are said to possess a degree of fertility but little inserior †.

Such an aftonishing progeny, were it allowed to arrive at maturity, and to add the whole of its number to the rest of the samily, would soon overstoon. Nature; and even the ocean itself would not be able to contain, much less to provide for the half of its inhabitants. Of the ova spawned by these different fish, however, probably not one in an hundred ever becomes a full-grown animal: They are devoured by the lesser fry that frequent the shores, by aquatic birds near the margin, and by

^{*} Dr. Lewenloch.

Wide Farmer's calculations, Philosophical Transactions, 1767.

the large fish in the deep water. Such as still survive; are sufficient for supplying the ocean with inhabitants; and of these, notwithstanding their own rapacity, and that of the aquatic sowls and terrestrial animals, enough is lest to relieve the wants of a great portion of the human race. Thus, two important purposes are answered in the economy of Nature by the extraordinary secundity of sishes: it preserves the species amidst numberless enemies, and serves to surnish the rest with a sustenance adapted to their nature *.

Among terrestrial animals, there are various degrees of fertility, according to their chance of destruction, from the want of courage, bulk, or strength: The largest are always least productive, and the smaller are more prolific, in proportion as the dangers increase, to which they are exposed +. The same observation may be extended to the inhabitants of the ocean: Among these, as the cetaceous fish refemble quadrupeds in their habits and conformation, fo they are distinguished by a similar degree of sterility. All the whales, and even the cartilaginous fishes, are not, perhaps, superior to terrestrial animals in their powers of production. Among the hungry and heedless inhabitants of the sea, they are distinguished by finer organs and higher fenfations. Their fize and courage place them, in a great measure, beyond the reach of danger; they nurse their young with tenderness and affiduity, and they protect them from injury with an obflinate intrepidity little inferior to that of the lion or the eagle. Hence the number of their young is few, proportioned to the dangers to which they are exposed: Amidf

^{*} Goldsmith's Nat. Hist. vol. VI.

[†] Buffon's Hist. generale et particulaire, tome 16.

midst that security which they enjoy, and the ease with which they can repel the hostility of the inimical tribes, they run but little hazard of being destroyed by their rapacity.

SECTION V.

Of the Growth, Longevity, and Dietical Uses of Fish.

As the dangers to which the progeny of all the spinous fishes, while in the form of ova, and in their nascent flate, are innumerable, and furrounding them on every fide, Nature has happily ordered, that they should remain but a short time in that defenceless condition. The period at which the different species arrive at their appointed fize, is not exactly afcertained; but the generality of naturalists agree in reckoning it extremely short *. Aristotle, and Pliny who copied him, seem both to have committed a mistake, when they assigned only two years for the life of the tunny +, a fish which approaches to to the fize of a whale. This species had various different names assigned it, till it arrived at its fifth year, when it received the appellation of a whale; a circumstance from which we must conclude, that the ancients in general entertained a very different idea of its longevity ‡. Of a fimilar

^{*} Rondeletius, de piscibus.

[†] Vita Thynnis longissima biennio. Plinii Hist. Anim.

[‡] During the first year, they were called Scordyla; the fecond, Palamides; the third, Thynnis; the fourth, Oreyni; and the fifth, Cete.

fimilar error, many of the moderns are guilty in afferting, that the common falmon arrives at its full growth in a year. Our most experienced fishers distinguish them also by different names, till they arrive at their fixth year, when they are supposed to have reached their full fize: Those of the first year they call smelts; of the fecond, fprods; of the third, morts; of the fourth, forktails; of the fifth, half-fith; and of the fixth, falmons *. It is probable, however, notwithstanding the immense fize at which many of them arrive, that fishes, in general, are of a quick growth, and that the period of their adolescence is small, in proportion to that of their lives. There is another peculiarity attending the growth of fish; and that is, the cetaceous kinds, which are by far the largest, inhabit chiefly the colder regions of the ocean, adjoining the poles. The reverse of this is the cafe with terrestial animals, who always diminish in fize as they recede from the heat; and by far the largest of them, the elephant, the camel, and rhinoceros, are only found in the warm or intratropical latitudes. It is probable, indeed, that the whales might prefer the more temperate regions of the ocean, and might even acquire a large fize there; were they not compelled to avoid them by the frequent diffurbance given by ships, and to seek that tranquillity, of which they are fond, in the unhofpital climes of the north +.

Hitherto, we have been examining fishes with regard to their external conformation, and have found them inferior to terrestrial animals in their organization, and in the number of their enjoyments. We are now to contemplate

£ 4 2. .

^{*} Vide Willoughby, lib. i. cap. 16.

⁺ Idem ibidem.

piate them in a more favourable point of view, and that is, in the extraordinary period of time during which they possess the humble existence which Nature has assigned them. Their longevity is far superior to that of other animals *. We have already seen what ample provision is made for supplying them with food, by multiplying the inhabitants of the sea: They are, therefore, in little danger of perishing from want; and there is reason to believe, that they are, in a great measure, exempted from diseases.

Most of the disorders incident to mankind arise from the changes and alterations incident to the atmosphere; but sithes reside in an element but little subject to change; theirs is an uniform existence; their movements are without effort, and their life without labour. Their bones also, which are united by cartilages, admit of indefinite extension; and the different sizes of animals of the same kind among sishes is very various. They still keep growing; their bodies, instead of suffering the rigidity of age, which is the cause of natural decay in land animals, still continue increasing with fresh supplies; and as the body grows, the conduits of life furnish their stores in greater abundance. How long a sish, that seems to have scarce any bounds put to its growth, continues to live, is not ascertained †.

Some species of the German carp have been known to live an hundred years +; those species, however, with which we are best acquainted, seem not to enjoy so great a degree of longevity. The salmon, we have remarked, takes only six years to reach its full size; if we

Vol. II. E allow

e Gesner de piscibus.

[†] Vide Bacon's werks.

allow four or five times that space for the period of its life, it will not exceed thirty years.

Different methods have been devised for ascertaining the age of fithes, fome of which will, perhaps, pretty accurately determine this matter. The ingenious Mr. Hiddostroam, a Swede, has attempted to compute their ages by the number of concentric circles observed in a transverse section of the vertebræ of the back; and it has been found, that each circle, like that in the fection of a tree, corresponds to a year of the animal's life. In confirmation of this fact, experiments have been made on different individuals of various fizes, but of the same age, whose concentric circles have always been of an equal number; whereas a young fish differs from an older one of the fame species, by having a smaller number of these circles. By this method of computation, feveral fish were found from fifteen to twenty years, but none gave indications of a greater age.

Another method of computing the age of fishes, practifed by M. de Buffon, is by numbering the concentric circles upon their feales *; but as this requires a more minute examination, it is, of consequence, liable to greater uncertainty. On examining the scale of a fish through a microscope, it exhibits a number of circles, one within another, resembling those in the vertebiæ of the back, and, like them, every circle represents a year of the fish's life. A scale of a carp, thus examined, announced the animal to be no less than an hundred years; a longevity less incredible, because confirmed by the testimony of several different authors †, some of whom have afferted, that this fish lives twice that period.

The

Vide Hist. gener. et part.

The dietical uses of fishes are to us the most important article of their history, a part that is happily free from that uncertainty and darkness in which many other circumstances relating to their manners and economy are still involved. All fish whatever, and particularly those without the tropics, are capable of being converted into wholesome food. Every European fish, while in seafon, is nutritive; the various methods of preparing and dressing them are detailed by the Authors to whom that province belongs. Such disquisitions constitute the history of an art, but they are not the objects of science.

Fishes, in general, when out of feason, are unwholefome, and even pernicious; and this is more especially the case with the oily kinds; such as the herring, the mackerel, the eel and the falmon. So little is this fact attended to, that the eating of the latter of these fishes, at an improper feafon, has been the immediate cause of an epidemic fever. Some of the fishes that frequent the shores of the West-Indian islands, are said to be poisonous: that they are naturally fo, however, may be justly questioned; though there can be no doubt of their becoming pernicious in certain circumstances. If they feed upon copper-banks, their food may contain fo great a quantity of the poison combined with that metal, as may render their flesh noxious; and the fame consequence will enfue if they devour the feeds of fome poisonous plants that grow in these parts of the globe.

The improvement of the arts tends greatly to extend the dominion of man over the inferior animals; the art of navigation, in particular, may be faid to have completed his conquest of the ocean, and brought a vast acments, by the capture of fo many animals whom Nature feemed to have placed beyond the reach of his power. A fleet of fishing vessels, manned by a few thousand European failors, is capable of taking a greater number of fish in a feason, than perhaps could be done by all the favages on the continent of America. But the superstitious regulations of the Romish Church has tended more than any other circumstance, to enhance the value, and increase the quantity of this species of food. To supply the imaginary abstinence of the devout or supply the imaginary abstinence of the devout or supply the imaginary abstinence of the devout or supply, ponds have been dug, and sish, like land animals, rendered domestic.

In the condruction of these ponds, it may be proper to observe, that, if intended for breeding, they ought to be of various depths, from fix feet to fix inches; for few fish will spawn in water of any considerable depth. Such pieces of water ought also to be partly filled with aquatic plants; because these afford shelter and nourishment to different kinds of intees, that contribute to feed the fishes: For a fimilar reason, fish pends ought to be furrounded with trees, that the infects frequenting them may afford an additional fumply for their fullonance. In arranging the fillies in these receptacles, attention ought to be paid to the different species that are introduced into the same pond. Carp and tench agree well together; but they frequently do not thrive when mixed with any other species. The perch is almost the only fish that can be fafely entrusted in a sheet of water frequented by the pike.

In populous countries, like China or Holland, where every article of food is in request, and every spot of ground is turned to account, great attention is paid to the structure and management of sish ponds: There experience has taught men to ascertain the quantity of every kind of sish, which as y given space of water can support. Ninety brace of carp, and forty of tench, may be supported in an acre of water; a greater number would languish and die. In transporting sish from rivers into ponds, it has been found that the young thrive much better after the operation, than the old; for at that period of life, they possess a power of more easily accommodating themselves to any change in their situation.

In some parts of Germany, where the demessication of fish is practiced, the ponds are constructed in a suite, one adjoining another; and, by means of a communication between them, they can empty the water of one pond, together with the fish, into another. After this is accomplished, the empty one is frequently plowed, and sown with barley: When the grain is in the ear, the water, with its inhabitants, are again admitted to feed upon the grain, and are thus more expeditionsly sattened than by any other management. In all fish ponds, the milters are the preserable fish, for they become much statter than the spanners.

Fish of discrent species, as well as terrestrial animals, are found capable of producing an offspring; but their cross breeding is a very obscure article of their history. Few species have hitherto been domesticated, and on these, but a very small number of well-attested experiments are known. A cross breed is faid to be obtained between the carp and the bream, and between the carp and tench:

That between the carp and tench was produced by mixing the milt of the former with the spawn of the latter; the offspring, which was examined by Mr. Pennant, bore a greater resemblance to the male than the semale parent *.

It is remarkable, that so few species of this class of the animal kingdom have been translated from their primitive haunts, and made subservient to man by domestication. Only three kinds have been transported from soriegn parts into Britain; the carp, the tench, and the goldsish. Double that number are domesticated with success upon the continent; but even this, in all probability, is but sew to what might easily be rendered subservient to the same purpose, by the industry of man. There can at least be entertained no doubt that those domesticated upon the continent would thrive equally in Britain.

The caprinus corufius is found indigenous in many of the rivers in England, where it is called the rudd. It is reckoned a superior fish to the carp in many respects, yet it has never been domesticated in its native country, though reared with great advantage in the fish ponds of Sweden. The cobetus fossilis is a fish unknown in this country; but is domesticated, with profit, in the ponds at Stockholm, and might, with equal propriety, be tranflated into those of Scotland. In the same manner, the cobetus barbatula, which was introduced into Sweden by Frederick I. might be made an inhabitant of our artificial ponds. The falmo tymalis of Linnaus, known in England by the name of graylin, is one of the best fishes both for fport and the table, and might eafily be transported here from the streams of Derbyshire, its native residence.

^{*} Fide Brit. Zool. Vol. III. class iv. p. 356.

refidence. Both eels and charr might be rendered useful pond-fish; as neither are too delicate for transportation; and, from experiments already made, the certainty of their thriving is fully established.

If the translation of fish has not been often attempted, it certainly is not because they are incapable of sustaining various degrees of heat, and of living in different climates. The necessity of procuring a supply of food; of seeking a safe retreat for propagating their species, or a temperature of that element in which they live, suited to their constitutions, compels the fishes, as remarkably as terrestrial animals, to make extensive migrations from one part of the sea to another. With regard, however, to this curious subject, we have but sew facts upon which we can depend.

The fish, like land animals, are either solitary or gregarious: Of the former kind, trout, salmon, pike, &c. the migrations are probably in quest of a proper place to deposite their spawn. The salmon, for this purpose, leave the sea, and mount the rivers in the beginning of winter, where they dig in the gravel, deposite their burden, and again return. The trout likewise ascends near the source of the rivutets at the season, when they enter the smaller branches that run into the main stream, for the purpose of spawning. It is then they are often seen in small rivulets upon the high grounds, in water so shallow as scarcely to cover their bodies.

Of the gregarious filhes that frequent fresh water, we know but little concerning their migrations. It is probable, that the perch and the minnow are stationary, and that they retire only to the margin of the river to deposit their spawn. The fishes, most remarkably gregarious,

are inhabitants of the fea; fuch as the cod, ling, hetring, pilchard, and sparling. The migrations of the herring are well known. In the months of June and July, it islues in vast shoals from the unexplored regions of the northern ocean; surrounds the British isles, and enters the bays and arms of the sea. These sithes are known to take up their residence in some particular loch or creek for eight or ten years, and then to resort to another for a similar period. The tythe, the calsish, the cod, and whiting, have all their seasons of migration. The last, in particular, are probably forced upon those immense journeys from the coasts of America to these of Spain and Africa, to avoid the pursuit of the cod, and other rapacious invaders.

From these observations, it would appear, that fish of the same species are capable of living in very different quarters of the globe, and of enduring various degrees of heat and cold; a circumstance which opens a vast field for the enterprise and ingenuity of man in transporting them, and rendering them subservient to the purposes of domestication. It is impossible to determine to what length this operation may be carried, or to ascertain how great an accession might thus be made to the sustenance of the human race.

CHAPTER II:

SECTION I.

Generic Characters of the Cetaceous Fishes.

The cetaceous fishes constitute the sirst order, according to the Aristotelian division; and have obtained from him that name, on account of their enormous size, and their producing a perfect animal from semen, and not from out.*. These animals, from the more perfect conformation of their organs, as well as their superior bulk, are sully entitled to the first rank among the inhabitants of the ocean. Among the terrestrial animals, we have seen some endowed with higher powers, and more various instincts, and there y challenging a superior rank among the kindred tribes. The same distinction obtains among the aquatic orders, where the whale, possessing the organization and many of the habits of quadrupeds, claims the first station among his companions of the deep †.

This order contains the whales of all the various kinds, who want teeth; the cachalot, with teeth in the lower Vol. III.

^{*} Willoughby, lib. ii. p. 26. † Goldfmith's Nat. Hift. vol. VI.

jaw; the dolphin, with teeth in both jaws; the porpeffs: and the grampus. Such are the marks by which the different tribes of this divition are diffinguished from each other; but the most striking and obvious character by which they are at once discriminated from all other animals, is their immense fize. The testimony of all ancient writers concurs in afcribing to them a bulk far furpassing the largest animal known upon the globe. In those books which Juba wrote to the fon of Augustus Casar concerning Arabia, whales of fix hundred feet long, and three broad, are faid to have entered a river of that country *. Pliny himfelf vouches the extraordinary fize of two hundred and forty feet in length, by eighteen in breadth; and Nearchus declares, that he faw on one of the islands opposite to the month of the Euphrates, a whale of an hundred and fifty cubits, cast out upon the thore. Arrian +, Strabo +, and Diodorus, mention nations of favages bordering on the Indian Ocean, which built their houses of the bones of whales cast out by the fea, using their jaws for door cases; and we find from Trobifber, that the same method of architecture was formerly practifed by the inhabitants of Greenland &.

Many later writers give a fimilar account of the magnitude of whales. Even within the arctic circle, they were formerly of a prodigious fize, when their capture was left frequent, and the fish were allowed to grow. We know, by the accounts of travellers, that within the torrid zone, where they are unmolefted, they are ftill feen one hundred and fixty feet in length ||. Those at present taken in the Greenland seas, seldom exceed eighty or nine-

ty

^{*} Pliny, lib. xxxii. cap. 1. † Liber viii. ‡ Liber xv.

[§] Vide Second voyage, p. 18; | Adamson's voyages, p. 174.

ty feet in length, and about twenty in breadth: Even thefe, however, must appear enormous to the spectator, and, when first beheld rising from the deep, must have struck him with aftonishment and terror. Those who frequent the northern ocean, to whom the frequency of the fight has rendered it less tremenduous, must enjoy a magnificent spectacle in beholding such an enormous animated mass tumbling amidst the waves, and darting through the water with incredible velocity! Such wonders does Nature present in an element where her operations are but partially known; perhaps the has fill greater wonders concealed in the deep, which we have had no opportunities of exploring. The whales are obliged to shew themselves upon the surface, in order to take breath : but who knows the fize of those animals that are fitted to remain for ever under water; and that have been increafing in magnitude for centuries *.

Though we now fee no whales two hundred, and two hundred and fifty feet, which we have good reasen to believe was the case two centuries ago, still the size of this order of beings is sufficient to distinguish them from all the rest of the animal kingdom. It is probable, indeed, from skeletons that have been found at different times, that there once existed terrestrial animals of much greater bulk than are to be found at present. Creatures of such an immense fize must have required a proportionable extent of ground for subsistence; and by being rivals with the human race for large territory, they must have been destroyed in the contest. A similar cause has diminished

^{*} The fea-ferpent, or kraken, is faid to be feveral miles in circumference, and to appear like a fmall ifland. To believe all fuch relations would be credulity; but to deny their possibility, would be prefumption. Goldfmith's Nat. Hist. vol. VI.

diminished both the number and the fize of the cetaceous tribes: Man had no fooner completed his fevereignty over the ocean by the invention of the compass, than these animals were so much thinned by continual capture or destruction, that they have never had time to arrive at that enormous bulk which the peaceable possession of the sea, in former ages, conferred *.

The external conformation of the filhes of this genus, no less than their fize, serves to characterise them among the other inhabitants of the deep. They are covered with a dark-coloured cinereous skin: they are moved, commonly against the wind, and with vast rapidity, by means of a horizontal tail, aided by three fins; two pectoral, and one back fin; but in some species, the last is wanting. The head is commonly extremely large, in proportion to the fize of the body, being in some equal to a third of the fize of the fish. The animal is directed to its prey by two small eyes, furnished with eyelids, and not superior in fize to those of an ox. As the cetaceous tribes all breathe by means of lungs, they have no branchise nor external apparatus for that purpose +. In the middle of the head, there is one, fometimes two orifices, through which they fpout water to a vaft height and with a great noife. With these orifices raifed above the furface, the whales fleep and breathe, genthy moving their fins, to keep them poifed upon the fummit of the water t. When immerfed below the furface, or while devouring their food, water unavoidably rushes into the throat and lungs, and is, in this manner, ejected every time they rife for a fupply of air. If the animal be wounded, it spouts the air and water with a violence fufficient

^{*} Vide British Zoology, class iv. gen. 1. † Aristot. Hist. Animal.

Willough. lib. ii. cap. I.

fufficient to overfet a fhip; and the noise it occasions, is heard like the discharge of cannon, at the distance of some miles.

Animals of fuch enormous firength and magnitude, we might imagine, would fprend terror and devastation all around them, and make an indiferiminate flaughter of the inferior tribes. No creature, however, is less voracious than the common whale: Almost no animal subflance is ever found in its stomach; it feeds, as some allege, upon different infects that float on the surface; according to others, upon the medusa or sea-bitubber. Its food, we are certain, must be extremely minute, for the capacity of its throat does not exceed four inches, a fize beyond all proportion, smaller than that of other sish.

The small quantity of food that suffices the whale, may justly surprise us, when we consider their size, and the numbers of these animals that often herd together. Had their voracity been proportioned to their bulk and numbers, the ocean itself would hardly have assorded a sufficient supply. The insects upon which they feed, are black, and about the size of a bean: They are of a round form, like snails in their shells, and are seen sloating in clusters together upon the waves. These, the whale such in great numbers, and bruises them with the barbs or pipes with which its mouth is internally covered †. Nourished with this food, it becomes the fattest of all animals, whether terrestrial or aquatic.

To a flender appetite, the whale adds peaceable and harmless manners: it pursues no other fish, but leads an easy and indolent life on the bosom of the waves, and is inosfensive,

inossensive, proportioned to its ability to do mischies. Among land-animals, we have had occasion to observe, that sovereignty does not always follow strength or fize *: The elephant and camel sly before the tyger and the lion; while the eagle possesses a decided superiority over the vulture and the offsich. The same law obtains among the inhabitants of the ocean; where the whale, if he holds the sceptre, holds it by a precarious tenure, for it may easily be wrested from him by his subjects. There is a strong analogy between his manners and those of the elephant: Both are the strongest and largest animals in their respective elements; neither offers injury; and each is terrible when provoked to resentment.

But these peaceable and innoxious habits do not equally belong to the whole of the cetaceous order: The finish differs from the great whale in this respect; it subfiles chiefly upon herring, and is often from driving vaft foods of them before it. Those of the cachalot tribe are fill more voracious, and commit greater depredations; They are furnished with teeth both in the upper and lower jaw; their throat, though inferior in capacity to that of the cartilaginous fishes, is much larger than in the common whale; and these powers of deglutition were probably conferred to gratify the cravings of an appetite proportionably more voracious. It is faid, that the fiftes of this genus are polleffed of a courage, that often proves fatal even to the animals of their own order: they purfue and terrify the purpelles to fuch a degree, that they often drive them ashore +.

The common whale, whatever honours vulgar prejudices may have conferred upon it, has no pretentions to

^{*} Vide Buffon, Hift, gen. et part. † Vide Lin. Syftem. Natura.

the fovereignty of the ocean: On the contrary, as it is a peaceable and inoffensive animal, it has many enemies disposed to take advantage of its disposition, and inaptitude for combat *. There is a fmall animal of the testaceous kind, called the whale-louse, that sticks to its body, as we see shells stick to the foul bottom of a ship. This creature infinuates itself chiefly under the fins; and, in defiance of all the efforts of the whale, it still keeps its hold, and lives upon the fat, which it is provided with instruments to extract. The filhermen, however, often witness the encounters of the whale with a much more terrible enemy. At the fight of the fword-fish, this largest of animals is feen agitated in an extraordinary manner, and leaping from the water as if with fear. Wherever it appears, the whale perceives it at a distance, and flies from it in the opposite direction. The whale has no infrument of defence except the tail: with that it endeavours to strike the enemy; and a fingle blow taking place, would effectually deftroy the adversary: But the fword-fish is as active as the other is strong, and easily avoids the firoke; then bounding into the air, it falls upon its subjacent enemy, and endeavours, not to pierce with its pointed beak, but to cut with its edges. The fea all about is feen dyed with blood, proceeding from the wounds of the whale; while the enormous animal vainly endeavours to reach its invader, and firikes with its tail against the surface of the water, making a report at each blow louder than the noise of a cannon +.

A still more fatal enemy of the whale, is an animal of its own order, called by the sistermen of New England the killer. Of ferocious habits, and furnished with strong sharp

[&]quot; Geldsmith.

Anderson's Dictionary of Commerce.

sharp teeth, these animals when they surround a whale seldom allow it to come off with life. They tear and mangle its sleth on all sides, till satigued with sighting, and overcome with wounds, it salls a prey at last to their surry; and after it expires, the tongue is extracted, the only part which they devour.

By the contlant hostilities of these various animals, the race of whales has probably been gradually diminishing in number for several ages *. From the largeness of their fize they cannot easily be concealed from their destroyers; and as they are distinguished by sterility among the sinny tribes, their destruction cannot soon be repaired: But of all the causes of the waste and diminution of this order of sishes, the interserence of man has operated by far the most powerfully. His hostilities has been incomparably more satal than those of all the rest of their enemies; and a greater number is probably destroyed in a season by the ingenuity of the sishermen, than is devoured by the rapacious animals in an age.

The inhospitable shores of Epitzhergen were found to be the great resort of the wholes; and for more than three centuries, notwithstanding the coldness of the climate, and the terrors of the icy sea, a great number of European ships have annually requested those dreary abodes, and at length thinned the number of their inhabitants.

The whale fishery was carried on, for the fake of the oil, long before the use of whale bone was discovered †. The substance which has obtained that name, adheres to the upper jaw; and is formed of thin parallel lamine, some of the longest four yards in length. Of these there

are

^{*} Vide Brit. Zoology, Class. iv. Genus. 1.

[†] Vide Anderson's Dict. vol. i. p. 442.

are about feven hundred in all: about two thirds of that number are of a length fit for use, the rest being too short*. The oil is extracted from different parts of the body; the tongue alone of some sish yielding from sive to six barrels.

As early as the beginning of the fourteenth century, the Bifeayneers were in possession of a very considerable trade to the coast of Greenland: They long enjoyed the profits of a lucrative trassic in train oil, and whale-bone, before the English attempted to obtain any share of that commerce. What probably first gave them an idea of the advantages to be reaped from it, was the accident of one of their ships bringing a cargo of whale-bone and train-oil from the bay of St Laurence, part of the burden of two large Bifeayan ships that had been wrecked there about the year sisteen hundred and ninety-four †.

A few years after that period, the town of Hull had the honour of first attempting that profitable branch of trade. At present it seems to be on the decline, the number of fish being greatly reduced by their constant capture for fuch a vast length of time. It is now said that the fishers, from a defect of whales, apply themselves to feal fishery, from which animals they also extract an oil, and turn the skins to good account. This trade however will not probably be of any long continuance, for these shy and timid creatures will foon be induced to quit those shores by being perpetually harrastied. We are informed too, that the natives of Greenland already begin to fuffer from the fearcity of feals in their feas. The fiesh of these Vol. III. G animals

British Zonlogy, ubi supra.

Hackluyt. Vol. iii. p. 194.

animals conflitutes their principal fublishence; and should they be at last extirpated, or desert the coast, that miferable people would be in danger of perishing through want*.

Before the year 1598, the whale feems never to have been taken on our coasts but when it was accidentally driven a-shore +. It was then deemed a royal fish, and the king and queen divided the spoil between them; the king afferting his right to the head, and her majefty by prerogative entitled to the tail t. A total revolution in the fashion of eatables, and the great quantity of these fish that are now imported, has rendered this prerogative of royalty of less importance, and even ludicrous: formerly however the whale as well as the porpoife, and dolphin, was probably a duh ferved at the royal board; and from its magnitude it must have held a very respectable flation there. Such dainties continued in vogue so later as the reign of Henry VIII.; for, in a household book of that prince &, it is ordered, that if a porpoise should be too big for a horse load, allowance should be made to the purveyor. Even in the reign of Queen Elizabeth, we find directions for the dreffing and ferving up of the dolphin with porpoife fauce; a composition of vinegar, crumbs of bread; and fugar ||

The flesh of the whale has always made a part of the food of fome favage nations. The natives of Greenland as well as the barbarous tribes that inhabit the vicinity of the fouth pole, eat the flesh prepared in various ways, and drink the oil, which is with them a first rate delicacy. The finding of a dead whale is an adventure confidence.

among

^{*} British Zoology, Class iv. Genus 1.

[†] British Zoology, idem ibidem.

[‡] Blackston, Comm. i. c. 4. § Archæolegia, Vol. iii-| Gaij. Opuscula-116.

emong the most fortunate circumstances of their wretched lives. They make their abode beside it; and seldom remove till they have lest nothing but the bones*. In the days of Willoughby, the eating whale was growing into disuse in England; and at present the Dutch sailors, as well as our own, will not taste it except in cases of urgent necessity: it is said, however, that the French seamen frequently dress and use it as their ordinary food at sea. The wretched inhabitants of the island of Feroe, who live one half of the year on salted gulls, are also, we are told, very fond of salted whales slesh: the fat of the head, after being well seasoned, they hang up in the chimney, and eat like bacon‡.

The internal structure of the whale we have already remarked, refembles almost in every respect that of quadrupeds: Like them they possess lungs, a bilocular heart, a diaphragm and urinary bladder. The precife shape and fituation of the viscera of these animals is indeed far from being fo exactly afcertained as might have been expected, from the trequent opportunity there is afforded of examining them. In those parts where they are caught in greatest abundance, the sailors are not very curious in inquiring into the structure of the parts; and few anatomists care to undertake a task, where the operator, instead of separating with a lancet must cut his way with an ax ||. It is therefore not yet known whether the whale has not one of its bowels entirely adapted for the reception of air, in order to supply it with that sluid, when it is obliged to continue longer than usual below

G 2

the

[&]quot; Goldfmith's Nat. Hift.

[†] Duram babent carnem setacea omnia; quamobrem liffis fale conditis fore in cibis utimur. Liber. ii. cap. 1.

a Jacobson's Hist. of Feroe.

^{||} Geldfmith's Nat. Hift.

the furface of the water. We are certain, that there is no fuch receptacle for air among the vifcera of terrestrial animals: The analogy however, is not here so strong as to justify us in denying the whale a particular provision of this nature, which the sailors affert that they have discovered, and which the necessities of the animal seem

to require.

As the cetaceous filtes refemble terrefirial animals in their conformation, to they are also distinguished by fimilar appetites and manners: Among them the act of copulation is faid to be performed more humane, and the female once in two years feels the access of defire. In the inferior tribes of fish we can differ hardly anything like pairing between the males and females, and have no vestiges of conjugal fidelity. The mutual attachment of the whales, however, exceeds whatever we are told of the constancy of birds.

Whenever a pair of whales are attacked by the fishers, they mutually assist, it is asserted, in the defence of each other; and when one is wounded the other still attends lending every aid in its power; and no motives of fear or self preservation can urge it to defert its associate. An instance is recorded * of one, which, after maintaining an obstinate consist in desence of its companion, that had been struck with a harpoon, and on seeing it expire under the wounds it had received, stretched itself upon the dying sish and yielded up its breath at the same instant.

The period of gestation among the cetaceous fishes is faid to be nine or ten months; the female commonly produces one, and never above two young. During the time of her pregnancy, and particularly at the birth of

her

iter offspring, she is uncommonly fat. The embryo, it is faid, when first perceptible, is about seventeen inches long; and the cub when excluded, is of a black colour, and ten feet in length. The two breasts of the whale are hid within the belly; but when she suckles her young, she can produce them at pleasure, when they are protuberant about two feet before her body. The teats resemble those of a cow; while the colour of the breasts, in some is white, in others speckled; in all, they are silled with a large quantity of milk resembling that of land animals.

From what has been faid concerning the procreation of whales, it appears, that these animals, in secundity, are far inserior to the rest of the inhabitants of the deep. Nothing, however, can exceed their care and tenderness for their offspring when produced. The semale whale carries her cub with her wherever she goes; and when pursued by the sishermen she keeps it supported between her sins. Even when wounded, she still class her young one; and when forced to plunge into the deep to avoid the strokes of her pursuers she carries it down along with her; but rises sooner than usual, to allow it time for respiration.

In defence of her young, the whale is faid to display a fierceness and courage far beyond what could be expected from an animal, so gentle and inossensive in its nature. IVuller + describes the conduct of a whale and her cub, when surrounded in an arm of the sea, in a manner which strongly displays the maternal tenderness of these animals. Being deserted by the tide, they were inclosed

on

^{*} Vide Goldsmith's Nat. Hist. Vol. 6.

⁺ Vide his Poem of the Summer islands.

on every fide; and the people from the shore observing that they had no means left of escape, poured down upon them in boats, armed with such weapons as the urgent occasion offered. After several ineffectual attempts to escape, the mother at last, by her superior strength, forced herself over the shallow into deep water: But though in safety herself, she could not bear the danger that awaited her young one; she therefore rushed in once more where the smaller animal was inclosed, and resolved, when she could not protect, at least to share its danger. Happily for the safety of these animals, the rapid inset of the tide soon cnabled both to escape from their enemies, though not till they had received many wounds, and tinged the sea all around with their blood.

It is, however, but for a short period that the young whales stand in need of this parental assistance: Their growth is fo remarkably rapid, that it may occasion some doubt concerning the accounts that are given of their extraordinary longevity. The cubs continue at the breaft, of the mother only for a year, during which period they attain to a confiderable fize, and are called fort-beads by the failors. The mother, at the end of that period, is extremely lean and emaciated, while her cub is fo large and fat, that it frequently yields above fifty barrels of blubber. The next year after they have left the breaft, they are called flunts; because they decrease in their fatness, and yield scarcely an half of the produce that is obtained from them when fuckling. After two years the young whales are called full fift; and though for a while they continue of an inferior fize, there is no mark by which their age can be afcertained.

Though the whales are gregareous animals, yet every individual



Plate I. CACHALOT WHALE

individual propagates only with those of its own kind; and without mixture of breed, they transmit an unpolluted race to posterity. When they are seen in shoals of different kinds together, or making their migrations in large companies from one ocean to another, their object probably is security and mutual desence. Hardly any instinct less powerful than that of self preservation, from the attacks of smaller but more powerful sishes, could induce them to an union by which the scarcity of food must be so greatly increased.

SECTION II.

The Common Whale*.

This is the largest animal known. In the north sea where it is most frequently taken, it measures from ninety to an hundred seet in length; and there is reason to believe that before the sishery had committed such vast depredations, there were many of this species seen of a far superior size †. In the warmer latitudes where they are less frequently taken, and consequently have time to gain their full size, they are still seen of the immense size of an hundred and sixty seet; a circumstance which seems to bestow credibility upon the relations of ancient writers.

Though

Balena, Rondelet. Willough. Ichth. La Balcine ordinaise, Briffon.

[?] British Zool. Class iv. Gen. I.

Though the ancients were acquainted with this species of whale, yet it does not appear that they knew its uses, or practifed the sishing of it. Aristotle has described it by the name of the bearded whale, from those hairs or strainers that surround the mouth to prevent the scape of its food, when the animal discharges the water from its mouth.* Pring has given it the name of Musculus, probably for the same reason.

It has already been remarked, that the food of this fpecies is the medufa or fea-blubber; and it is probably the necessity of procuring this food that consines the animal in its residence to the artic circle. Few of them are ever seen so far fouth as the *British* shores, though the ancients mention a large kind that obtained its name, from frequenting our coasts †.

The head of the common whale is equal to one third of the fize of its body: The fiftule, or two orifices for species has no teeth: in their room are fituated the black horny lamine called whale bone, so long used in the ladies stays, in the construction of umbrellas, and for various other purposes. These lamine give off a part of their substance which constitutes those bristles that surround the mouth, the supposed use of which has already been described. Closely consined by these bristles lied the tongue, the tenderest part of the animal, which was formerly salted up as a great delicacy §: It is now extracted only for the sake of the oil; of which it contains no less than six barrels.

About

^{*} Musingros Hift. anim. Lib. iii. cap. 12.

[†] Quanto Delphinis Balæna Britannica major. Pliny, Sat. E.

[#] British Zoology.

[§] Roudelet, apud Willough, p. 37.

About four yards distant from each other appear the eyes, externally not larger than those of an ox, but constituting a pretty large ball within. This picturesque visage is rendered tremenduous by the large opening of the mouth, which, when the jaws are extended, is no less than eighteen feet wide*. On the back of this animal there is no fin: but the two lateral ones are exceedingly large; and with them, as we have icen, it supports its young. The tail is broad, semilunar and horizontal; it is of such vast strength, that by a stroke of it upon the surface, the water is so much agitated, as to overset a boat if it is near. The penis is included in a strong sheath, and is seven feet in length †.

The colour of whales varies very much; the back being in some red, and the belly generally white. Some are marbled with black and white, while others are entirely black, and some white. Their skin is smooth and slippery, and their colours in the water are extremely beautiful. In the belly of one described by Rondeletius there was found no kind of sish, only a fort of mucus, foam, water, and sea weed.

The Fin Fish t.

This species is distinguished from the common whate by the fin upon the back, placed low, and near the tail is. It is sometimes found in the *British*, but is more frequent in those tracks of the northern ocean, where the Vol. III.

^{*} Idem ubi fupra. † British Zool.

[‡] Le Gibbar. Briffon. Balæna Phyfalus. Linnæus

⁵ British Zool,

whale fiftery is carried on. It is, however, a booty which the fiftermen feldom choose to pursue: The whale-bone adhering to its upper jaw is short and knotty, and therefore of very little value: The blubber also yielded by this species is very inconsiderable in quantity; and these circumstances, added to its extreme serceness and agility, render the capture both dissicult and dangerous; hence our seamen generally neglect it.

But meagre as this animal may feem to those whose object is the procuring of oil, it is held in great esteem by the miserable *Greenlanders*; for its slesh assords them a food which to men so poorly supplied is very agreeable.

This fish is generally of the same length with the common whale, but of a much more flender conformation. The lips are brown like a twifted rope: The spout-hole is as it were fplit in the top of its head, through which it blows water with much more violence, and to a greater height, than the common kind. The fishers are not fond of feeing it; for on its appearance the others retire from those seas*. It is impossible to determine whether this fpecies be the same with the Physalos and Physater of the ancient writers, fo vague are the terms in which they fpeak of that fish +. If that particular name was assigned it from its faculty of spouting water, or blowing, the habit is not peculiar to any one species, but common to all the whale kind. It would appear from the name given it by Linnaus, that he believed this to be the animal spoken of by these writers. Rondeletius and Gesner have been of the fame opinion; but the description the first of these writers has given of this animal is in many respects

P British Zool.

[†] Vide Oppian Halcut, L. 1. 1. 368. Oelian Hift, anim, and Plin. lib. Ex. c. 5.

respects erroneous; of this nature certainly is the assigning it sharp serrated teeth*. There are seven different kinds
of the whale properly so called; sive of these are enumerated among the British sishes. The common whale, the
sin-sish above described, the pike headed whale, the round
lipped, and the beaked. All these differ from each other
in size and sigure, as their names imply: Their manner of
living is also somewhat different, the last described being
more active and sierce than the rest. None of them have
a large swallow, when compared with other kinds of sishes;
none of them accordingly are very voracious; and if compared to the cachalot, that enormous tyrant of the deep,
their manners will appear harmless and gentle.

The Narwhal, or Sea-Unicorn +.

The narwhal is about fixty feet long, of a more flender make than the common Greenland whale; and its fat is in less abundance. It inhabits the feas of Iceland and Greenland, and is feen in the fame northerly regions with the rest of the cetaceous tribes ‡. Nature has, however, distinguished it from every other inhabitant of the deep, by that formidable weapon in the form of a tooth, which projects from its upper jaw. Amongst all that variety of armour which she has conferred upon her different tribes in the animal kingdom, she has contrived no instrument of destruction so dreadful as the horn of the narwhal.

H 2

Tt

^{*} Rondelet. apud Willough. cap. iv. page 41.

⁺ Monoceros, Willough. page 42.

¹ Vide Bartholin, apud Willough, page 42.

It feems yet doubtful whether this extraordinary wead pon of this fifth grows naturally fingle or double in the animal. There is a skull at present in the Stadthouse of Amsterdam, which is armed with two of these teeth, which demonstrates, that in some subjects at least, this instancent is double*: But those which are taken with one tooth, give no fort of indication of having been deprived of the other by any ascident: there appears no socket, nor any remains of a second tooth in the opposite jaw; all there is plain and even.

This extraordinary instrument generally springs from the upper jaw on the left fide; into the focket of which it reaches a foot and an half. It is ftriated, and twifted in spires, as we sometimes see a bar of iron; its length is from feven to eight feet, and of the thickness of a man's arm: It is of a white colour, harder and heavier than ivory t. From the fize of this weapon, most naturalists confider it as an horn, rather than a tooth; but it refembles in every respect the tulks of a boar, or an elephant; it rifes like them, from a focket in the jaw; it is of the fame flrong fubflance, and posleffes the fame folidity. Willoughby regards it as the only real example of an unicorn afforded by nature; and after a minute examination of all the fubiliances that are imposed upon the public for the horns of the unicorn, he pronounces them impofitions on the credulity of mankind t.

This naturalist had the greater merit in making a discovery of this nature, because in his time the capture of whales was not very frequent, and the means of detec_

tion

[&]quot; Goldsmith's Nat. Hist. vol. 6.

⁴ Idem ibidem.

Fide Ichthyol, lib. ii. p. 43.

tion were proportionably few in number. The tooth of the narwhal about a century ago was univerfally afcribed to fome land animal: it had often been dug up among fossil substances, and from that circumstance it was naturally bestowed upon a terrestrial owner. Pliny had long ago described an animal resembling a horse, with a single horn springing from the middle of its forehead *: Upon this animal, which a farther knowledge of nature has proved to be sictious, the tooth of the narwhal was unanimously conferred; and the sinding of so precious a remain, was considered as a fortunate incident, that afforded a strong testimony of the veracity of that historian.

But it was not the curiofity of mankind alone, nor the rareness of the object, and the singularity of its form, that brought the narwhal's tooth into such high repute in the different countries of Europe. A medicine was prepared from it, which was long given out by the quacks as an infallible specific against poison, and malignant fewers †. At length, however, these frauds were detected and exposed by one of the privy counsellors ‡ who had a concern in the whale sishery, and received by the return of his own vessels a number of those teeth, some of which were of the enormous length of three yards.

The error of supposing this armour of the narwhal a horn, has led some writers to suppose, that as among quadrupeds the semale was often found without horns, so these instruments of defence were only to be found in the male. This, however, has often been contradicted by actual experience; both sexes are found armed in this manner:

^{*} Histor. Natural.

[†] T. Bartholin. Act. Med. An. 16736

t Henry Muller,

manner; and in all the varieties of the horn, whether wreathed or finoeth, bended or firaight, it is uniformly flrong, fharp, and deeply fixed. There can be no doubt, but that an inftrument of this nature is intended for the defence of the animal on which it is bestowed. It is thus that the narwhal uses it; whenever it is urged to employ this terrible instrument, it drives directly forward against its enemy, and pierces him through.

But notwithfinding this implement of war, and its amazing velocity and strength, the narwhal is one of the most harmless and peaceable inhabitants of the ocean. It wants teeth for chewing, and a throat for swallowing any bulky prey: Of consequence it commits hostility against no animal; but is constantly seen sporting inossensively among the great monsters of the deep, never attempting to injure any of them. It is called by the Greenlanders the forerunner of the whale; for wherever it is seen, that fish feldom sails soon to appear. The manners of these two species nearly resemble each other; the food of both is those infects which we have already described; and both are peaceable and innocent, though qualified by their strength or their arms to spread general destruction *.

So little does this fish avail itself of these implements with which nature has provided it, that they appear rather an impediment, than a means of defence. It is at no pains to keep them in repair for action; but on the contrary, the tooth is constantly seen covered with weeds, since, and all the filth of the sea. In one instance they evidently operate to the defenction of the owners; for the narwhals being gregarious animals, they are no sooner

de Goldsmith's Nat, Hist.

fooner attacked by a fishing vessel, than they croud together in such a manner, that they are mutually embarassifed by their tusks, and are prevented from finking to the bottom. In this situation the harpooners seldom fail of striking one or two of those that are longest detained upon the surface of the water; and the quantity of the oil which they produce, renders their capture an object of very considerable emolument.

SECTION III.

GEN. III. The Cachalot.

THE fishes of this genus are not of such an enormous fize as those last described: They are, when full grown, from fifty to fixty feet long, and fixteen feet in thickness. Their heads are still more disproportioned to the fize of the body, than that of the common whale: in the latter animal it is equal to a third of the body; in the former it constitutes an half. The cachalots are distinguished from all the other cetaceous tribes, by having sharp arcuated teeth in the lower jaw: Their bodies being more Mender, they are more active than the Greenland whale; are capable of remaining longer at the bottom; and yield a fmaller quantity of oil. The tongue is commonly fmall, but the mouth and throat are fo capacious that the animal could eafily fwallow an ox*. The teeth are about feven inches long, exceedingly thick and hard; they enter, when the mouth is shut, into a number of cavities in the upper jaw prepared for their reception +.

This

[&]quot; Goldsmith's Nat. Hist.

[†] British Zool. Class iv. Gen. 2.

This formidable conformation of the mouth and throat feems to indicate an extraordinary degree of voracity in these animals. The history of the cachalot corresponds to these appearances: for while the stomach of the whale is feen to contain hardly any thing but froth, that of the cachalot is crammed with a variety of different kinds of fishes; fome half digefted, others whole; fome fmall, others eight or nine feet long. The cachalot, therefore, is probably one of the most rapacious sishes of the deep: and is as destructive among the lesser tribes, as the whole is harmless. But it is not to the smaller sishes alone, that this animal is formidable; among thefe the contest is foon ended, for it can devour thousands at one swallow; it purfues and terrifies those of its own order, the dolphin, and the porpelie to fuch a degree, that they are frequently driven ashore in endeavouring to escape.

Of the cachalot there are no less than seven varieties: That of a black colour, with two sins; the cachalot with a white back, and the same number of sins; the species with its spout-hole in the neck; that with the spout near the mouth; that with three sins, and sharp pointed teeth; that with three sins, and sharp edged teeth; and, lastly, the cachalot with three sins, and slat teeth. All these were indiscriminately termed spermaceti whales, till Mr. Pennant borrowed that name from the French, by which they are now distinguished.

From the smallness of its size, as well as its sizeceness and agility, the capture of the cachalot would seldom be attempted by the sishers, were it not for the sake of those valuable medicines, spermacet and ambergris,

Vol. III. I which

^{*} Dale's Harwich. p. 413.

Y Vide Brit. Zool. Class iv. Gen. 2.

which these animals have been found to contain. The various purposes to which these substances are applied, both as drugs and articles of luxury, have rendered the cachalot, which supplies them, a sish in great request, and its capture the most advantageous object in the Greenland trade.

Spermaceti is the name erroneously given to that subflance which is found in the head of the cachalot, and which is by no means the femen, but the brain of the animal. Goldsmith gives the following account of the method by which it is extracted *. The outward ikin of the head being taken off, a covering of fat appears, about three inches thick; and under that, instead of a bony skull, the animal has only another thick skin, that ferves for a covering and defence of the brain. The first cavity or chamber of the brain, is filled with that fpermaceti which is supposed of the greatest purity, and is of the highest value. From this cavity there is generally drawn about feven barrels of the clearest spermaceti, that, thrown upon water, coagulates like cheefe. Below this there is another chamber, just over the gullet, which is about feven feet high; and this also contains the drug, but of less value. It is distributed in this cavity, like honey in a hive, in fmall cells, separated from each other by a membrane like the inner skin of an egg. In proportion as the oily substance is drawn away from this part, it fills anew from every part of the body; and from this is generally obtained about nine barrels of oil. Befides this, the fpinal marrow, which is about as thick as a man's thigh, and reaches all along the back-bone to

ing

^{*} Nat. Hift. vol. vi. p. 219.

the tail, where it is no thicker than one's finger, affords no inconfiderable quantity.

Formerly the spermaceti was obtained but in small quantities, and was fold at a very high price, from the supposition of its great efficacy as a medicine. Though it still enters into the compositions of the apothecary, yet it is rather to give a consistency to his medicines than to add to their virtue: and since the art of converting the whole oil of the cachalot into spermaceti by boiling has been discovered, the article has decreased rapidly in its value. It has now fallen below the price of wax, and is used instead of it for candles.

Ambergris is the other medicine, for which mankind are indebted to the cachalot; and this substance, rather than the former, should have obtained the name of spermaceti, because it is found in the place where the seminal vessels of other animals are commonly situated. For a long time the world was taught to believe, that ambergris was a substance to be found only in small quantities upon the surface of the water. The trade in these articles was originally in the hands of men of obscure and suspicious characters; and this was one of the arts by which they endeavoured to add to the myslerious nature, and value of the commodity. Time, which reveals the secrets of the mercenary, has at last discovered that this medicine is the produce of the cetaceous sishes *.

Among the intestines of the cachalot is found a bag three or four feet long, silled with liquor of a yellowith colour, and thinner than oil; and in this sluid, the ambergris is feen sloating in round lumps, from one to twen-

I 2

^{*} Goldsmith's Nat. Hift. ubi supra.

ty pounds weight, and never above four in the fame fish. These balls of ambergris, the purposes of which, in medicine and perfumes, are so well known, are not indiscriminately found in every fish: it is only the oldest and strongest that yield it in any considerable quantity.

The Blunt Headed Cachalot*.

This species sometimes visits the coasts of Britain: A dead one was cast ashore near Edinburgh in the year 1769, which measured sifty-sour feet, from the mouth to the tail; and its greatest circumference was thirty feet.

The head of this animal is of an enermous fize, far exceeding the proportions of the whale. The upper jaw projects five feet beyond the lower; and its length is about fifteen feet, the other being only ten. Near the fnout, which is quite blunt, and near nine feet high, is placed that orifice peculiar to the cetaceous order, by which they fpout the water. The lower jaw is armed with forty-fix teeth, all pointing outward to meet the fockets, where they enter into the upper. The teeth are about feven inches in circumference at the bottom, fharpening as you approach the top; they are all bent, and like the teeth of the other cetaceous fishes, they are white,

^{*} Le cachalot a dents en faueilles. Briffon. Physiter. Microps. Lin. Syd. 4 British Zeology, class iv. gen. 2.

white, and polished like ivory. Far back, and towards the hind part of its monstrous head are placed the eyes, which are very small in proportion to the fize of the animals whose motions they are intended to direct. The back sin of this species is wanting, and in its place there is found a large protuberance: The two pectoral sins are placed hard by the corners of the mouth, and are about three feet long. The penis is seven feet and a half; the tail bifurcated, and sourteen feet from tip to tip. Such are the monstrous dimensions of this unshapely animal. In its great outlines we still recognize the general characters of the order; and though inserior in size to the whale, it is far more tremenduous in its aspect, and sierater in its manners *.

The round headed Cachalot is

This fish is described by Sibbald ‡, who mentions a shoal confisting of an hundred and two that was cast on shore at the same time upon the Orkney isles. According to that writer, it wanted the spout-holes, that are reckoned characteristic of this order of sishes; it is probable, however, that what he has described as nostrils, was this

^{*} Vide Phil. Tranfact. vol. LX.

[†] Physiter Catodon, Lin. Syst. Le Petit Cachalot, Brisson,

i Sib. Phalain. ix.

this opening, which nature, as we have already feen, has destined for a different purpose.

This fpecies is far inferior in fize, to that above deferibed; the largest of that great number just mentioned did not exceed twenty-four feet in length. Its head is round, and the shape of the teeth the reverse of those of the blunt headed kind. They are smaller at the back, than at the top, where they are quite slat, marked with concentric lines. Their bottom is pierced with a small orisice, bearing no resemblance to the large cavity, deferibed in the former species. The back sin is wanting; instead of it is a rough knotty kind of space *.

^{*} British Zoology.

SECTION IV.

GENUS IV .- The Dolphin.

In reviewing the different tribes of cetaceous fishes, we find them, in proportion as they diminish in fize, growing still more active and voracious. The sishes of this genus, which comprehends in it three different families, that of the dolphin, the grampus, and porpesse, are all much less in bulk, than the common whale. They have all sins upon the back, and like the rest of the whale kind, they have heads disproportioned in fize to the rest of the body. Each of the three species have both jaws armed with formidable teeth; for nature has not conferred upon them an extraordinary voracity, without bestowing the means of gratifying it.

From their great agility, the fishes of this genus are not frequently taken. They seldom remain a moment above water; sometimes, indeed, their too eager pursuits expose them to danger; and a shoal of herrings allures them out of their depth. In this situation, the wretched animal continues to bounce and sounder in the shallow water, till it is knocked on the head, or till the returning tide again comes to its relief. The porpesses are often allured up the Thames in this manner, till they are surrounded with

with boats, from which the fishermen shoot them with fire arms*.

In their internal structure these animals resemble quadrupeds, and the other cetaceous sishes. Conception is begun by the introduction of semen, and after ten months, a persect animal is produced. They bring forth their young only in the summer season; in the month of October the setus has been extracted from a dolphin but newly begun to form ‡. The young, after being produced, continue to follow the parent, till they have attained their sull size, which is in about ten years. The longevity of these sishes was computed by Aristotle to be upwards of thirty years: The method practised to ascertain it, seems but an aukward one: it was, by cutting off a part from the tail sin, and returning the animal to the fea; and if the sish happened afterwards to be taken, its age was guessed at from that circumstance.

. The Dolphin §.

Concerning the figure of the dolphin, properly fo called, the ideas of the ancients were exceedingly erroneous. Upon their marbles and coins, they are uniformly defigned of

[·] Goldfmith's Nat. Hift.

[†] Mistot. Hist. anim. lib. vi.

[‡] Willough. lib. ii. p. 30.

[§] Delphinus Antiquorum, Willough, Le Dauphin, Briffon.

Plate II.





of a crooked form: From the descriptions of the poets, it appears that they entertained the same inaccurate notion of the shape of this animal*. Before a storm the dolphins are observed to be in great commotion, and frequently to leap above the surface of the water. It is probably in this attitude, almost the only one in which they were then seen, that they have a crooked appearance to the eye. The natural stape of the dolphin is almost straight, the back being very slightly incurvated, and the body slender.

But it is not in the external form alone of the dolphin that the ancient writers were mistaken: Their philosophers and historians seem to have contended who should invent the most extravagant sables concerning it. This fish was celebrated in the earliest time for its fondness of the human race. It does not dread mon, fays Pliny, as a ftranger; but comes to meet him in the ship, and to sport around the vessel: hence it was distinguished by the epithets of boy loving, and philanthropitl t. It was confecrated to the gods; and therefore was honoured with the title of facred fish. Various are the services which this race are faid to have performed to man: Pliny, Melian, and many of the other early writers fet no bounds to their belief of the tales related concerning the dolphin's attachment to the human species: Piny the younger, who lived in an age lefs remarkable for credulity, makes an ap logy for the flory of the enamoured dolphin of Hippo, which he narrates in a beautiful manner &.

Vol. III. K From

^{*} Tumidumq: pando tranfilit dorso mare. Senec. Trig. Agam. 450.

[†] Bellonius apud Willough. p. 30.

[‡] British Zoology, vol. iii. p. 65.

[§] Vide Epift, lib. ix. ep. 33.

From what cause the ancients were prejudiced in favour of the dolphin it is not easy to determine. The figure is fuch as could create no partiality; its manners, which are fierce and rapacious, could still less endear it; and it does not appear that this fish shews any attachment to mankind, more than the rest of the cetaceous tribes. It perhaps has arisen from commiseration, on hearing that plaintive mean, by which, when taken, it expresses its fufferings. In a veffel where feveral dolphins were confined, fays an old writer, I spent a night of great pain and uneafiness, so feelingly did these wretched creatures express the mileries of their condition, in eries and lamentations, refembling the human. Their fufferings forced from me tears of compassion; and while the fish rman was afleep, I forced the one that feemed to fuffer the greatest agony overboard into the fea. This act of tenderness availed me nothing; for the moaning of those that remained behind, feemed only to be encreated; who made figns too plain to be mifunderstood, that they wished for a similar deliverance. Thus I fpent the night in unavailing forrow for fufferings that I could not alleviate *.

These prejudices in savour of the dolphin, are now so wholly obliterated, that even the common people regard them in a very different light. Their appearance is far from being deemed a savourable omen by the seaman; for their boundings, sports, and srolies in the water, are held to be fure indications of an approaching storm †. It is not known whether these motions of the dolphin are the gambols of pleasure, or the effects of sear. They pro-

bably -

^{*} Gillius apud Willough. page 31.

^{&#}x27;s British Zoology, and Willoughb. lib. ii. p. 30.

hably are occasioned by some presentiment of that turbulence and commotion which is soon to take place in their element; a commotion which obliges the smaller sishes to sink to the bottom, and which of course deprives them of their accustomed prey.

Almost every thing marvellous in natural history difappears on farther acquaintance, and more accurate examination of the subject. This has been the fate of all those extraordinary circumstances related of the dolphin. So great is the number of those animals that play around this island, and so frequently are they taken, that experience and ocular testimony, soon contradicted those tales that had been the growth of credulity and ignorance.

The fwiftness of these animals in the pursuit of their prey, and their extraordinary avidity to fecure it. are almost the only peculiarities (if such they can be called) which farther experience has afcertained to belong to them, or at least to characterise their manners. Belon* long ago remarked, that they eafily came up with the thip in which he was, though in full fail before a brifk gale; and that they played around her, with a velocity refembling that of a bird in the air. This extraordinary swiftness of the dolphin seems to be the combined estect of agility and strength: by means of it they are enabled to make dreadful havoc among the falmon, herring and mackrel They are frequently fren fighting pitched battles with the amint. The two different kinds of fish range themselves on each fide, like contending armies. In one of these engagements, at the opening into the Dar-

danelles,

^{*} Hift. Pifcium.

[†] A fish in the Mediterranean refembling a salmon.

danelles, the areie were vanquished, and put to a total rout by their victorious enemies *.

The dolphin, like fome other kinds of cetaceous animals, have been observed to emigrate from one sea to another. They have been known to pass from the Mediterranean into the Black Sea; and after having remained for some time there, to return to their former haunts. When they are taken, as sometimes happens, in the nets placed for other sishes, if they are not strangled by their own efforts to escape, they will live for a long time after being brought to the shore: One has been carried alive from Montpellier to Lyons †.

In former times the flesh of the dolphin was reckoned a great delicacy: it was bought by princes at a high price, and presented at their most formal entertainments. This Rondeletius declares, he had more than once obferved, though not among his own countrymen, where the practice of eating these fishes had entirely gone into disuse. This was also the case in England in the days of Willoughby ‡. At a more ancient period, however, these fishes were not only eaten, but held in as high repute among the English, as any other nation. Dr. Caius says, that one which was taken in his time, was thought a present worthy of the Duke of Norson, who distributed part of it among his friends. It was roasted and dressed with perpesse sauce; a composition of fine white bread mixed with vinegar and sugar §.

The mout of the dolphin is narrow and pointed, not unlike

[#] Hift. Pifcium.

[†] Vide Gefner and Rondeletius.

^{*} Vide Hift. Piscium lib. ii. p. 31.

[§] Caij opuscula, p. 116.

unlike the beak of fome birds; and hence the French call it the fea-goofe *. Its teeth are forty-two in number, placed in both jaws, and bended a little towards the mouth. They are placed at fuch a distance, that each tooth enters into a space between two in the opposite jaw.

The skin of these fishes is smooth, of a dusky colour on the back and sides; while the belly is white. The form of the tail is semilunar; that of the back sin, high, triangular, and placed nearer to the tail than the head?

The Porpeffet.

The fishermen do not distinguish this fish from the dolphin, which it nearly resembles: Both are known by the
name of porpesse among the English failors; and the
French with indiscrimation call them marsonins. The
latter word is derived from the German, and the former
from the Italian language: the import of both is the
the same, and expresses the resemblance which this sish
bears to a hog §. It is probably from the particular shape
of the nose, that the animal has obtained this name
so generally among the modern nations of Europe. The
shout projects considerably, though not so long as that of

^{*} British Zoology, L'oie de la mere.

[†] Idem, Class iv. Gen. iii.

i Le Marfouin. Briffon. Delphinus Phorcana, Lin. Syft.

[§] Meerschwain denotes a sea-swine, and Percopelee may be translated hog-fish.

the dolphin: It is furnished with very strong muscles, which enable the fish more readily to turn up the fand; for by this means the porpesse, as well as the swine, procures a great part of its food. When prey fails at the surface, it dives to the bottom in search of sea worms, and fand cels which it digs up there.

The porpeffe is diftinguished from the dolphin, by the fuperior thickness of the head, as well as the shortness of the nose: It is commonly of smaller size; the body gross and fat towards the head, but tapering away in the form of a cone, till it becomes slender at the tail *. The colour of the back is generally black, and the belly white: This, however, is not an uniform character; for in the river of St. Lawrence there is a white porpesse; and there is in the British channel a small species called thorn-backs, that are various, sometimes brown, white and spotted †.

Both jaws of the porpesse, as well as those of the dolphin, are provided with teeth; but in the former they are much more numerous, and of a smaller fize than in the latter animal. They are forty-eight in all, moveable, and so situated as to lock into each other. The eyes are small, resembling those of the human species, both in size and in the distribution of their humours ‡. It has no branchiæ, nor any aperture in their place; the spout-hole is upon the crown of the head, of a semilunar shape, and divided internally by a cartilaginous membrane, which appears at the summit of the head like the comb of a cock.

Towards the note of this animal there have been ob-

^{*} Willough. Hift. Pifc. page 28.

[†] British Zool. class iv. gen. 3.

[‡] Dan. Major apud Willough.

ferved no less than fix concealed apertures or foramina leading to the brain, which have been supposed to be its organs of smell; a sense which it probably possession a greater degree than other sishes.* The tongue is slat, pectinated at the edges, and sastened down to the bottom of the mouth †.

The structure of the viscera of this fish appears more complex, and indicates an organization somewhat superior to the rest of the tribe; we are persuaded however, that a more minute examination of the parts would discover an exact conformity between it and the different cetaceous tribes. The porpesse, according to Willoughby \$\frac{1}{2}\$, has three stomachs: the first, large, wrinkly and muscular, for reducing the shells of fish which it devours; the second was also large, empty, and furnished with smaller muscles; the third appeared a kind of adjunct to the other two, and contained a quantity of assimulated chyle, of a livid colour.

The whole body of the porpeffe, immediately below the skin, is covered with a coat of fat resembling lard, of an inch thick. Various uses, in the economy of the animal, are affigured to this substance: it is supposed to defend the body from injury and cold; to diminish its weight so as to make it equiponderate with the water, in order that the sich may alternately rise and sink without effort, while employed in the different operations of breathing and collecting food. Beneath this coat of sat lies the slesh, red, thick, and muscular, resembling that of a hog §.

Of all the cetaceous fishes, the porpesses are found in greatest

^{*} Dan. Major apud Willough." † British Zool, ubi supra.) † Hist. piscium, page 33. § Idem ubi supra.

greatest abundance around the British isles. In such vast numbers do they appear in some places, that the sishermen are not safe to venture among them in a small boat. When gamboling upon the summit before a storm they appear almost to darken the surface of the water *. In times of sair weather, they are seen herding together, and pursuing the shoals of macked with great impetuosity. Their manner of hunting after their prey is distinguished by the same eagerness and disposition to act in concert, that is observeable among dogs when pursuing a hare †. They are seen urging them from one bay or creek to another, deterring them from the shallow water, and driving them towards each other's ambush, with all the art of an experienced greyhound.

The Grampus ‡.

This animal feldom exceeds the length of twenty-five feet, but is remarkably thick in proportion, the breadth being to the length, as ten to eighteen. It was with reafon that Pliny called it an immense heap of sleth, armed with dreadful teeth §. From its resemblance to a wine cask in the thickness and rotundity of its form, it obtained the name of orca among the ancients.

The nose of the grampus is flat, and turned up; the under

^{*} Pennant's Brit. Zool. † Gold mith's Nat. Hist.

¹ L'Epaulard, Briffon. Delphinus Orca L n. Syft.

[§] Cujus imago nulla representatione exprimi potest alia, quam carnis immensa dentibus truculenti s. Lib. ix, cap. vi.

under lip is so thick and heavy, that when the animal is on its belly, it is separated from the upper *. The teeth are sixty in all, thirty upon each jaw; and between each is a space adapted to receive those of the opposite side, when the mouth is shut †. Those upon the fore parts are blunt, round, and slender, the innermost sharp, and of great thickness. The spout hole in this species, is placed at the top of the neck; the animal is covered with a strong skin, like leather, black upon the back, but displaying a large spot on each shoulder, of the same colour with the belly, which is white; the sides are marbled with black and white spots ‡.

The grampus is of all the cetaceous fish the most fierce and voracious; it does not even spare the porpesse and whale, which are animals of its own order and of superior fize; the latter of these sish it is said to tear and mangle in such a manner with its teeth, that it bellows like a bull from the pain of its wounds ||. The sishers who formerly repaired to the coasts of America, prohibited the savages from destroying this sish; for by its assistance they were aided in the capture of whales, seals, and other animals: whilst the grampus pursued these, and the other monsters of the deep, they were frequently obliged to betake themselves to the shallow water in order to escape its merciles teeth, where the seamen were ready to kill them with javelins and harpoons.

This species is a profitable capture for the fishers, as it is said to yield a very considerable quantity of oil; like the porpesses, it sometimes appears upon the British Vol. III.

[&]quot; Vide Will. Orca. Lib. ii. p. 40.

⁺ Sibbald, Phalæn. 7, 8. | Brit. Zool. Classiv. gen- iii.

Willough, ubi fupra.

coasts; but its great resort is off the North Cape of Norway. The number and position of the fins are the same as in the dolphin; and it resembles all the other cetaceous animals in swimming against the wind, and in that tumbling and agitation which it discovers on the approach of a storm,

CHAPTER V.

SECTION I.

General Characters of the Cartilaginous Fiftes.

In the organs prepared by nature for the respiration of animals, there is a great difference between those formed within the body, called lungs, and that apparatus at each fide of the head, known by the name of gills. Among all the terrestrial animals, and the cetaceous fishes, breathing is performed by the former of these contrivances, and among the spinous fishes it is carried on by the latter; between these two orders of the animal kingdom, there is placed a third, which unites them in the great scale of being, and partakes in some measure of the nature of each. These are the cartilaginous fishes, which, while they are furnished with apertures on each fide of the neck, corresponding to the gills of the spinous fishes, are also supplied with lungs within the cavity of the body, analogous to what we have already feen among the terrestrial and cetaceous tribes.

From this double conformation in the organs of breathing, it would at first view appear, that the cartilaginous tribes were a kind of privileged order of animals, pofsessing a greater variety of endowments than the rest of T. 2

the animal kingdom. It would appear that they are capable of subsisting indiscriminately in either of these two elements, to each of which the structure of their parts feem to have a reference; it was no doubt this aspect of the cartilaginous sishes, which presented itself to Linnaus when he arranged them in the same order with those animals, properly amphibious *, that are capable of subsisting either in the water or in the air. Of this power, however, this class of beings is by no means possessed; for, though they are capable of supporting life for a longer time on shore, that the spinous sishes, yet there is hardly any of them that can live there above a few hours at a time †. Whatever, therefore, be the structure of their organs, there is none of them properly amphibious, or that justly merits a place among that order of beings.

But although the double structure of the organs of respiration in the cartilaginous sishes, will not justify us in placing them among the sishes, tortoises, lizards, and other animals that are strictly amphibious, it forms, however, a very palpable mark of distinction between them and the cetaceous sishes on the one hand, and the spinous upon the other; from the latter they are known, because, though they have apertures corresponding to gisls, they universally want the bony opercula which covers them ‡.

These apertures by which they breathe, are placed always near the head, but have not uniformly the same position: In some they are placed beneath, as in the rays, and other slat sish; in others on the sides, as among the shark

^{*} He has termed them amphibia nantes. Vide Systema Naturæ.

[†] Willough. Icthyol.

⁵ Pennant's Brit, Zool. Class iv. Div. ii.

Mark tribe; and among a third class, they are fituated on the top of the head, as exemplified by the pipe fish.

From these foramina, there are two bending cylindrical dusts, that run to the lungs, and are supposed to convey the air that gives their organs their proper play: The heart, however, has but one valve; and hence, their blood does not perform that double circulation which obtain among the cetaceous kinds, and the lungs seem rather an internal assistant to the gills, than for supplying the same purpose which they do in quadrupeds, for they want both the pulmonary vein, and artery.

Another flinking character by which this order of the animal kingdom may be diftinguished, is, that their flesh is supported by cartilages instead of bones. In the cetaceous tribes we have seen, that the bones were hard, and filled with marrow, resembling those of quadrupeds, while these parts in the spinous tribes, are small, slender, and pointed, resembling thorns, and generally solid throughout. The size of other animals increases with their age, and stops at a certain point; but from the pliancy of the bones in this tribe, they seem to have no bounds placed to their dimensions, and it is supposed that they grow larger every day till they die †.

A third general character of these sistes from their manner of generation; some of them are oviparous, while the greater part produce living young; the same duplicity of character attending them in this respect, which we have already taken notice of with regard to their conformation. In those that are viviparous, the progeny is excluded from an egg hatched within the bo-

dy `

dy of the female; this egg confifts of two parts, the white and red, as in those of birds, and instead of a shell, it is lodged in a case formed of a thick tough substance, resembling a horn that has been softened in boiling water *.

The oviparous fishes of this class produce their young nearly in the same manner; in these as well as in the viviparous kinds, the egg is hatched for some time in the belly of the semale, but before the setus is disengaged from it, is excluded from the womb, and left till time and the warmth of the sun bring the inclosed offspring to maturity. Both differ remarkably from the whales and the quadrupeds, whose eggs start from the ovarium, and enter the uterus at the time of conception, where they remain only a few days, till they leave that state, and are formed into a complete setus.

There are other discriminating characters belonging to this order, less discernible indeed, but such as mark a disference between them and the adjoining tribes. The spinous sishes we saw had no aparatus for hearing; it is probable, however, that the cartilaginous are furnished, though imperfectly, with the organs of this sense; upon each side, near the eyes, there is an aperture which opens into the mouth, the supposed use of which is partly to convey the water into the mouth, which is expelled through the gills, and partly to convey sounds to the censorium. In this respect, therefore, the cartilaginous sishes may be deemed a more accomplished race than many of their companions of the deep.

The

British Zool. ubi supra.

The fishes of this order are possessed both of gills and lungs, and are thereby in some measure capable of deriving aid from both elements, to their necessities and enjoyments; the manner of their parturition is also various: and we are now to contemplate them in the double enjoyment of another qualification. The males of this race copulate with their females, more hominum; and in them the external organs of generation are two-fold *. These instruments project forward from each side of the abdomen; and whether it be that these animals possess more warmth of constitution, or a greater capacity of bringing their offspring to maturity, they generally choose colder seasons and situations for producing their young, than other sishes; many of them propagate in the midst of winter, and far from the shore †.

The cartilaginous fishes, though not so remarkable either for fatness or size, are in general more voracious than any of the tribes we have already reviewed; their livers are indeed fat, and are sometimes employed for the production of oil; it is not, however, from them that man derives either his most pleasant or salubrious food; they are impure and immoderate seeders, and their sless favours of that impurity \$\ddots\$. Of a considerable portion of them, the mouth is placed below the head; a contrivance of Nature, for which an old writer assigns some curious reasons: Their snout, says he, is too small to be divided; and their voracity is so keen, that their own life requires that it should not be allowed a ready or complete gratification §.

From

^{*} Idem ubi fupra. † Goldsmith's Nat. Hist.

[‡] Will. Ichthyol. Lib. iii. p. 45.

³ Vide Rondeletius de piscibus. Lib. iii. cap. 4.

From this conformation of the mouth, a great part of the cartalaginous fishes are obliged to turn their back downwards in laying hold of their prey; a circumstance that requires time, and Fords an opportunity to the smaller fish of making their chape. They devour every kind of fish or flesh, and to their extraordinary voracity, had Nature granted the power of easily apprehending their food, the other kinds of fish must long ago have fallen a facrifice to their gluttony *.

* Idem ibidem.

SECTION II.

GENUS V .- The Lamprey.

OF this genus, Linnaus has enumerated only three species, though it is probably diversified into a greater number of different kinds *; his catalogue confifts of the fea lamprey, the river or leffer lamprey, and the pride. He makes mention of the murana, that fish so much celebrated among the epicures of ancient Rome, although fo nearly allied to this genus, that it feems to merit a place there. There is a species of lamprey esteemed a great delicacy by the modern inhabitants of Italy, different from ours, and probably the same sish with the ancient Murana, as it is reared in ponds, and fed in the fame manner. The genuine characters of these fishes are : A long and flender shaped body, resembling that of a snake; the skin has no scales, but is covered with a slimy glutinous mucus; they have feven apertures for breathing upon each fide, and one like the cataceous fishes upon the top of the head; both the pectoral and ventral fins are wanting.

Vol. III.

M

Wide Syft. Nat. p. 394

The Murana *.

This species, we are told, obtained its name from its continually floating upon the surface of the water, by reason, as was supposed, of its extraordinary fatness. In shape it resembles an eel; the snout more protuberant, sharp and compressed: Its colour variegated with shades of yellow, brown, and black. The head is small, but the or pening of the mouth capacious. The margin of each jaw is surrounded with a single row of very small teeth: upon the snout there are two apertures, and above the eyes other two; the sormer, as is supposed, are the organs of smell, and the latter of hearing. The aucients believed that this animal was endowed with a very acute sense of hearing, and that the sishermen could allure it to the brink of the water by hissing, and by that means take it §.


Like the other fishes of this genus, it wants the pectoral and ventral fins; but a little beyond the head there arises a fin, which stretches along the back to the tail, and turns round to the belly, terminating at the anus: This fin is covered with the common legument of the body, and is easily laid bare by flaying the animal.

The bite of the murana was deemed to poisonous by the fishermen, that in taking the animal, they used every precaution to prevent its effects: They tumbled them immediately

Muræna omnium auctorum. Will. Ichthyol.

[†] Παρα το μυρειν. Vide Macrob.

Will. p. 103. § Plinii. Hift, Nat.





cures

mediately from the net into the boat, where they laid hold of the head with a forceps, rubbed its frout against fome hard fubstance to destroy its teeth, and beat it on the tail, where the life was supposed to lie *. That the principle of life, among eels in general, is placed in the tail, is still a prevalent opinion; and when taken, it is not by beating the head, but the tail, that the fisher endeavours to deprive them of animation.

Poisonous as the bite of these animals may be, and hideous in their external figure as they certainly are, yet these circumstances did not prevent the luxurious citizens of ancient Rome from adding them to the endless list of delicacies with which they pampered their appetites †. This species, the basse (Lupus), and a kind of mullet (Myxo), formed that pride of Roman banquets, the tripatimam; fo called, according to Arbuthnot, from their being served in a machine of three bottoms 1.

The flesh of this animal must be various, according to the manner in which it is fed; it is faid, however, to be in general white, tender, and of an agreeable flavours. It was reared with much care in the fish ponds of Italy, and fold at a high price; of its estimation among the ancients, we have full evidence in all the classics, that have either praifed good eating, or ridiculed gluttony. A fenator of Rome, whose name will be transmitted with infamy to posterity, was highly complimented for the delicacy of his murænæ. Tigellinus, Manucius, and all the celebrated epi-M 2

^{*} Plinii Hift. Nat. lib. 32. cap. 2.

⁺ Pliny, Ovid, Juvenal and others.

Atque ut luxu quoque aliqua contingat auctoritas figlinis Tripatinam, inquit Fænestella apellabatur, summa cænarum lautitia: Una erat Muramerum, altera Luporum, tertia Myxonis pifcis. Plinii Hift. Nat. lib. 35.

[&]amp; Willough. lib. 3. p. 104.

eures of his time, were loud in his praises: No man's siste had so rich a slavour, were so nicely sed, or so exactly pickled. Augustus hearing so much of this man's entertainments, desired to be his guest; and soon sound, that same had been just to his merits; the man had indeed sine muranae, and of excellent slavour. The Emperor was desirous of knowing the method by which he sed his sish to so excellent a reliab; and the glutton, making no secret of his art, informed him that his way was to throw into his pond such of his slaves as had at any time displeased him. Augustus, it is said, was not much pleased with his receipt; and instantly ordered all his ponds to be filled up. Goldsmith, who narrates this story, adds, that it would have ended better, had he ordered the owner to be slung in also.*

The Sea Lamprey t.

This is a British fish, of which the names lampetra, and petromyzon, lately given it, are expressive of a habit this animal possesses of sucking and adhering to stones. The mouth is of a round form, placed rather obliquely below the nose, and resembling that of a leech. Like that animal, it possesses the extraordinary power of sticking to any substance to which it is applied; and that with such sirmness, as not to be drawn off without some difficulty.

^{*} Goldfinith's Nat. Hift.

[†] Petromyzon Marinus, Lin. Syft. La Lamproye de Mer. Belon.

[‡] The first lampetra, a lambendo petras; and the other, from Merges and Muraw, because they are supposed to lick the rocks.

culty *. One that weighed but three pounds, has been known to adhere so sirmly to a stone of twelve pounds, that it remained suspended at its mouth. This amazing power of suction must arise from the animal's exhausting the air within its body by the hole over the nose; while the mouth is closely fixed to the object, and permits no air to enter †. The weight which the lamprey is by this means able to sustain, will be equal to that of a column of air of the same circumference with the animal's mouth.

The lamprey fometimes grows fo large, as to weigh four or five pounds; its colour is dufky, irregularly marked with dirty yellow. In the mouth are placed twenty rows of small teeth, disposed in circular orders, and placed far back near the throat, four, five, and fix, in each row ‡. The branchiæ are situated within the seven apertures, that are found on each side of the neck. Though in shape this animal resembles an eel, yet it is of a thicker and more clumfy form.

The fea lamprey, we should imagine from its name, was only produced in the falt water; it is found, however, very frequently at the opening of large rivers, where they join the ocean. At certain feasons it is found in many of the *British* rivers, and also in the *Irish*. They are fea fish; but, like salmon, quit the salt waters, and mount the rivers about the end of winter, and after remaining there for a few months, return again to the ocean §.

As our manners are probably still far short of that senfuality and extravagance which distinguished the ancient

^{*} Pennant's British Zoology. + Goldsmith's Nat. Hist. vol. 6. p. 270

^{\$} Willough. Ichthyol. p. 106.

S Brit. Zool, ubi fupra. Vide etiam Willough.

cient Romans, we do not hear fo many fanciful encomis ums on particular dishes, nor fuch enormous sums paid to procure them. Accordingly, the lamprey known among us, has obtained no very extraordinary character. It is differently estimated, according to the season it was taken, or the place where it was fed. The best season for them is in the months of March, April and May; for they are more firm immediately after their arrival in the rivers from the falt water, than afterwards. Towards the Summer, and in the hot weather, after they have depolited their spawn, they are observed to be much wasted; and their fish becomes flabby, and bad eating *. Those caught in feveral of the rivers in Ireland, the people will not venture to touch +; and throughout the whole of Scotland they are held in detestation: while those taken by the English in the Severn, are considered by them as the most delicate of all fish whatever. Prejudices, though of opposite kinds, feem to operate in each of these cases, whether in producing difgust or predilection.

At a very early period, we find, the lamprey was reckoned a great delicacy by the English: That they are a heavy furfeiting food, the experience of one of their monarchs fatally testified; for the death of Henry I. was occasioned by eating too plentiful a meal of these fishes ‡. Notwithstanding this accident, they seem to have continued in high esteem, because we find Henry IV. granting protection to such ships as brought overlampreys for the table of his royal confort; and his successor issuing out a warrant to William of Nantes, for supplying him and his army with these sishes, wherever they might happen to march §.

In

^{*} British Zoology.

⁺ Goldfmith's Nat. Hift.

[†] Vide Hume's Hift, of England.

S Rymer ix. 544. apud Pennant,

In proof of the ancient predilection of the English for this fish, it is a custom, from time immemorial, for the city of Gloucester to prefent annually his majesty with a lamprey pye, covered with a large raised crust. As the gift is made at Christmas, it is with great difficulty the corporation can procure any fresh lampreys at that early season, though they offer a sum far exceeding their usual price *. The various methods of taking the lampreys, it is not the business of this work to describe.

The fishes of this genus, from some peculiarity in their conformation, generally swim with their body on a level with the surface of the water; and, it is said, can easily be suffocated, by being immersed in it for any confiderable time. From this circumstance, it is probable that they require a constant supply of air for breathing, and that they are possessed of lungs, though no naturalist has observed them in giving an account of the viscera of these animals.

Among this tribe, copulation is performed ventribus commissis, contrary to the manner of the greater number of fishes: When engaged in the business of procreation, they generally frequent the shallow water, where their habits admit of being easily examined. The female, when ready to spawn, digs a hole in the mud, where she deposits her ova; and in this operation, the power of suction, which we have already described, is of singular advantage; for, should she meet with a stone, though of considerable bulk, she raises and carries it out of the way †. After the ova are excluded, and buried in the excavation thus formed, the parent remains in the

^{*} British Zool, class iv. gen. iv. † Balnerus apud Willough.

neighbourhood till the young are quickened into life. She is then feen, with her numerous family playing around her; which, as foon as they have acquired fufficient frength, the gradually conducts to the fea.

The River Lamprey *.

This species is smaller than the former, seldom exceeding ten inches in length; and is distinguished by transverse lines of a blackish colour. The mouth is of the same round form as in the preceding sish, and is possessed of the same adhesive power. On the upper part of it, is a large bifurcated tooth; on each side there are three rows of very minute ones; and on the lower part there are seven, the exterior of which on each side is the largest.

Like all the other fishes of this genus, there is between the eyes a spout-hole or orifice, resembling that of the cetaceous fishes, and probably destined for a similar purpose, that of ejecting water from the throat and lungs. The back fin of the river lamprey is not continued uninterrupted to the tail, but is broken off; and behind it there rises a second, which goes round the tail, terminating at the anus.

This species, as well as the larger kind, is also eaten, potted and highly seasoned: By some it is preferred to the sea lamprey, being milder tasted. All these dishes

are

^{*} Lampetra medium genus, Will. La lamproye d'eau doulce, Belon. Petromyzon fluveatilis, Lin. Syst.

[†] Willough. page 106, ‡ British Zool. class iv. gen. iv.

are

are probably heavy and unwholefome, more highly recommended by the epicure than the physician *. They are taken in fuch vast numbers in the Thames, the Severn, and the Dee, that they constitute a considerable branch of trade. They are fold to the Dutch, who use them for bait in the cod fishery: besides what are confumed in England, 450,000 have been exported in a feafon for this purpose, at two pounds per thousand.

The Pridet.

THIS species is exceedingly small, being from four to eight inches. It is a British fish, but found also in the rivers of of Germany and Italy. In the river Isis, and other streams near Oxford, they are abundant, where, inflead of concealing themselves, they roll in the mud, and never are observed to adhere to the stones, like the other lampreys 1.

The back is of a livid colour, approaching to black; the belly filvery, and more resplendent, than the larger kinds. The mouth is of an eliptical shape; within it are fix or feven teeth, and above them a small semicircular bone. Upon the top of the head is a fmall tube, or aperture, like the cetaceous fish: The belly fwells and contracts alternately, as in those animals which breathe by means of lungs. It has two dorfal fins; the first rifing about the middle, and supported by small tendons, that N

* Willough. ubi fupra.

VOL. III.

[†] Lapern, or pride of the Isis. Will. Petromyzon Branchialis. Lin. Syft.

[#] British Zoology.


are scarcely visible: The other takes its rise immediately behind it, surrounds the tail, and terminates in the middle space, between the tail and the anus*.

Willoughby describes two other species of the lamprey; the mucu peculiar to the marshes and lakes of Brazil; and the blind lamprey, a small round sish, no larger than May worms: The last marked round the body with 84 transverse lines, or annuli, resembling those upon the terrestrial worm †.

Willough. page 105.

† Fide Icthyol. loco citato.





SECTION III.

GEN. VI. The Ray.

This genus is eafily diffinguished from all other kinds of fish, by a broad, flat, and thin conformation. The five branchial apertures on each fide, are placed under the animal; they want the bony opercula, and are marked by the same peculiarities, that distinguish the other cartilaginous tribes. The mouth, in all the fishes of this genus, is situated quite below the head; while the body of the greater number is beset with spines or prickles, resembling those on the branch of a thorn, a circumstance from which they obtained their ancient name *; that of ray, which we now apply to the genus, is derived from the latin.

This genus comprehends all the rays, thornbacke, and flairs, or skates, which are distinguished by manners as singular as their external form. Willoughby has deferibed fourteen different species, and Linnaus nine; of which no less than eight frequent the British coasts +.

N 2 With

Puros, which fignifies the berry that grows upon a prickly farsb,

[†] Systema Nat. and British Zool.

With regard to their dietical uses, these sishes are very different; some of them affording a plentiful supply of salubrious sood, while others are noxious and terrible; with respect, however, to their nature, appetites and conformation, the similitude among them is perfect and entire.

Of all the larger fish, the rays are the most numerous; and their numbers are in a great measure owing to their fize, and to the protection of those frightful spines which nature has afforded them. There is none of the rapacious tribes, except, perhaps, the cachalot and white shark, that has a fwallow fufficiently large to receive them; and even these are, probably, often deterred from their purposes of destruction by the armour with which their prey is covered *. Of some the fize is such as to defy all the powers of destruction which even the shark himself is known to possess: In England some have been taken upwards of two hundred pounds weight; but that is far inferior to their enormous bulk in other parts of the world. Near the island of Guadaloupe in the West Indies, a ray was killed, thirteen feet eight inches in breadth, and twenty-five from the fnout to the tip of the tail. This member itself contributed largely to this prodigious dimension; for it was twenty inches broad at its infertion, and tapered to a point, by which it terminated fifteen feet behind the body of the animal+.

The fishes of this genus probably attain to a much larger fize, than that of any individual that has ever yet been examined: It is only the smallest of the kind that approach the shores; the larger continue forever prowling

at the bottom, in the unfathomable caverns of the ocean, where they probably continue growing for ages. The utmost bulk of the ray cannot therefore be ascertained; but it has been supposed, that they are the largest inhabitants of the sea *. When, however, they surpass the size commonly observed by the sishermen, they become totally unsit for food; the negroes, and starved savages of America, indeed, cut them up, even of the largest size; and the tenderer parts, when salted, are reckoned among their delicious morsels.

These fish generate in March and April; at which time they fwim near the furface of the water, feveral of the males purfuing one female: With fuch avidity of defire do they engage in the work of procreation, that the fishermen frequently draw up both male and female still adhering together, though only one has taken the bait +. The females are prolific to an extreme degree, there having been no lefs than three hundred eggs taken out of the body of a fingle ray. These eggs, containing the fætus, acquire a tough horny covering after descending into the womb; for while they continue in the ovarium, they are attached to it, like the small eggs in the body of a hen; after they drop into the womb, the formation of the animal becomes visible, and the shell is formed by the concretion of the fluids of that organ; when come to their proper maturity, the young are excluded, but never above one or two at a time, and at the interval of three or four hours. The females begin to cast their purses, as the fishers term it, in the month of May, and continue producing young till September. In October, when their

Coldfmith's Nat, Hift. + British Zools

their parturition ceases, they are exceedingly poor, and then they improve gradually during the winter and spring, till May, when they are at their highest perfection, and when they again begin to undergo the same functions*.

The rays generally frequent those parts of the sea where the bottom is black and muddy, where their voracity leads them indiscriminately to devour every living animal which they can surprise: But although their appetite is ravenous and indiscriminate, they become mere delicate with regard to a baited hook; they devour below any putrid substance whatever, but if the bait has been taken up and suffered to lie for any time in the open air, they will not touch it: they appear by their manner to perceive the line, and to dread it; but the impulse of their hunger is too great for their cautien, and, even though they perceive the danger, if thoroughly hungry, they devour, as if regardless of destruction †.

Both English and Dutch carry on the fifting to a confiderable extent; the feafon at which they begin is early in the winter; and the boats in which the fiftermenput to fea, are of different fizes, according to the distance of the place where they intend to fish. The vessels used in the British Channel, called cobles, are of one ton burden, rowed with three pair of oars, and admirably constructed for encountering a mountain fea. When they go out to fish, every person is provided with three lines; each man's lines are sairly coiled upon a stat oblong piece of wickerwork, the books being baited, and placed very regularly in the centre of the coil. Each line is furnished with

two

two hundred and eighty hooks, at the diffance of fix feet two inches from each other; the hooks are fast and to lines of twisted horse hair, twenty-seven inches in length.

When fishing, there are always three men in each coble, and consequently nine of these lines are suffered together, and used as one line, extending in length near three miles, and furnished with above two thousand sive hundred hooks; an anchor and buoy are fixed at the sirst end of the line, and one more at the end of each man's lines; in all, four anchors, and sour buoys made of leather or cork.

The line is always laid across the currents; the tides of flood and ebb continue an equal time upon our coast; and, where undisturbed by winds, run each way about fix hours; they are so rapid, that the fishermen can only shoot and haul their lines at the turn of the tide; and, therefore, the lines always remain upon the ground fix hours; the same rapidity of tide prevents them from using hand lines, and, therefore, two of the people commonly wrap themselves up in the fail, and sleep, while the other keeps a strict look out, for fear of being run down by ships, and to observe the weather; for storms often rise so suddenly, that it is sometimes with extreme difficulty they escape to the shore, though they leave the lines behind them.

Such is the account which M. Pennant gives * of the manner in which the taking of these sish is conducted in the channel; but there are annually larger vessels, of twenty-sive tons burden, that repair at the time of Lent to the *Dogger Bank*, where they sish for turbot, cod, ling,

and

[?] British Zoology.

and skates; there the business is carried on in a more expeditious manner, for as the fishers have no occasion to wait the returns of the tide, each man takes along with him twice the number of lines, which he continues to bait, haul, and shoot, without interruption.

But this method of fishing, operofe as it may feem, and this extent of line, though it runs three miles along the bottom, is trifling when compared to the exertions made in the Mediterranean by the Italian fishers; there they go to fea in a Tartan, a vessel much larger than ours; and they bait a line of no less than twenty miles long, with above ten thousand hooks. This line is called the parisina; and the fishing goes by the name of pielago *. A piece of tackle of fuch enormous length, it is impossible to hawl and shoot in the same space with the English lines: it remains for a confiderable space in the sea, and cannot be taken up in less than twenty-four hours. By this apparatus, they not only take rays, but sharks and other fish; some of which are above a thousand pound weight. When a fish of this magnitude is found at the line, the fishermen are provided with an harpoon to dispatch them before they are brought on board.

^{*} Fide Goldsmith.

is

SECTION IV.

The Skate *.

THE skate is the largest of this genus, and thinnest in proportion to its bulk: In these colder latitudes it is seldom found above two hundred pounds weight. Willoughby mentions one sold in the market of Cambridge, which, when dressed by the cook of St. John's college, dined an hundred and twenty students †.

The noie, though not long, is sharp pointed; above the eyes there rises a set of short spines: The whole upper part of the body is a pale brown, resembling the wet sand with which the animal is said to cover itself. The body, towards the sides, is thinned down into two sins, that have the appearance of wings; and at the buttocks there are others of a thick sleshy substance. The tail of this species is short, obtuse, of a roundish form, but compressed a little towards the sides ‡.

The lower part of the body is white, plentifully marked with a great number of minute black spots, which are supposed to be the glands, by which that slimy mucus which covers the fish is secreted. All sish covered with a similar glutinous matter, are observed to possess the same organs of secretion. The skull of the skate

* Raia Batis, Lin. Syft. 'The Skate or Flaire, Will.

VOL. III.

Vide Ichthyol. page 71.

† Idem ibidem.

[§] Nicolas Steno apud Willough.

is filled with a copious collection of clear glutinous matter, through which all the nerves that are distributed to the different parts of the body, appear transparent. Among these are seen the optic and olesactory nerves; the latter proceed through the apertures, all along to the nose, and afford full proof that this animal possesses the sense of smelling.

The Sharp Nofed Skate *.

Willougher describes a sish of this species, which he examined at Rome in 1664: They are frequently taken on the British coasts; and are distinguished from the former by their long, narrow, and sharp pointed note, not unlike the end of a spontoon. The mouth is very large, and surnished with a number of small sharp teeth, hending inwards: The tail is thick, and terminates in two small sins.

This is supposed to be the bos of the ancient naturalists, a fish they describe as of an enormous size ‡. Oppian gives an account of its sondness for human sloth, and the method it takes of destroying men, by overlaying them, and keeping them down with its vast weight, till they are drowned. Other writers give the same relation, and confirm the character of these anthropophagi §. In the fouth they are so well known, that the persons employed

^{*} Raia oxyrinchus, Lin. Syst. Idem, Willoughby.

[†] Pennant's Brit. Zool.

[‡] Ευρυλαλος πανλεσσιν μετ ιχθυσιν. Haleut.

[§] Vide Phile de propriet, anim. et Ulloa's voy. i. 132.

employed in the pearl fishing, have learned from the fatal experience of many of their own number, how much they ought to be guarded against. These unhappy divers, are sometimes surrounded, and wrapped up by the fins of the manta, for so this animal is there called, till they are sufficiented: The negroes, therefore, take care never to go down without a sharp knife to defend themselves against this terible enemy.

The Torpedo, or Electric Ray ..

This species inhabits our seas, where it grows to a large size, some weighing no less than eighty pounds; its colour is a dirty clay; the head and body are round, and but indistinctly separated; the letter extremely thin, and attenuated towards the edges. Belind the eyes are two wide foramina, which Willoughby supposes, are intended for conveying sound †: they are beset with fix cutaneous rays on their inner circumference, and communicate with the mouth.

The torpedo can live only twenty-four hours out of the fea, and but little longer if put into fresh water; it inhabits these places where he bottom is fandy, and buries itself superficially in it, by slinging the fact over upon its back with a vibration it gives its extrunities.

O 2 . It

^{*} Raia Torpedo, Lin. Syft. Cramp-fish, Will,

[†] Ichthyol. page 81.

It is in this concealed fituation that the torpedo aftoniffes the unwary passenger, who inadvertently treads upon it, by the exertion of a power perhaps the most extraordinary and mysterious in nature.

The narcotic power of this animal has been taken notice of in all ages *; it is so powerful, that when the fish is alive, the instant it is touched, it benumbs the arm, and sometimes the whole body †. The same effect is produced, even when it is touched with a stick, or tread upon by a person who has his shoes on. Oppian stretches the matter probably too sar, when he alleges that it will benumb the sisherman through the whole length of the line and the rod ‡.

The shock given by this animal, most nearly resembles the shock of an electrical machine, sudden, tingling, and painful: It is thus described by Kempfer; " The inftant," fays he, "I touched it with my hand, I felt a terrible numbness in my arm, as far up as the shoulder. Those who touch it with the foot, are feized with a stronger palpitation, than those who apply to it the hand. This numbrefs bears no refemblance to that which we feel when the vein has been a long time pressed, and the foot is faid to be afleep; it rather appears like a fudden vapour, which, passing through the pores in an instant, penetrates into the very fpring of life, from whence it diffuses itself over the whole body, and gives real pain; the nerves are fo affected, that the perfon ftruck, imagines all the bones of his body, and particularly those of the limb that received the blow, are driven out of joint; all this

^{*} Vide Galen, Alian and Gillius.

[†] British Zoology,

f Haleut.

this is accompanied with an univerfal tremor, a fickness off the stomach, a general convulsion, and a total suspension of the faculties of the mind: In short, such is the pain, that all the force of our promises and authority, could not prevail upon a seaman to undergo the shock a second time *."

The substance which the torpedo discharges, and which produces effects so violent and instantaneous upon the human frame, is probably electric matter, but by what means the animal so soon collects it, no naturalist has been able to explain. Great as its powers are when the animal is in vigor, they are impaired as it declines in strength, and totally cease when it expires †; they impart no noxious qualities to the fish, when used as food; for the French, who find it in plenty on their coasts, very commonly dress and eat it ‡. Galen, we are told, recommends it in epeleptic complaints, while Rondeletius and Willoughby, decidedly condemn it in all cases, as a watery, soft, and fungous aliment §.

The narcotic power of the torpedo, however it may affect it as food, is ferviceable to the animal in two important respects; both as a means of defence against voracious fish, and a method of procuring subsistence from among the smaller tribes; the former, when electrified, are deprived of all possibility of seizing their prey; and the latter, after having unwarily approached the torpedo, and received the shock, are incapable of making their escape [].

The food of the torpedo is furmulletts and plaife; the former is fo fwift, that it is impossible for the torpedo to take

[•] Kempfer's travels. † Salvianus and Willoughb. p. 82. ‡ British Zool. § Rondel de piscib, and Willough, thethy I. p. 32. ‡ Plinii Hist. Nat.

take it by purfuit: and as this fish has been found in its stomach, it justifies Pliny's account of the manner in which this animal secures its prey. The torpedo is taken at Torbay, off Pembroke, and at Waterford in Ireland; it is caught with the other stat si h, with the trawl, and is commonly found in water forty sathom deep, in company with the congenerous rays *.

Those animals are said to bring forth their young about the time of the autumnal equinox †; but from a disfection of one made by a French gentleman, they seem capable of supersectation, and of consequence must produce young at different times. Ronacletius ‡ mentions two species of the torpedo, and Willoughby describes an American kind after Margrave, of a foot and nine inches in length, and seven in breadth. It is without teeth, and has two spiracula below the neck §.

The Fire-flare, or Sting ||.


Were we to credit all the marvellous accounts which the ancient naturalists have given, of the venom lodged in the armour of this all, we would unavoidably dread it, as an animal fill more formidable than the torpedo \(\Pi\). The weapon in which nature is faid to have lodged this poison,

^{*} British Zool. † Aristot. Hist. animal. † De piscibus. § Vide Ichthyol. p. 80.

^{||} Pastinaca marina lavis, Will. La Pastenade de mer. Belon.

I Pliny, Ellian, and Opiam.

Plate V.



LONG NOSED RAY



poison, grows like a spine from the tail; it is about five inches long, and about sour inches behind the body. According to the tremendous sables of Pliny, it is surnished with a venom so potent, as to a set even the animal creation. Trees that are struck with it, instantly lose their verdure, and rocks themselves are incapable of resulting this powerful virus. It was with the spine of the trygon, or sire-slare, that the enchantres Circa armed her son; and with this irresistible weapon, he, unintentionally, became the murderer of his sather Ulysses. What a happiness it is that these animals do not frequent the woods, and are so seldom seen ashore!

Succeeding naturalists finding the taste for the marvelous had decreased in their age, have been obliged to give the history of this fish a greater degree of verisimilitude; curtailing it of its powers against inanimate nature, they have contented themfolves with making it formidable to the fishermen to Their opinion has been fanctioned by the name of Linnaus, who afcribes poisonous qualities to no less than three different fithes; the fire-flare, the torpedo, and the tetrodon lineatus t. It may be juffly doubted, however, whether any fith has a fpine charged with actual poison; as far as the fling is concerned, we know that this is not the case: The spine is its peculiar weapon of offence, and is capable of giving a very bad wound, which being commonly inflicted on the tendonous parts of the body, and by those who are constantly working in cold water, is fometimes painful, and difficult to cure §.

But

^{*} Historia animal. - † Vide Gesner, Aldrovandus and Rondeletius † Syst. Nature. § Brit. Zool.

But the spine of the fire-flare, though unattended with any poisonous qualities, is still a formidable weapon.—
There is reason to believe, that in early times, before the use of iron, the spears and darts were headed with this bone, instead of that metal. The arrow of many of the South American tribes, are still pointed with the bone; of this sish; and from its hardness and sharpness, it makes no contemptible weapon *.

The fire-flare, formidable as it has been deemed, is never rejected by the fishermen when found in his tackle; he commonly cuts away the tail, and exposes it for sale, deprived of this instrument of mischief. This species never grows to the size of those above described; the most common sized weigh about eight or ten pounds, and are about two or three feet from the snout to the end of the tail †; the nose is very sharp at the point, but short; the mouth is small, and provided with granulated teeth; the eyes are protuberant; the irides of an oblong shape, and of the colour of gold. The whole body of the animal is smooth, slat below, but rounded above, and more elevated in the middle than any of the other rays ‡.

The fire-flare casts its spine, and renews it annually; and that it may at no time be without a proper instrument of desence, it is sometimes seen with the new one growing up, before the old drop off. That nothing relating to this animal might be dirested of the marvellous, its fish was said to have the peculiar quality of exciting concupitence; and when used in a decoction of oil, it was reckoned a cure for the leprosy: The point of its spine, when introduced into a diseased tooth, made it splinter and drop from the jaw ||.

^{*} Robertson's Hist, of South America. + Will, p. 67.

[#] Brit, Zool. || Vide Gefner.

The Thornback *.

THE thornback is the last of this genus which we shall particularise; and its figure is so well known, that a short description will suffice. It is of a rhomboidal shape, broader, however, than long, and distinguished from all the other species of the ray, by three rows of spines upon the back, and sive upon the tail, all pointing towards its end. The mouth is small, and sitted with granulated teeth; between the nose and the eyes are a few spines; others are scattered without any order upon the pectoral sins.

The colour of the upper part of the body is that of pale ashes, marked with streaks of black, and the skin rough, with small tubercles like shagreen. The belly is white, crossed with a strong semilunar cartilage †. Willoughby has given a long detail of the internal structure of this species; by which it appears, that it has two distinct organs, of a curious conformation, one on each side, for generating and excluding its eggs: In these are deposited innumerable ova of various sizes ‡; hence, the thorn-back must be very prolific. It begins to generate in the month of June, and continues production till September; its young, as long they are incapable of breeding, are called maids; they are good eating at all times of the year, but the old only begin to be in season in the month of November.

VOL. III.

P

The

^{*} Rain clavata. Lin. Syst. Idem Rondeletii et Will.

⁺ British Zool.

The fishes of this species are found in great plenty, and make a considerable article in our fish markets. They are fond of devouring herring, mackrel, and cels; but their common food is probably the different kinds of ground fish; crustaceous animals, such as the crab and lobster, have been found in their stomachs, which are supplied with a powerful solvent, capable of speedily reducing the shells into a chyle sit for their support.

SECTION V.

GEN. VII. The Shark.

HE fishes of this genus are the most formidable and voracious of all the tyrants of the ocean; if some of the rays were invested by the credulity of the ancients, with imaginary powers of destruction, those possessed by the shark are real. The spine of the fire-flare might often prove hurtful, but the fangs of the fhark are always fatal. The genus is distinguished by an enormous mouth, placed beneath the head, armed both above and below, with an hundred and forty-four frightful teeth, which the animal has the power to raife or depress at pleasure; to this hideous apparatus of destruction, the shark joins a fierceness and rapidity, far superior to that of the other rapacious tribes: The fmallest of the race is dreaded by almost every fish in the sea; for there is hardly any of them that dread an encounter with animals far fuperior in fize, and more formidable in appearance. The white shark, which is the largest of this tremendous race, maintains a terrible despotism in the ocean; for, with that amazing velocity peculiar to his tribe, he unites the flrongest appetite for depredation. He approaches nearly to the five of the whale, but far furpailes that animal in strength and colority, in the formidable arrangement of his teeth, and in his insatiable desire of plunder *.

The head and throat of the shark are larger than the proportion to the body, which is slender and tapered away almost to a point as it approaches the tail. The teeth are broad at the base, but slender and very sharp above; they are disposed in sive rows all around each jaw, and at their insertion are surnished with strong museles, by means of which the animal raises or depresses them; when the shark is at rest, they all lie slat in his mouth; but when he comes forward to lay hold of his prey, these dreadful instruments are suddenly raised, and the devoted animal, seized by them, instantly dies, pierced with an hundred wounds.

The animals of this genus have a wild and ferocious afpect, that faithfully indicates the malignity of their nature: Their eyes are oblong, funk in the fockets, and placed longitudinally in the head, where they appear almost overhung by the skin, and are full of malevolence and sire: The fins, to which they owe their rapidity of motion, are larger, and more numerous than those of most other sithes: The body, though slender, is possessed of very strong muscles, the source of that assonishing impulse communicated by the tail. The whole external parts are covered with a rough and hard skin, which was used by the ancients in polishing wood and ivory; but is now applied to cover instrument cases, the skin of the greater dog sith furnishing a preserable shagreen †.

The

^{*} Goldsmith's Nat. Hist. vol. vi. 238.

at British Zoology, Class iv. gen. 6;

The shark thus fitted by his external conformation for rapin and bloodshed, is endowed also by nature with a courage and activity that gives effect to his rapacious inflinds. No fish, it is faid, can swim as fast as he; none is so constantly employed in swimming; scarcely a ship croffes the Atlantic, that is not attended by some of these rapacious monsters, for days together; and fuch is their velocity, that they outstrip the best sailing vessel, and play around her without any fymptoms of exertion. With fuch voracious appetites, and fach powers of gratifying them, one would imagine that thele animals might have long fince destroyed the inferior tribes, and thinned even the ocean itself of its inhabitants; but happily for the fafety of the smaller fitnes, these plunderers, by the particular fituation of their mouths, are obliged to turn upon their fide before they can feize their food; and while this evolution is performing, their prey often finds time to escape *.

But although the devastations committed by the shark, are thus providentially limited by the aukward situation of his mouth, still his depredations are extensive, and in every warm climate he is the terror of the sailors; for of all kinds of food, he is said to discover the strongest predilection for human stell. If by any accidence a seamon has sallen a victim to this destroyer, he is seen for some time afterwards havesing about the same place, and making several returns to the space expecting another repast of the same booty: The wake of a ship is a sevenite refort with those mimals, and numberless are the instances in which accident, or the improvident rathness of seamen,

have favoured their defigns upon the human race. Allong the coast of Africa, where the sharks are numerous, many of the negroes, it is faid, who are obliged to fish for their sustenance, are annually devoured by them. The inhabitants of these dreary regions are sirmly of opinion, that the shark prefers the slesh of a black man to that of a white; and that in consequence of this unenviable diffinction, if two persons of different colours are both in the water together, his choice falls always upon the negro *.


Whether there be any foundation for this opinion of the negroes, it is of little importance to inquire; that the shark discovers an avidity for this species of food, is a fact that has been too often verified by fatal experience, to admit of a doubt. Mr. Pennant relates the story of the master of a Guinea ship, who, finding a rage for suicide prevail among his flaves, from a notion the unhappy creatures entertained, that after death they would be reflored to the families, their friends, and their country; to convince them, that at least some difgrace should attend them here, ordered one of their dead bodies to be tied by the heels to a rope, and fo let down into the fea: Though it was drawn up again with great fwiftness, yet in that flort space the sharks had bit off all but the feet +. Gold/mith tells another, occasioned by the same barbarous traffic, of a woman who was thrown out alive, to deter the flaves from felf murder: When the rope was drawn to pull her up into the veffel, part of her body appeared in circumflances too shocking to be mentioned.

^{*} Idem ibidem,

[†] British Zoology.



Plate VI.



SECTION VIL.

The Angel *.

This fish possesses the character of extreme voracity, common to the sharks and rays, two genera which nature seems to have intended this animal should, in some measure, connect together: In its external sigure, it partakes of the nature of both, while it differs from each in the situation of the mouth; that organ being placed in the extremity of the snout, and not below the head †.

The angel grows frequently to the weight of an hundred and fixty pounds: Its body is covered over with a filthy mucus, beneath which lies a rough skin, employed formerly by the workmen as a shagreen ‡. It was from this circumstance, that the fish obtained the name of file among the Greeks §; while the Romans called it fquatina, from the filth with which the body is covered ||. The colour of the back and sides is brownish, that of the belly white. The pectoral sins are large, and seem, like those of the rays, to be a prolongation of the body: It was from these wing-like sins that this fish received its English name ¶.

The

^{*} Squalus Squatina, Lin. Syst. L'Ange, ou Angelot de Mer, Belon.

[†] Brit. Zool. ‡ Plinii. Lib. ix. Cap. 12.

[§] Pan, Arift. Hift. Anim.

[|] Squatina, Vide Willough. page 80.

I Idem ibidem.

The ventral fins are placed in the same manner as the pectoral, and to them is annexed on each side the double penis, that characterises this and the former genus. Upon the back, and farther down than the anus, there are two dorfal fins: The tail is biturcated, the upper lobe rather larger than the under; a character belonging to all the sharks.

This species is described by Rondeletius as an inhabitant of the Mediterranean; it is also seen frequently on the British coasts, where it preys voraciously upon the statish that frequents the bottom of the water. The slesh is in no request, but seems loathed in every country, on account of its rank flavour and hardness: Its ova, when dried, are used for medicinal purposes, and are supposed by the Italian sinermen to be efficacious in removing a diarrhea*. The Turks upon the Barbary coast, apply the skin in the manusacture of sheaths for swords, which they are said to sabricate with peculiar neatness †. None of the uses, however, in which the parts of this malignant monster are employed, seem capable of compensating the dangers to which seamen are continually exposed, from its voracious habits.

^{*} Rondeletius de piscibus.

The Basking Shark *:

This is the largest animal of this genus, and differs in some remarkable characters from the rest of the tribe; it inhabits chiefly the cold climates, and has long been known to the inhabitants of the west of Scotland and Ireland, where it is distinguished among the sailors by the name of the basking shark, though better known in the former country by the name of the Sail-sish.

The latter of these names has been given it, because it is seen at a distance on the top of the water, with all its dorsal fins appearing above the surface, and spread like sails. Both in its manners, and external form, there are peculiarities in which it dissers from the other sharks. It is far from being either voracious or sierce: Its food consists chiefly of sea plants, no sish having been ever found in its bowels; but the green, and half digested remains of algre: As its habits resemble those of the whale, Linneus supposes that it is supported by the same food, afferting that it devours only meduse.

Mr. Pennant has given a very accurate description of this species, and seems first to have discovered its affinity to the sharks, by pointing out the sive transverse apertures of the gills on each side of the neck: formerly is had been considered as a kind of whale ‡. The oil arising from their capture, is now made an article of trade in Vol. III.

Squalus Maximus, Lin. Syst. Sun-fish, Smith's Hist. of Corke, † Systema Nat.
‡ British Zool. class iv. gen. vi.

the west of Scotland, where we have seen from fix to eight barrels taken from one fish, though it is only the liver that produces it. They lie quiet and motionless on the furface, commonly on their bellies, but fometimes like tired fwimmers, upon their back; and it is in this pofture that they allow a boat to approach them, fometimes till it is within contact, without attempting to escape. Then the harpooner strikes his weapon into the body, as near as possible to the gills: But so insensible are they to the wound, that they do not feel its pain, till, by the united strength of two men, the instrument is forced very deep into their flesh. As foon as they feel themfelves wounded, they fling up their tail, and dive to the bottom. Frequently in their agonies, they coil the rope about them, attempting to difengage the harpoon by rolling upon the ground.

Having at length discovered that all efforts of this kind are vain, they swim away with amazing rapidity, and with such violence, that a vessel of seventy tons has been towed after them against a fresh gale. They sometimes run off with two hundred fathoms of line, and with two harpoons in them; and will employ the sishers for twelve, and sometimes twenty-sour hours before they are subdued. When killed, they are either hawled on shore, or, if at a distance from land, to the vessel side; where the liver is taken out and melted into oil, in kettles provided for that purpose *.

The basking shark is from fifteen to thirty feet in length; the colour above of a leaden colour, and below a pure white; the form of the body, like that of the shark kind, is slender and tapering. The upper jaw projects

^{*} Vide Pennant ubi fupra.

jects beyond the lower, and is rounded at the end; the mouth is placed beneath, and furnished with a vast number of small teeth. The dorsal sins are placed differently from those of the rest of the genus: the first rises nearer the head than the tail; the second is much smaller, and is situated near the extremity. On the under part of the body there are sive sins; two pectoral, two ventral, placed below the second sin of the back, and a small annal sin. Near these the male animal is provided with a double organ of generation; and between them in the semale, is situated the pudendum. The oil produced by these animals, is said to be preferable to every other for the manufacture of wool; and the skin furnishes worksmen with an excellent shagreen.

The White Shark *.

This shark is probably the most ravenous and formidable of the order, and grows to the immense weight of four thousand pounds †. It was in the body of one of this species that an entire corpse was found; a circumstance no way incredible, when we consider their strong appetite for human sless, and the amazing power of deglutition which an animal of this size must possess. In all hot climates, it is this species in particular that is the terror of the sailors; it is they that constantly attend the ships,

* Squalus Carcharias, Lin. Syst. Lamia, Plinii Hist. Nat.

⁴ Gillius.

watching till fome failor drop overboard, or attempt to bathe; in either case, the unfortunate seaman perishes without redemption; for this voracious creature, will even dart up above the water to meet him in his fall.

It is in this fish that Rondeletius places the prophet Jonas, when he lay so long unhurt in bowels; and he says, that the language of Scripture gives no repugnance to his idea; for among the ancients all large sish whatever were termed whales *.

The mouth of this species is proportioned to the immense bulk of the animal: it is furnished with fix rows of teeth, the number of which probably varies, according to the age of the fish. Fossil teeth of the white shark are frequently dug up in Malta, and fometimes in England: in the former country, they have been feen four inches long, which must have belonged to animals of vast fize, if the proportions laid down by Grew, between the body and the teeth of this species, are just +: He afferts, that those in the jaw of a shark of two yards long, are not above half an inch; the fossil teeth found in Malta would, according to this calculation, belong to an animal fixteen yards in length Linnaus has enumerated fifteen different species of sharks; but if the appearances of the teeth, shewn in the collection of the curious, can be trusted to, he is far short of their actual number; for many kinds are feen, of which no mention is made by him.

Pliny, Oppian, and feveral of the ancient naturalists, were acquainted with this species, whose mouth they justly describe as situated far beneath the head, obliging the animal to turn almost upon its back, when it devours

its prey*. The eyes are large, round, and staring, capable of being easily rolled in the fockets. The dorsal, and pectoral fins are very large, and aid the efforts of the tail, in giving rapidity to the fish's motion. The tail itself is semilunar, and capable of giving the most dreadful strokes; hence, the sailors, as soon as they take one of these fishes, so far duarm it, by cutting off the tail.

The Blue Shark +.

This species has obtained its name from the fine azure colour of the back; it is most frequently an inhabitant of the hot climates, where it is scarcely less formidable to seamen, than that immediately preceding. The body is less rough than those of the rest of the genus. The eyes are of an eliptical shape, but the irides perfectly round; behind them, there are none of those foramina, which are observed in other species ‡.

The fish is of an oblong shape; the nose long, sharp, and compressed towards the point; the nostrils are placed transversely. Although most frequent in the warm climates, the blue shark is often seen on the British coasts, and particularly those of Cornwall, during the pilchard feason. It is there caught by the sishers, with a large kind of iron hooks, that are made for the purpose.

y ondeletius

Supina vescantur. Plinii. Lib. ix. cap 24.

⁺ Squalus glaucus, Lin. Syft. Galeus glaucus, Rondel.

i Willough. page 49.

Rondeletius ascribes to this species, only two rows of teeth; but their number probably depends upon the sishes age; and if so, no conclusion can be drawn with regard to it, from a single instance *: They are granulated, and of a triangular shape †. This naturalist declares, that he was an eye witness of the blue sharks fondness for human sless; having seen one pursue a boy a considerable way along the shore, and making a spring at him, nearly bit of his legs.

It is concerning this species, that *Ælian* relates such extraordinary instances of parental affection ‡. It is said by him, that the blue shark permits its small brood, when in danger, to swim down its mouth, and take shelter in its belly. This sact is said to have been observed by *Rondeletius*, an able naturalist; and it is narrated by *Pennant*, a very accurate ichthyologist, as a part of the history of this whole genus that is entitled to belief. In sact, it is no more incredible, than that the young of the opposium, should seek an assylum in the ventral pouch of its parent; a sact too well known to be contested §.


[&]quot; De Piscibus.

[†] Lin. Syst. Nat,

[‡] Ælai. An lib. i. cap. 16.

[§] Vide British Zool, article, Blue Shark.





The Long Tailed Shark, or Sea Ape *.

THIS species has obtained its name from the extraordinary length of its tail, which generally exceeds that of the body, the upper lobe extending greatly beyond the lower, almost in a straight line. Its fish has an abominable fetid, and rank fmell, refembling that of the land fox + The body is more thick and round than any of the sharks; the snout is sharp; and a little below is a fmall mouth, furnished with pointed teeth. The colour above is cinereous, below white. Rondeletius afferts, that he was an eye witness of this animal allowing its young to retreat into its belly to avoid danger; for he faw an old one diffected upon the shore, in the body of which were all its young. At first he imagined that they had been devoured by the parent in place of food; but upon examining them, they were all alive, and fo entirely unhurt, that there could remain hardly any doubt of their having fled there as a refuge from danger t. The extraordinary length of the tails of the young, forms no objection to this fact; for at that age they are foft and pliant. Willoughby supposes that this fish was only found in the Mediterranean; it has fince, however, been taken in the British feas §.

The

^{*} Vulpes Marina, Rond. Sea-fox or Ape, Willough...

[†] Salvianus. ‡ De Piscibus.

[§] Pennant's British Zoology.

The other fishes of this genus that frequent our coasts, are the tope, or sea dog of the ancient naturalists; the greater and lesser spotted sharks; the smooth shark; the porbeagle, and the picked shark: These all so nearly correspond in their general characters to those already described, that there seems no occasion farther to particularize them.

SECTION VI.

GENUS VIII. The Angler.

The fishes of this genus were by the ancients called sea frogs *, from their resemblance to that animal: The English have given them a still more hideous name, that of sea toads, or sea devils; appellations which these animals seem to merit, by their extraordinary deformity. Of all the monsters contained in the ocean, they are perhaps the most ugly and deformed. Their head and body are joined together, forming one round slat mass †. The head and mouth are far larger than all the rest of the body; the latter being in some a yard wide, and surnished all around with an infinite number of small teeth. The pectoral sins are broad and thick; behind each venetral sin there is one aperture.

Vol. III. R

^{*} Βατραχος, Aristot. Oppian. Rana piscatrix, Pliny and Ovid.

[†] Willough. Ichtchyol. page 83.

The Common Angler *.

It is in its tadpole state that this animal resembles the frog: its colour upon the back is dark blue, marbled with some white spots. The lower jaw projects considerably beyond the upper, which gives the animal a grinning aspect. The eyes are placed in the upper superficies of the head, somewhat prominent, and their irides white. About half an inch from the corner of the upper jaw are two hairs, the one sour, and the other six inches in length. It is by means of these, according to Pliny, that this animal sisses; it puts out these shender horns among the mud, enticing the little sish to play around it till they come within reach, when it springs upon them, and devours them †.

Around the edges of its head are small fin-like appendages; outwardly there appear no spertures of the nose, but there are two within the mouth, supposed to convey smell ‡. About the middle of the back there rises a fin supported by ten rays; the tail is not forked, but the rays stretch somewhat beyond the membranes of the fin. The ventral sins are broad, thick, and sleshy, jointed like arms, and on the inner side divided into singers, resembling what is seen on the foot of a mole §. With these instruments

²

^{*} Lophius piscatorius, Lin. Syst. La Grenouille de Mer. Belon.

[†] Plinii, lib. ix. Vide etiam Cicero de Nat. Deor.

[#] Willough. ubi supra. § British Zool, and Willough.

instruments Belon afferts, that the animal walks at the bottom of the water.

Another pair of fins is placed farther out, at the verge of the body, and below there are the apertures of the gills, differing so considerably from those of other fish, that they have been supposed by some, to be a retreat for the young. They are large orisices, and at the bottom appears the branchiæ, resembling the teeth of a comb.

These fishes grow to an extraordinary fize; some are taken near Scarborough, between four and five feet in length, and whose mouth are near three seet wide. The fishermen no sooner take them, than they give them their liberty, from the supposition that they destroy the dog sish; and in support of that idea, some of these voracious animals have been found in their stomachs.

Willoughby has four different species, which he has classed under this genus; three of which are Brazilian sishes, described by Margrave, the Guacucuia, the Guapperua, and the Acaramucu: Of these the Guacucuia or Brazilian water bat, resembles our sishing frog, in the shape of its body and the fins; but differs from it in the smallness of the mouth, in wanting teeth, and in the long protuberant snout, that resembles a horn. The whole body is variegated with white and black spots, of which the pencil, and much less the pen, is unable to convey any adequate idea *.

R 2

Willough, page 89.

GENUS IX .- The Sturgeon.

The fishes of this genus have only one narrow aperture on each fide; a circumstance in which they resemble the anglers above described. They have the mouth placed far below, tubular, and unprovided with either teeth or jaws *: Hence, these animals collect their food by suction; and notwithstanding their fize, which is often as large as that of the shark, they are the most peaceable and harmless of all animals. The sturgeon is as unwilling, as it is incapable to injure other fish: It slies before the most feeble, and timid of the race; and not unsrequently falls a victim to its own apprehensions and fears †.


The sturgeons are gregarious and migratory: To the former of these habits, they are probably compelled by the timidity of their nature, and their incapacity for self defence: all the weaker animals that are peaceable and inossensive, commonly trust to their numbers for security; and hence, herd in great crowds together. The necessity of procuring food, of avoiding cold, or of providing a proper place for the production of their young, determines

British Zool.

[†] Goldsmith's Nat. Hist.



Plate VIII.



determines the migrations of the sturgeon, as well as those of other sish. During the winter months they mount the rivers; while in summer, they betake themselves to the ocean; probably, however, never removing at any great distance from the shore *.

The Common Sturgeon +.

RONDELETIUS, Gefner, and Aldrovandus, have tedious disputations, whether this animal be the acipenser, or Elops of the ancients; a matter of very little moment, since, if the animal be accurately described, and well known, it matters but little by what name it was anciently distinguished.

The great refort of the sturgeons is the rivers of Russiand Germany; and as they are frequently found in the Mediterranean, at the mouths of the Italian rivers, it is highly probable that at least some species of them were known to the Romans: Pliny, however, as well as Ovid, mention the acipenser as a foreign fish ‡.

The common sturgeon is an inhabitant of the British feas, from which it ascends the rivers, and is sometimes taken in the nets set for salmon: It is there, however, sound only in small numbers; the greater part consumed in this island is brought either from North America, or the Baltic. In the lakes of Frischehaff, and Gurrischaff, and in the rivers in the neighbourhood of Pillau, they

Willough. Ichthyol.

[†] Acipenfer Sturio, Lin. Syft, Acipenfer, Rondelet.

Pliny, Lib. ix. cap. 17. et Ovidii Halleut.

are taken during the summer months in great abundance, and sold over all Europe at a very high price. The shores in that vicinity are all divided into certain districts, and let as we do the salmon sisheries; some of them bring a rent of six thousand guilders, or near three hundred pounds per annum *.

In these large rivers they are fometimes found of two hundred pounds weight, and are taken in nets made of fmall cords. They are dull and spiritless animals, making but small resistance when entangled; and as soon as they are brought out of the water, appear like a lifeless lump. Their flesh is every where in high estimation; it has the delicacy, whiteness, and folidity of veal, and when roafted, is reckoned one of the greatest niceties that can provoke the appetite of an epicure +. It is commonly pickled, and packed up in casks, when fent to foreign markets: Of late, we have received a confiderable quantity of these fish from North America, where they are caught in May, June, and July; it is at that feafon that they are feen leaping above the furface in the rivers, and falling again upon their fides, with a noise that is heard at the distance of some miles t.

It is of the ova of the sturgeon that the Russians make the caviar, by taking out the membranes found among them, and drying them, after having been washed with vinegar. The best caviar is said to be afforded by a small species called the sterlet, sound in the Yaik and Volga §.

Icthyocolla or ifinglass is made of the sound of this as well as the other kinds of sturgeon ||. The ancients were acquainted

^{*} Brit. Zool. † Will. p. 249.

¹ Catefby Nat. Hift. of Carolina.

[§] Strahlenberg's Hist. of Russia, p. 337.

[|] Vide Phil. Trauf. lvii. 354.

acquainted with the composition of this glue, so useful in many species of manufacture: It was probably made up by them in a similar manner, from a sish that had this very name *: It is useful as a glue for paper, musical instruments, and cementing every slender sabric of wood. It is well known to the varnisher, the apothecary, and the clothier.

The fize of the sturgeon depends upon the place of its refidence; where it is confined either in fresh or falt water, it is comparatively fmall; but where it alternately enjoys both, it grows often to the length of eighteen feet, and weighs from four to five hundred pounds; of this magnitude one was caught in the river Esk, in Britain; and in the larger rivers of the continent, some attain even to a superior fize to. The body of this fish is long, and of a pentagonal form, on account of five rows of large bony scales, which proceed from the head along as far as the dorfal fin, one upon the back, and two upon each fide. The head is fmall and protuberant at the fnout; the eyes are fmall in proportion to the fish, and the irides are of a filver grey: The mouth, without either jaws or teeth, is fituated confiderably below the extremity of the nofe, and in the middle space between them, spring out a few bristles: The first pair of fins are placed immediately befide the gills; the fecond pair near the anus, and a third between that and the tail; upon the back their is but one fin.

The sturgeon differs from the other cartilaginous sistes in the manner of its generation; like the spinous tribes it is oviparous, and deposits its spawn in the winter months ‡.

^{*} Plinii, Lib. xxxii. cap 7.

[†] Rondeletius.

⁴ Brit. Zool.

The Hufo of Germany.

This species is supposed to be peculiar to the rivers of Germany; particularly the Danube and those streams that slow into it. From the Black Sea it ascends that river, and is caught in nets placed in different places to intercept it, nearly as far up as Vienna, where they are exposed to sale every Friday, to the number of fixty or an hundred. The siesh is soft, glutinous, and slabby; but it bears falting, and is exported from Wallachia, where it is principally taken, to the different markets of Europe*.

This fish, like the rest of the genus, is gregarious and migratory; and it is upon its excursions from the sea to the fresh water, that it is taken, during the months of October and November. The husos are from fifty to sour hundred pounds weight, and are said to possess such strength, that they frequently drive the sisherman overboard with a stroke of the tail †.

It is chiefly for the ifinglass which it affords, that this fish is taken: The manner of preparing this substance, is by cutting the skin, the entrails and the fins, into small pieces, and leaving them to macerate in a proper quantity of warm water; they are afterwards all boiled together upon a flow fire, until they are disloved and reduced to a jelly; this jelly is spread upon instruments made for the purpose, and upon drying it assumes the form of parchment;

^{*} Gesner de Aquatil. p. 59. † Idem ibidem:

parchment; when it is rolled up, and exported in the shape in which we see it in the shops of the venders *.

The fnout of this species is remarkably long, and surnished below with small barbs; on the belly are two pair of sins, on the back one; above, the body is black; the lower parts yellow, as in the perch; the whole surface is destitute of sins, and the slesh is supported by cartilages instead of bones. The genuine icthyocolla that produces the isinglass is supposed by Willoughby, to be a dissertent sish from the huso.

Vol. III.

S

^{*} Goldfmith's Nat. Hift.

[†] Vide D. Will. apud F. Will. Ichthyol. p. 243.

SECTION VII.

GENUS X .- The Diadon.

THE cartilaginous class of fishes abounds with animals of anomalous and monstrous figures, which no pen can describe, and which even the pencil of the artist cannot imitate; fuch are the endless variety of orbs, of triangular and quadrangular fish described by Clusius, and after him by Willoughby *. They are mostly exotic animals, exhibiting the rich exuberance of nature on the warmer coasts of America, India, and the Mediterranean. From all these hideous animals, the British seas are happily free: Their utility to man cannot be discovered; their deformity is obvious, and the venemous nature of many of them has been fo often confirmed by fatal experience, That it can no longer be doubted. From the distance of their fituation, as well as their unattractive form, their history is but little known; we shall select, therefore, a few only for a particular discussion.

^{*} Ichthyologia, p. 143

The Globe *.

This fish is feldom a visitor of the British coast; one, however, of the species was taken at Penzance in Cornwall. Its more common residence is off the shores of Carolina. It is of an oblong shape, of a foot and eight inches in length; the length of the globular belly, when extended, is above one foot, and its circumference, when in that state, is above two: This singular power of instating its belly, till it acquires a globular shape of such magnitude, seems to have been intended by Nature as the defence of this animal; it is then of a size less easily laid hold of by its enemies, whilst the great number of spines with which its belly is surrounded, renders it an object of terror *.

The globe, upon the back, is of a rich deep blue; the fides and belly are white, shagreened or wrinkled, and beset with innumerable small spines, each attached to she skin by four processes. It has two pectoral sins; a dorsal one far back behind the globe, and opposite to that an anal sin.

Tetrodon lævigatus Lin, Syst. † Pennant. Class iv. gen. 9.

The Oblong Diadon *.

This species disters from the former, not only in wanting the globe, but in being much shorter; it resembles a carp, or some thick sish cut through by the middle. It sometimes grows to a very amazing size; some have been taken sive hundred pounds weight †. It is commonly little more than two seet in length; so that the thickness, to afford nearly such a weight, must be very much disproportioned to the animal's size.


The body of this species is black above, and dappled; below it is silvery; it wants scales, but is covered with a thick, hard, and rough skin. The mouth is small for the size of the sish; between it and the eyes there are two apertures for the nostrils; but there are several other oringes in the head, the use of which has never been ascertained by any naturalist, though Willoughby supposes that two of them correspond to the organs of hearing in other animals ‡.

The pectoral fins are placed immediately behind the eyes; they are fmall, and of a roundish shape; the dorfal and the anal fins are placed high, and at the very extremity of the body; the tail fin is semicircular, and fills up the whole abrupt space between them.

^{*} Oftracion Lævis, Gronovius. † Borlafes's Hift, Cornwall, † Ichthylogia, p. 151.



Plate IX.



LUMP SUCKER







JURA SUCKER

The short diadon is still a more aukward figure than this; it is shorter and thicker, having the anal and dorfal sins set still higher, which gives the body a more abrupt appearance. The slesh of both these is rank and unsavoury: When boiled, it turns into a kind of glutinous jelly, resembling boiled starch after it is cold; and from experiments made upon leather and paper, it has been found to answer the purposes of glue *. Linnaus has enumerated seven species belonging to this genus †.

? British Zoology.

A Syft. Nat. 414.

GENUS XI .- The Sucker.

The Lump *.

It is tribe contains only three species, all remarkable for the singular desormity of their shape. The body is deep and thick; the back arched, and the belly distinguished by an oval aperture, surrounded with a sleshy and muscular substance, and edged with small thready appendages, which operate as so many claspers. It is by means of this appartus that these animals adhere with such vast force to any substance to which they apply themselves. The tenacity, with which the sucker adheres to the bottom, has been tried by putting it into a pail of water, where it sixed itself so firmly, that on taking the sish by the tail, the whole vessel, with its contents, were listed from the ground, although it held some gallons †.

The lump sucker, which is the largest of this genus, increases to the weight of seven pounds, and the length of nine inches. The back is arched and sharp, of a black-ish colour, dilated and variegated with brown. The belly is flat and reddish: it wants fins, but the whole body is defended with sharp black tubercles. On each side there are three rows of large bony scales, and on the back

Lumpus Anglorum, Will.

back a fourth. It has two pectoral, a dorfal, and an anal fin; the eyes are covered with a lax skin, the irides of a pale red; the lips are thick, fat, and ruddy; the orifices of the nose appear one on each side, raised above the surface of the rest of the skin*.

The lump fish is caught on several parts of the British coasts; and though its flesh is held in no great estimation. it is frequently feen for fale in the London market. The great refort of this species, is the north sea upon the coasts of Greenland. While it prowls upon the top, the feals fwarm beneath, watching an opportunity to devour it. Infinite numbers fall a prey to these animals, who swallow all but the skin; vast quantities of which, thus emptied by the feals, are feen floating above during the fpring months, when the fuckers approach near the land, for the purpofe of depositing their spawn. Every district where the seals carry on their depredation, is eafily diftinguished by the failors, on account of the fmoothness of the water: Wherever any oily fish has been devoured below, the water at the top becomes tranquil, it being uniformly found to be a property of oil to calm the agitation of the waves t.

The inhabitants of the barren tracts of Greenland, who are obliged to draw a great part of their fustenance from the sea, avail themselves also of this sish; the roe is remarkably large, and in the summer months they take them on account of it; when boiled, it forms an extremely sat, oily sood; a kind of repast, of which the necessities of the Greenlanders have rendered them exceedingly fond ‡.

Willough. page 208. † Phil. Trans. 1774. p. 445-

[‡] Crantz's Hist. Greenland.

The Sea Snail *.

THE whole of this fish, the head as well as body, is fore and unctuous, easily soluble into a kind of oily sluid; it is on this account that it has obtained the name of snail. Its habitation is not properly in the sea, but about four or sive miles from the mouths of large rivers, where the water is beginning to be salt. The body of this species is transparent, of about sive inches in length; and when newly taken, the colour is pale brown †.

The head is thick and round; the mouth without teeth, each jaw being only a little rough. The aperture of the gills is fmall, covered with a valve or operculum, which fprings from the base of each of the pectoral sins. These sins are very broad, thin and transparent; and below the throat, they almost unite together. Below the throat, there is observed a round spot resembling the impression of a seal, the place which the animal applies to those substances to which it means to adhere. The anal and dorsal sins continue without interruption, till, like those of the eel, they meet at the tail ‡.


^{*} Liparis nostras. Will. Cyclopterus liparis. Lin. Syst.

[†] Will. Append. 17.


[#] Brit. Zool.



Plate X



SEA PORCUPINES



OSTRACION



GENUS XII .- The Sea Forcupine *.

This animal is placed among the orbs or globular fift, by Willoughby; of all the inhabitants of the fea, it is the most completely furnished by nature with defensive armour. Instead of scales, the whole skin is covered with strong sharp spines, which are firmly inserted into it at the base. These are all laid slat when the animal is at rest; but when alarmed, they all around stand upon end, so that the sish is inaccessible upon every quarter. There are various kinds of these animals; all, however, seem equally possessed of the power of raising or depressing their spines at pleasure, and by that means of increasing the terror of their appearance in proportion to the approach of danger.

Together with the power of erecting their spines, they possess also that of instating their bodies on the appearance of an enemy. This operation is performed by means of a bag within the body, which, by being filled with air, instates the fish to a prodigious size: An animal of this tribe, that at first view seems small and inossensive, is no sooner alarmed or provoked, than it swells to the view, the whole body becoming visibly rounder and larger; while in the mean time all its prickles stand upright, and threaten the invader on every side †. The Americans often amuse themselves with the barren pleasure of catching these frightful creatures, by a line and hook baited with

Vol. III. T a piece

^{*} Hiftrix, Clufii Exot. Hiftrix, Lin. Syft.

d Goldsmith's Nat. Hift.

a piece of fea crab: The animal approaches the hook with its fpines flattened; but when hooked and flopped by the line, all its fpears are inflantly crected, and the whole body is armed fo completely at all points, that it is impossible any where to lay hold of it; it is therefore dragged to fome distance from the water, and there it quickly expires.

The body of this species is about twenty inches long, and twenty-nine in circumference cound the thickest part. The mouth is a little prominent, and when opened, about three inches in diameter; the eyes are pretty large, and rendered horrible in their aspect, by four spines vising immediately above each of them. The two pectoral sins are about three inches long, and sive broad; the dorsal and anal sins are nearly of the same dimensions.

⁹ Vide Clufium apud Will, p. 140.

GENUS XIII .- Ofracion *.

Ostracton is the generic name adopted by Linnaus for nine different kinds of anomalous fish which he has enumerated, all inhabitants of warm climates, and all agreeing in having the body covered with a bony coat of mail †. Some of them want horns; others have two, and some four: They are of various shapes and sizes, and from their appearance, hardly seem entitled to rank with this class of the animal kingdom.

The species here described was fent from the East Indies, and is about fifteen inches long, and four deep: The forehead appears flat, and even concave, on account of two large eye-brows that preject over it; the eyes are about an inch in diameter; below them are the two fmall foramina of the nostrils; the mouth is hardly an inch in width, and furnished with very minute teeth. The body is of a quadrangular shape, and moved by five fins; of these, the two branchial are square, being two inches broad by two long: Refembling thefe, are other two fituated near the tail, the one above and the other below; the tail fin is three inches in length, by three and a half in breadth. The whole body is covered with a crustaceous substance, refembling that of the pale blooded fishes of that order. This crust is strengthened and ornamented by a great T 2 number


Oftracion prior Aldrovandia

[†] Systema Naturæ, genus 136.

number of tubercles of triangular shape, and in substance is formething of the nature, partly of skin, and partly of a shell. There are some varieties of this species *. Of the orbs, or globular sish, Willoughby enumerates no less than thirty different species; concerning which, however, very little is known, but the name and the external sigure: His orbs comprehend two or three different genera of the Systema Nature. Linneus having improved the arrangement given by preceding naturalists.

^{*} Will. Ichthyol. p. 148.





GEN. XIV .- The Pipe Fish.

This genus comprehends all those fishes that are known by the name of sea adders, and sea horse: Linneus has enumerated seven different species, of which the generic characters are, a long and tubular nose, no orifice to the gills, but the breathing aperture situated on the back part of the head; the body is covered with a strong crust; and in all, the ventral fins are wanting *.

The Longer Pipe Fish +.

This species, as described by Sir Robert Sibbald ‡, is two feet long, the nose an inch in length, and compressed laterally; at the end of it is a small orifice for the mouth. The thickest part of the body did not exceed the circumference of a swan's quill; trexangular from the head to the dorsal fin, the rest quadrangular. The belly, carinated, and marked along the middle with a dusky line §.

The

^{*} Vide Systema Nat. † Syngnathus barbarus, Lin. Syst. ‡ Scotia illustrata, 24. Tab. 19. § British Zool.

The colour of this fpecies is an olive brown, marked with bluish lines, and, toward the tail, with brown spots. The pectoral sins have each twelve rays, the anal two, and the dorfal, which was two inches long, has forty. Between the vent and the tail, is a groove six inches and an half long, covered by two valves. This is the receptacle for the young; and as these animals are viviparous, on treading upon this part, hundreds of living young, are seen to creep out around the parent sish.

There are three species of the pipe sish sound in the British seas; the longer, above described, the shorter, and the small pipe sish, or ophidion of Linnæus. The last seldem exceeds sive inches in length, is very slender, and tapers off to a point at the tail, having no sin there. The pectoral sins are also wanting, and the body is covered with a smooth skin, instead of the crust peculiar to this genus. This species is not viviparous, and consequently the groove under the belly, does not contain living young, but eggs, which are large, and not very numerous.

The Hippocampus, or Sea Horife *.

I'ms animal has obtained its name from the resemblance its head bears to that of a horse +: It bends the body into different curvatures, like an eel; and is marked all over with circular incisions, resembling those of the insect tribe.

^{*} Hippocampus, Lin. Syst. Cavallo. Marino Italor

[†] Gesner de piscibus.

tribe. It is only nine inches in length, and often lefs, while it: thickness is not above an inch: The fnout is oblong, without any suffere at the mouth; at the lower end of which appear the foramina, with an operculum to cover them*. The body is furnished with acculeated knobs, and is of a pentagonal shape, till it reaches the anus, from which, to the tail, it is quadrangular. The eyes are round and prominent; behind them, in place of the gills, which are here wauting, there are two small sins, resembling the external ears of terrestrial animals. Under the belly there are two crisices; one for voiding the excrement, and the other for the emission of the young in their egg state.

The colour of this animal is a dark green, the belly variegated with light blue spots. In the middle of the back rifes one dorfal fin, with fixteen rays. Some species have hair along the back, refembling the mane of a horse. They are caught in different parts; that which Willoughby describes, he saw at Baic, not far from Naples. It had no mane; he concludes, that that ornament is either characteristic of one of the sexes, or of a different species. In his plates, he has given the sigure of three different kinds †.

^{*} Willough. page 157.

[†] Vide Willough, Ichthyol, p. 158.

CHAPTER VI.

ORDER III .- Spinous Fishes.

SECTION I.

General Characters of this Order.

WE are now come to that order of the finny tribe, to which the later naturalitts have exclusively appropriated the name of fish: According to them, the cetaceous kinds are beafts, that have taken up their abode in the ocean; and the cartilaginous, an amphibious band that are but half denizens of that element; the fpinous fishes being aione entitled to that appellation *. After detailing the history of the two first orders, we have seen that there are particularities in their conformation, which, in a philosophical view, perhaps justify the arrangement of these naturalists, to whom science is much indebted. While therefore, we have availed ourselves of their method of placing the different genera, our veneration for the ideas and the language univerfally established among men, has induced us to rank all the inhabitants of the fame element in one class, and to distinguish them by a common name.

The

The first peculiarity that strikes us, with regard to the lishes of the spinous order, is the greatness of their numbers. Not only are the individuals of each family more numerous; but the variety of the kinds is also far greater. Upwards of four hundred different species of spinous fishes are already known and described; while the cetaceous and cartilaginous orders, when taken together, hardly amount to a fifth of that number *. The former, are in general inferior in fize; and it is conformable to a law, which obtains in every department of the animal kingdom, that the fmaller the productions of nature are, the more numerous and diverlified in form does the yield them. A very valuable purpose in the economy of providence, is gained by this conflitution of the animal kingdom; for, fince the finaller tribes are in general deflined to become the prey of the larger, an adequate provision is made for the supply of every kind; none entirely perishes through want; none is ultimately extirpated by depredation.

It is by the numbers, therefore, of the spinous sishes that the other orders are preserved, and their own perpetuated. In them, as we have already observed, generation is performed, not by producing a living animal, or by hatching a distinct egg, but by spawning innumerable ova, that are quickened into life by the heat of the sun, and are desined to supply the annual waste of millions. Hence the powers of secundity in this order, exceed belief, and in a short space defy calculation. A single herring, if suffered to multiply unmolested, and undiminished for twenty years, would shew a progeny greater in bulk, than the globe itself. It is owing to this exubevant

^{*} Linnaus has four hunded spinous fishes.

f Goldfinith fays above ten times greater, Nat. Hift, vol. vi. page 335.

rant fertility, that the herring, the pilchard, and fome others, are obliged to migrate annually, from the artic regions, in shoals of such vast extent, that for miles they are seen to darken the surface of the water.

But the amazing propagation of fishes, which we witness along our coasts and rivers, bears no proportion to the vast quantities that swarm in the warmer latitudes of the Indian ocean. The inhabitants of some of the islands there, are, it is said, under no necessity of providing instruments for sishing: As they approach the shore, they are found in great numbers, in the plashes, where the water remains after the ebbing of the tide. In some places where these swarps are dried up by the sun, the sishes are lest in such shoals, that they communicate by their putrefaction, a noxious and unhealthy tincture to the atmosphere.

Happily, however, for the purity of that element, and the health of those beings which it supports in life, the waste of these fishes is nearly proportioned to their secundity; and the balance of nature is exactly preserved. The shark, the porpesse, and the cod, we ought therefore to consider not so much in the light of plunderers and rivals, as that of benefactors to mankind: Without their exertions, the sea would soon be overcharged with the burthen of its own inhabitants; and that element, which at present distributes health and plenty to the shore, would in a short time load it with putrefaction *.

The generation and growth of fishes, and particularly of the oviparous, are involved in great obscurity. The common opinion, as we have already stated, is, that impregnation

dem ubi supra.

which

pregnation is performed without the body of the female, by the male ejecting the smilt, and mingling it with her ova. Something, however, like pairing and copulation, has been observed among the sish kept in ponds. At a a certain period, the sexes are seen struggling together among the grass, at the brink of the water. It is then that the scales of some grow rough, and lose their lustre; that others grow thin, lose their appetite, and become slabby. Should their copulation be established by farther observation, their manner of generation would be more analogous to that of the rest of the animal kingdom:

It is never yet afcertained, whether all the fishes of this order, when they first attain to animation, and burst from the egg, leave it perfect animals, or in a tadpole state, as is the case with the frogs, and many of the lizards. The young frog is first ushered into life, with an enormous head, and slender tail; but the tail foon after drops off, the head diminishes, the legs appear, and the tadpole is metamorphofed into the quadruped. when it changes its element, as well as its form. A species of the lizard also, which is excluded from the shell without legs, acquires them by degrees, and after some time leaves the serpentine shape. Some fishes, it is probable, in like manner fuffer a change, though too imperceptible to attract the notice of the observer. In support of this idea, it is well known, that during the month of July, there appear in the Thames innumerable shoals of small fish, called white bait, that are universally allowed to be the young of some species abounding there. They have no roe; a circumstance which proves them to be young: they refemble no other fish exacily; from

U 2

which the conclusion is, that they undergo a change before they arrive at maturity.

The general character by which naturalists distinguish the spinous sishes from every other, is that bony operculum, which, in this order, universally covers the gills or each side. By these coverings, the gills are alternately opened and shut; and the spinous sishes breathe by these organs alone, without any assistance from lungs. Hence, as these animals partake less of the conformation of quadrupeds, than the two orders which we have already examined, so they can in general remain a shorter time out of their proper element: When taken from the water, they testify their suffering, by panting more violently, and at closer intervals; the thin air surnishes not their gills with proper play, and in a sew minutes they expire.

But the spinous tribes are not all equally incapable or supporting life in the open air; fome are tenacious of it, even in that element. The eel will live feveral hours out of water; and the carp has been known to be fattened in a damp cellar. The manner in which this procefs is conducted, is by putting the animal in a net wrapt up in wet mofs, the mouth only difengaged, for the convenience of feeding: The nourishment with which it best agrees, is white bread and milk; and upon this food it will fatten more rapidly, and become better flavoured than when fed in the pond. It is necessary, however, that the net be dipped frequently in water, and kept hanging in a damp vault. From this experiment, it would appear, that the want of moisture to the gills, is the cause of death in these animals; and could that be Supplied, the vital functions might be carried on nearly as well in the one element, as in the other.




Plate XII.





EELS



DIVISION I. Of Apodal Fiftess

SECTION II.

GENUS XV. The Eel.

In forming this genus, nature feems to have made a near approach to the reptile tribes; like these animals, the body is long, slender, and slexible. Excepting the small pair of pectoral, it may be said to have no sins; for the dorsal, anal and tail sins are united in one web, which surrounds a large portion of the body. The apertures to the gills are small, and placed behind the pectoral sins: they are covered by ten branchiostegous rays.

The eel differs from almost every other fish of that order, of which it is placed at the top, in the manner of its generation. It is viviparous, and is impregnated in the same manner which obtains among the cartilaginous tribes. The ancients entertained very extravagant notions concerning the generation of these animals. Aristotle afferted, that they were neither male nor female, had no ova nor semen*. Hence it was believed that they sprung from the mud, or that the incrustations scraped from

from their bodies by the fiones, received animatica. Rondeletius rashly adopted this opinion, from having obferved that they were generated in pools, from which all the mud and water had been for a while extracted. This was a phenomenon, for which they could in no other way account, than by the spontaneous generation of eels: But later observations have ascertained, that ponds are often supplied with these fish, in the same manner that vegetation is spread, by transporting the seeds of plants. The heron, or other water sowl, may drop the eel, when carrying it to its young, or the young may be ejected undurt from its bowels, as the seeds of plants are voided by land birds, without being injured by the operation of the stomach.

In their habitation, the fishes of this genus are still more singular, than in their manner of propagating their young. They can reside either in selt or fresh water †; and what is still more surprising, they are in some measure independent of either; for they sometimes leave their native element, and wander, during night, along the dewy meadows, not only for a change of habitation, but in quest of prey ‡. Of these nightly excursions, the snail is commonly the victim, being devoured by the eel as it passes along.

There is no animal more vivacious than the eel; when drawn from the water, it will furvive blows that would have killed an animal ten times its fize; and even after it is cut afunder, the different parts are feen to move §. It is, however, so casily destroyed by cold, that to avoid it, it beds itself deep among the mud, and continues, like

the

^{*} De piscibus.

† Brit, Zool, class iv. gen. 12,

[†] Rondelet. de Piscibus. 199.

[§] Idem ibidem.

THE EEL. 159

the ferpent tribe, in a torpid flate during winter. Some have been known to take shelter under a rick of hay in severe weather, and even there, have all perished from excess of cold *. Though fond of hiding themselves in the mud, they are incapable of living in thick turbid water; and hence, when a river is disturbed by a slood, they are frequently suffocated by the impurity of the stream †.

The Common Eel t.

THE external figure of this fish is too well known to require a minute description. The body is long, round, and tapering, compressed towards the tail, and above of a blue colour, tending to yellow on the belly. The river eel has no scales, the body being all over lubricated with a viscous substance, ferving at once to protect the animal from cold, and external injury. Those which are fed in pure running water, are brighter in their colour, as well as more delicate in their sless.

The irides are white, and near the eyes, are feen the orifices of fmell. The branchiæ are four in number, covered with a ikin, and extremely fmall: the aperture externally communicating with them, is so minute, that it occasions their sufficient in muddy water, and is probably the cause why they are capable of living so long in the open air ||.

There

^{*} Albertus apud Gesner, page 45. † Willough. page 110.

[†] Muræna Anguilla, Lin Syst. Anguilla Plinii, lib. ix.

[§] Idem ibidem. | Rondelet. de pifcib. lib. 1. cap. 9,

There are feveral varieties of the common cel, probably occasioned by the different food on which they are supported. These animals are all extremely voracious; they devour carrion or any putrid substance that falls in the way*: They are capable of subsisting equally in fresh water, or in falt, and thrive either in a stream or in a pond, and even in wells; but though a fish almost universal, naturalists seem agreed, that there are none in the Danube, nor in any of those streams which slow into it; while they are found in all the branches of the Rbine;

This species often grows to a pretty large fize, some weighing seventeen pounds; there is indeed an inferior kind in the *Thames*, and about *Oxford*, which neither attain to the same fize nor fatness; they are known by the largeness of the head, and the roundness of the snout, and have there received the appellation of grigs. Every species of the eel was deemed among the *Romans* contemptible food, according to *Juvenal*, from their foul feeding, and their resemblance to a snake.

The Conger Eel §.

This often grows to an enormous fize; fome are taken eighteen inches in circumference, and ten feet long, weighing

^{*} Brit. Zool. † Rond. ubi supra. ‡ Albertus. § Muraena Conger, Lin. Syst. Le Congre, Belon.

weighing upwards of an hundred pounds. A fishery of congers established at Mount's Bay, in Cornwall, forms a very considerable article of commerce. They are annually exported to Spain and Portugal in a dried state, where they are grinded down into a kind of powder, and are used in enriching their soups.*

They are caught by a fort of line called a bulter, baited with pilchards; when taken, they are flit up, that a part of the fat may exude from them before they are falted, and fit for use; and so considerable is the quantity of juice that thus escapes, that a fish of a hundred weight will not dry to above twenty-five pounds. M. Pennant supposes that a fishery of congers might be established with advantage in the Hebrides, could the aversion of the natives to this tribe be overcome.

This species is distinguished by the same voracity as the former; it devours other fish, crabs, and even carcafes. The mode of its generation is probably the same with the common sels; but, however this be, it is certainly prolific, for the number of its young that annually ascends the Severn is prodigious; they are there called elvers, and during the month of April they swarm in such shoals, that they are thrown out upon the shore with small sieves made of hair, and fixed to the end of a pole; a man will in this manner take out as many at one tide, as will fill a bushel †.

The conger differs from the common eel, in having the eyes larger in proportion; the irides of a bright filvery colour, the lateral line marked with a row of small spots; the edges of the dorsal and anal fin black, and in having a greater number of bones.

^{*} Brit. Zool,

[†] Vide Gesner, apud Will.

The Sea Serpent *.

THIS hideous animal may be referred to the genus of eels, which in its external figure, it nearly refembles. It is generally about five feet in length, slender, and almost entirely of the same thickness, till near the tail, where it tapers off in a fmall point. The upper part of the body is of a dirty yellow, the under, bright blue. The faout is long, flender, and prominent, divided by a frightful mouth; armed on the inner parts with small teeth, eight or nine of a larger fize being placed before, and on the middle of the palate. The eyes are small, of ar owing shape, placed obliquely in the direction of the body, the irides of a gold colour, interspersed with brown spots. This species has only two pair of fins, inferred near the gills, and strengthened by fixteen cartilaginous rays. There are two spotted lines, arising from a common point on the back part of the head, and ftretching in a parallel direction along the fides, till they terminate about two inches from the extremity of the tail. The tail is not compressed as in the eels, but round, and fimbriated, with no fins; neither the anal nor dorfal reaching to its extremity +.

There are feveral other kinds of fea-ferpents, fome spotted, and others red; but as these are mostly foreign sishes,

Serpens Marinus. Rondel. Sea ferpent, Will. 107. T Wi llough, page 108.

files, whose history is altogether unknown, we forbear to enumerate them, and to tire the reader wirh the barren and uninstructive discription of their forms *. One species, however, of the eel tribe deserves our notice, because it has been found to possess the same narcotic powers as the torpedo; it is the gymnotus electricus of Linnaus, called by the English of Jamaica, the torporific eel: This fingular fish is only found in the interior parts of South America, particularly in the lakes of Surinam, from whence it has been transported to Jamaica. Different attempts have been made to introduce it into Europe, but without fuccess. If touched with the hand, it communicates a firong flock to the arm; and the fame effect is produced by applying it to a metallic rod, whereas it is harmless when touched with a piece of wood. From these experiments it appears certain, that the matter discharged by the torporisic eel, partakes of the nature of the electric fluid. The tribe of gymnoti in which Linnaus places this animal, comprehends five different species; of the eels, properly so called, he enumerates only feven kinds +.

Vide Syft: Nat;

Vide Rondel, et Aldrovand.

SECTION III.

GENUS XVI -The Sea Wolf *.

This voracious fish inhabits the northern parts of the ocean; it is found on the coasts of Ireland, Greenland, and Norway, but seldom ventures farther south than those parts of the German Ocean which wash the shores of Britain; and Hollond. The back, fins, and sides, are of an azure thue; towards the belly it is white, and the whole skin is smooth §.

What particularly diffinguishes this animal, is its large mouth, and formidable teeth; the fore teeth are ten or twelve in humber upon each jaw, round, conic and sharp; behind these are the grinders, round and slat, about twenty-four in number; upon the bones of the palate are three rows of similar teeth. There is no sish seems more completely armed for devastation than the sea-wolf, and mone more willing to use the instruments with which nature has supplied it; it will gnaw even the anchor of a ship, so that the noise is heard above; the marks of its teeth are plainly discernible on weighing it up ||.

The

^{*} Anarhicas Lupus. Lin. Syst. Cat-fish, Sibbald.

⁺ It is there called Steinbeisser, \$ School. 45.

S Brit. Zool. Class iv. gen. 13. Will. p. 130.

The teeth of this fish are frequently dug up in a petrified state, where they are called busonites, or toadstones; formerly they were much esteemed for their imaginary virtues; they were sometimes set in gold, and worn as jewels *; the teeth of three large sishes of this species, were seen at Scarborough, every one of which were either broken or dissigured by the hard substances which these animals had attempted to macerate †.

Both the dorfal and anal fins of this fish extend almost to the tail, which is round at the end, and strengthened by fourteen rays; the pectoral fins resemble two small wings, being about five inches long, by seven broad. The food of the wolf fish is crustaceous sishes, the shells of which it can easily comminute. These animals are oviparous; and their young, for some time after their production from the egg, are of a greenish cast, resembling the sea weeds among which they then reside; some of them have been caught in the Frith of Forth.

^{*} British Zool.

[†] Willough, p. 130;

GENUS XVII .- The Sand-Eel, or Lance *.

THESE fishes somewhat resemble the eel in their external form, as their name imports. They are dug or hooked up from the fand in the shallow pools that are left by the reslux of the tide, commonly for bait, though they are themselves reckoned delicate food. The sand-eel has frequently been found in considerable quantity in the belly of the porpesse, which confirms the account we gave of that animal digging up the sand in searching for its prey. Linnaus has only one species belonging to this genus, that which we frequently see dug up along our shores.

The body is nearly a foct long, of a fquare form, but rounded towards the fides, which are divided, each by a ftrait line proceeding from the head, and terminating in the tail; the lower jaw projects beyond the upper, and when extended, the gape of the animal is very wide; there are no teeth, the long fharp tongue moving in a mouth entirely fmooth; the pectoral fins are placed near the gills; the dorfal and the anal fins are fupported by numerous rays, and proceed nearly as far back as the tail ‡.

^{*} Ammodytes Tobianus. Lin. Syft. Ammodytes Gesneti

[†] Willough. p. 114.

GENUS XVIII .- The Ophidium *

THE fishes of this genus are two; the one has a number of barbs hanging over from the lower jaw, while the other wants these, but in all other respects appears the fame. The ophidium has a great refemblance, in shape, to some of the tribe of eels, but is of inferior fize. Belonius afferts that those in the Mediterranean are not above a handbreadth. The back is cinereous; the fides of a filver colour: There are no fcales, but in their place a number of oblong fpots here and there difperfed over the body; the mouth is large, and exasperated with a number of small teeth round the palate and jaws; the eyes are large, covered with a pellucid membrane: There'is only one pair of fins near the branchiæ; the dorfal fin takes its rife not far behind the head, from whence it proceeds along the back, furrounds the tail, and terminates below at the anus. The body is exactly divided by a lateral line, which extends along each fide from the head to the tail +.

Ophidion Plinii. Ophidium, Guan. † Vide Guan. Hist. Piscium, p. 117.

GEN. XIX .- The Stromateus *.

LINNAUS enumerates two species under this gerus, the hatola and the parda to The characters are, a round flat body, covered with a smooth glutinous skin, and destitute of scales i. The fize of the fiatola or lampuga, as it is called by the fishermen of Rome, is nearly a foot, and its weight a pound and an half; the colour of the upper part of the body is pale azure, that of the belly filver; the whole beautifully ornamented with yellow fpots; upon each fide are two lines, the one straight, and the other incurvated like a bow &; the fnout is flat, the mouth finall, and the tongue fmooth, fleshy, broad, and moveable; the eyes are fmall, and covered with the common membrane that furrounds the head; the pectoral fins are furnished with a great number of rays, and the ventral are wholly wanting; the dorfal fin rifes about a third part of the total length from the head, is supported by forty-fix cartilaginous rays, and is prolonged till within less than an inch of the tail; the anal fin nearly refembles it in shape, and takes its rife still nearer to the tail.


Willoughby observed these since they are reckoned most delicate food, and bring a high price; none of them, have yet been found in the British seas.

[·] Stromateus, Rondel. Callichthys, Belon.

[†] Syft. Nat. 169. † F. Guan, Hift. Pifcium. p. 173. Will. p. 156.



Plate XIII.



GENUS XX .- The Sword-Fish *.

Or this genus there is only one species, which on that account is allowed to retain the generic name of sword-fish, an appellation given this animal almost in every country, and evidently derived from the peculiar conformation of the upper jaw; it projects about four times the length of the lower one; is compressed at the top and bottom, and sharpened towards the point. This enormous snout is three feet long, resembling a sword in shape; its substance is rough and hard, but by no means capable of piercing and sinking vessels in the sea, as is afferted by Pliny;.

The fword fish grows to a large fize, the head alone, being in some instances known to weigh upwards of seventy pounds §. The body is long and slender, thick towards the head, but tapering off into a small fize as it approaches the tail: The colour above is black, and on the belly of a filvery white: The mouth is without teeth; the lower jaw terminates like the upper one, in a sharp spear-like point, but is greatly inferior in length: The dorfal sin takes its rise above the gills, and continues till it nearly reaches the tail; it is supported by twenty-fix rays; the sirst of which is by far the strongest and highest, the Vol. III.

^{*} Xiphias Gladius, Lin. Syst. L'Heron de Mer. Belon.

[†] Will. p. 161.

[†] Rostro mucronata esse, ab hoz naves persossas mergi in oceano. Lib. 32

[§] Brit. Zool. Class iv. gen. 16.

other gradually diminishing, till the four last, which again rise higher, and give the appearance of a second sin. The tail branches into forks, exactly resembling a crescent: There are on each side, a little above the tail, two triangular protuberances, formed by the skin, resembling the spurious sins of the tunny: the pectoral sins are placed at the gills; the sirst ray of these is also by far the longest, which gives them the appearance of a scythe. The anus is placed about one third part of the body from the tail, and below it are two anal sins connected by a common membrane. The sword-sish is exceedingly voracious, and is a great enemy of the tunny, which discovers its fear as soon as it approaches. Ovid and Belon take notice of its hostilities against this timid prey *.

The fword-fish fometimes frequents the British feas, but is much more common in the Mediterranean; the Straits of Messina are particularly famous for it; and it was probably upon a promontory there, that the speculatores, or perfons employed to watch and give notice of its approach, were stationed. Willoughby informs us that he went himself to Scylla to be a spectator of the Italian method of killing this animal: The spies above, on seeing the sword, make signals to the boats below, directing the failers where to steer: As soon as the vessel reaches the spot where the silhes are, one of the most skilful of the fishermen gets upon a mast, erected for the purpose, and directs the boat till it comes within reach of the particular sish at which he intends to aim; he then comes

^{*} Ac durus Xiphius, ictu non mitior enfis; Et pavidi magno fugientes agmine Thunni. Halieut. 97.

down and pierces it with a fpear; this instrument being fastened to a rope, the animal is allowed to struggle till it is overcome with satigue and pain, when it is either taken up into the vessel, or drawn ashore, according to the size of the fish *.

The flesh of this animal is whiter than that of the tunny, nourishing and not unpleasant in its flavour. The inhabitants of Sicily reckon it one of the first delicacies, equal to the sturgeon, and purchase it frequently at the price of sixpence the English pound †.

There is another fish nearly allied to the above, which is peculiar to the American seas, and called by the Brazilians guebucu. The snout is sharp, bony and hard; its length is sixteen inches, the lower jaw reaching above one half of that space. The length of this species is sour feet, and its thickness one; it is distinguished by two remarkable long bony substances, resembling a rod, in the place of its ventral sins, which it has the power of solding down into a kind of surrow made in the belly for their reception; the dorsal sin is three feet long, and capable of being hid in the same manner.

^{*} Vide Strabo. Lib. i. apud Will.

[†] Willough. page 163. ‡ Idem ubi supra,

Division II .- Jugular Fishes.

GENUS XXI .- The Dragonet. *

THERE are three kinds of the dragonet enumerated in the fyftem of nature, the lyra or gemmeous, the dracunculus or fordid, and the *Indian*: They are all defittute of fcales; and are of a smooth, uniform body, attenuated towards the tail. The mouth is very small, thickly set with aceth, upon the jaws, tongue and palate †. The eyes are prominent, somewhat large, placed vertically, and near to each other; hence the name of ouranoscopi, which these animals have obtained from the old naturalists ‡. The upper lip is in this genus double; and it is also distinguished by a three forked spine, rising from each of the epercula of the gills, and lying backwards along the head.


The gemmeous dragonet, is found as far north as Norway and Spitzbergen, and fouth as far as the Mediterranean: it is not unfrequently upon the Scarborough coasts, where it is taken by the hook in thirty or forty fathoms water. It grows to the length of ten or twelve inches, and is


^{*} Callionymus Lyre, Lin. Eyft. Yellow Gurnard, Phil. Tranf. No. 29;

[†] Guan, Hift. Pife. 121.

[‡] Vide Gronov. Zooph. No. 206.

Plate XIV.







COMMON WEEVER



eften found in the stomach of the cod-sish. It breathes in the manner of the cetaceous sishes, by means of two orifices in the crown of the head, through which it forces the water received by the mouth *.

The ventral fins are jugular, and placed before the pectoral, a circumstance by which all the fish of this division are distinguished; they are placed at a distance from each other, are broad, and of one piece. The pectoral fins are round, of a light brown colour, the rays aculeated, and extending beyond the membrane that connects them like launces. The dorfal fins are two; the first of a form so singular, that it at once distinguishes this species from every other. The first ray is cetaceous, and extends in a curvated direction, almost as far as the tail. The rays of the second dorfal fin are of an equal length, except the last, which extends a considerable way beyond the rest: The anal sin is placed exactly opposite to it, and resembles it very nearly, both in shape and size.

The tail of this species is long, rounded, and supported by ten rays. It has obtained the name of the gemmeous dragonet, from the variety and brilliancy of its colours: When taken out of the water, the yellow, the white, and rich cærulean blue, with which it sparkles, make a fine appearance; the last colour in particular glows with inexpressible splendour.

The fordid dragonet is inferior in fize, as well as beauty; being only about fix inches in length; and about two in thickness. It is of an olive colour above, and white below, The fides are marbled with small spots of filvery

Brit. Zool, clas iv. gen, 17.

filvery blue; and upon the top of the head is a triangular fpot of a purplish colour. It frequents the British feas but rarely; Willoughly, however, afferts that it is frequently feen in the fish markets of Rome and Genoa: Its flesh refembles that of the bull-head, and its manner of feeding is the same.

GENUS XXII. The Uranoscopus *.

Of this genus there is but one species, a fish about nine inches in length, with a large, flat, and circular head. The lower jaw projects beyond the upper, and is turned upwards at the end; the eyes are fmall, prominent, and almost contiguous. The face is flat, and has the appearance of confantly staring upwards. Both the jaws are armed with sharp teeth, as is also the palate and whole infide of the mouth. The whole face, but particularly the covers of the gills, are rough and unequal with tubercles, the bases of which are surrounded with spines; and at the extremity of each of the opercula, where the head unites with the body, there are two pointing backwards, remarkably sharp and large. The uranoscopus has two dorfal fins; that nearest the head is small, of a black colour, and supported by three sharp rays: The fecond is higher and broader, and contains fourteen rays: The ventral fins in this species are placed far before the pectoral, and feem directly under the eyes: It is frequently caught in the Mediterranean, and is faid to be a delicious morfel +.

^{*} Trachinus, Artedi Uranoscopus. Aldrov.

[†] Gesner de piscih.

GENUS XXIII. The Weever *.

THERE are several species of this genus found on the British coasts, although Linnaus only mentions one, the draco, a fish of along and compressed figure, marked upon the fides by a number of cinereous lines, that arise from the middle of the back, and proceed along the fides towards the belly. The two lateral lines that extend from the gills to the tail, are placed much nearer the back than those of any other fish. The branchice are yellow, the belly white, and the whole body is covered with fmall thin scales. The eyes are placed remarkably near to the fnout; the irides are yellow, punctuated with black fpots. The lower jaw projects beyond the upper; both armed with fmall teeth. The apertures of the gills are uncommonly large, and each of the covers is furnished upon the upper angle, with a strong sharp spine. The dorfal fins are two; the first confisting of five sharp prickly rays; the fecond is supported only by cartilaginous rays, and is prolonged almost to the tail. The pectoral fins are fituated lower than ufual; the ventral fins are jugular, each strengthened by fix rays. The anus is not far from the throat; and from it to the tail, there proceeds a long fin fortified by thirty-two rays, which project beyond the membrane that connects them +.

The

^{*} Trachinus Draco, Lin. Syst. Draco Marinus, Plinii.

[†] Fide Willough. page 288.

The weever is frequent in the Mediterranean, where it grows from ten to fifteen feet in length; it is capable of inflicting a very fevere wound by the spines of the first dorfal fin, which, if neglected, becomes cancerous, according to Rondeletius *. For feveral hours, the part affected is attended with a violent burning, and shooting pain, fometimes with an inflamation, that reaches from the arm to the shoulder. These essects are commonly supposed to proceed from something venemous, lodged in the spines, which look indeed suspicious; for those of the first dorsal sin are dyed with a blackish substance. Various remedies are applied to relieve the pain of this wound. Rondeletius recommends the flesh or brains of the animal: our fishermen rub the parts affected with sea fand; while those of Scarborough successfully use fiale urine, applied warm.

This animal is excellent food, and therefore is often taken, notwithstanding the noxious quality of the spines: it buries itself in the sand, leaving only its nose out; and if trodden on, immediately strikes with great force, each blow directed with a degree of judgment that evinces a consciousness in the animal of its own noxious powers f-Willoughby describes three species belonging to this genus; the first is the sish above described; the second a larger kind, seen at Rome, with beautiful variegated colours; and the third a Brazilian sish, called niqui by the natives.

^{*} De piscibus, 300. † Brit. Zool. class iv. genus 18.

GENUS XXII. The Cod .

This genus contains a very numerous, and well known tribe of fishes, a considerable part of which frequent the British coasts, and contribute more largely to the sub-sistence of the inhabitants of Europe, than any of those samilies which we have hitherto reviewed. Their general characters are, a smooth head; seven slender branchiostegous rays; an oblong body, covered with small deciduous scales. The number and situation of the sins are various; the teeth are small, and numerous, placed upon each jaw, and in the upper part of the mouth †.


The flated migrations of most of the fish that compose this genus, is one of the most remarkable circumstances in their history: in these annual voyages, in the immensity of their numbers, and in their social habits, they bear a strong analogy to birds of passage. The cod, the haddock, and the whiting, issue forth in immense shoals from the artic seas, very early in the spring, and after having dispersed over the temperate latitudes, again regularly return to their northern retreats about the same time of the year. The necessity of procuring food has been assigned as the cause of their annual migrations from the artic seas; and their retreat thither has been ascribed to the security that these unfrequented tracts are supposed to afford them, while they deposite their spawn.

But

^{*} Afellus Major Vulgaris, Willough. Gadus Morhua, Lin. Syft.

[†] British Zool. class iv. gen. 19.

Plate XV.







WHITING POUT



WHITING





But although the cod undertakes annual excursions of confiderable length, it still may be regarded as a local fish; for it never ventures into the warmer tracks of the ocean. None are found in the Mediterranean *; and few in those parts of the Atlantic of the same latitude. They are in greatest perfection, and seem to prefer that space lying between the fiftieth and fixtieth degrees; fuch as are caught beyond it, being always inferior, both in quantity and quality +. Their grand refort for centuries past, has been on the banks of Newfoundland, and other fand banks off Cape Breton. That extensive flat seems to be the broad top of a subaqueous mountain, every where surrounded with a deeper fea. Hither the cod annually repair, in numbers beyond the power of calculation, to feed upon the worms that fwarm upon the fandy bottom. Here they are taken in fuch quantities, that they fupply all Europe with a confiderable quantity of provision. The English have stages erected all along the shore, for falting and drying them; and the fishermen, who take them with the hook and line, draw them as fast as they can throw them out I.

This immense capture makes no sensible diminution of their numbers; for after their food is consumed in these parts, or when the season of propagation approaches, they take their departure for the polar seas, where they deposit their roes in full security, and repair the waste which has been occasioned by death, or the depredations of their enemies. They annually make their appearance on the coasts of section, Norway, and Britain, gradually diminishing in their numbers, as they proceed to the Z 2

Rondeletius de piscibus. † British Zool, ubi supra,

de Goldfmith's Nat. Hift, vol. vi. page 325.

fouth, and ceasing altogether before they advance to the straits of Gibraltar *.

Before the discovery of Newfoundland, the greatest fisheries of the cod were on the coasts of Iceland, and the western isles of Scotland, where the English resorted in quest of them, as early as the beginning of the sisteenth century. Our right of sishing in these parts, however, was not acknowledged by the government of Denmark †, till the reign of James I. whose marriage with a princess of that country, secured to his subjects that indulgence of which they availed themselves so completely, that they had then a hundred and sifty ships employed in the Iceland sishery.

Even on the banks of Newfoundland, the French, Spaniards, and Portuguese, had originally a far larger portion of the sithing, than the British: In 1570, the former nations had upwards of three hundred vessels employed in that trade, when those of the English did not exceed sity t. Matters, however, have since been reversed; and the English shipping on that coast has intendedly increased; it is now superior to that of any other nation, and the trade is deemed a valuable accession to the wealth of individuals, as well as to the naval power of the empire §.

This immente fithery is conducted in a tract of the fea, apitated by a perpetual fwell, and involved in continual darkness, by means of a thick fog, that constantly hangs

over

^{*} British Zool. † Rymer's sæd. xvi. 275.

Hackluyt's coll. voy. iii. 132.

^{§ 15000} British framen are at prefent employed in this sishery. British Toology.

over it: The bait used is herring, a small sish called capelin, a shell sish, and bits of sea sowl. The natural sood of the cod is small sish, testaceous animals, such as crabs and whelks; and their digestive powers are so strong, that they dissolve every substance which a voracity almost indiscriminate, leads them to devour. Their sight is probably very impersect; for almost every small body that is agitated by the water, attracts their rapacity, stones and pebbles not excepted, for these are often sound in their stomachs.

The founds of the cod fish, are reckoned a great delicacy, and frequently brought from Newfoundland, salted up by themselves: They are employed by the fishermen of Iceland, in making isinglass; are obtained, by carefully separating them from the back-bone, to which they adhere after the fish is cut up *.

The general weight of the cod fish on the British coasts is from fourteen to forty pounds; some have indeed been caught near eighty, but those of the middle fize are most esteemed for the table. Their time of spawning is from January to April, when they deposit their eggs in rough rocky ground. After having been exonerated of a load containing frequently three millions of young, the parent recovers its plumpness sooner than almost any other fish; and is caught in good condition, during almost the whole summer.

Schonfeldt remarks a kind appointment of providence in the immense fecundity of this fish, and in that abundant supply which it affords to the inhabitants of those bleak

^{*} Phil. Trans. 1773. † Vide Schon, apud Willough. pape 166.

bleak and frozen countries, that are unfit for the production of grain. The ichthyophagi of these barren regions, says he, not only furnish themselves with a substitute for bread, by drying this sish, but send a vast quantity of their surplus stores to add to the supply of other nations. The numbers and sertility of these sish, seem indeed amply to justify the grateful exultations of this writer; for they are such as will forever basse all the efforts of man, and the voracity of the inhabitants of the ocean, to exterminate their race; and will secure to every age an inexhaussible supply of wholesome provision.

The cod is a fish fe well known, that it requires not any particular description; the colour, which is commonly finereous, and sported with yellow on the back and sides, varies much according to the age and residence of the animal.

The Hadock +.

This species appears upon the coasts of Britain annually, particularly those of Yorkshire, about the middle of December, and in such vast shoals, that they cover a tract frequently of many miles. They do not venture far from the shore; when the sishermen cast their lines beyond their limits, which is commonly about three miles

from

^{*} Vide Schonf. apud Willough. page 166.

[?] Gadus Ælglefinus. Lin. Syft. Tertia afellorum fpecies,

from land, they are feldom any taken. The larger hadocks leave the coast as soon as they begin to be out of season, seaving behind only the smaller fish. This remark is applicable to all the fish that appear on the Yorkshire coast, except the mackrel, which alone remains after becoming unfit for use *.

The hadock is the most common species in the London market, as it is not only plenty, but in season during the greatest part of the year. The larger sish begin to roe in the middle of November, and continue so till the end of January, when they are unsit for use. Those of the middle size recover about the beginning of May, and are in season till February; such as are incapable of breeding remain at all times sit for the table. In stormy weather none of this species take the bait; the sishermen affert, that they then ooze in the bottom of the sea, and shelter themselves there, till the agitation of the water has ceased: In proof of this, they allege that those which are taken immediately after a storm, are covered with mud upon the back.

The common fize of the haddock is twelve inches in length, of a dark colour upon the back, and the skin covered with small scales. On each side about the middle is a large black spot, the prints, as it is said, of the singer and thumb of St. Peter, when he held this species, and took the tribute out of its mouth: A mark which superstition has extended to the whole race of hadocks, at once to attest and commemorate that miracle; unfortunately, however, the haddock is not the only sish that has been supposed to be thus distinguished by the marks of the apostolic touch †.

Of the same genus, but inserior in size, are the pouts, blinds, and poors. The latter is the smallest of the cod kind yet discovered; and it is the only one sound in the Mediterranean. It is taken near Marseilles, and in such quantities as sometimes to prove a nuisance. It seldom exceeds six inches in length, and is unsit either for being dried or salted; if the sirst is attempted, it grows as hard as an horn*. The whiting is the most delicate and wholesome food of any of this genus: It appears in vast shoals on our coasts in spring, and is caught with the line, from one to three miles from the land.

The Coal-Fish +.

All the species of the cod, which we have mentioned, are distinguished by a barb under the lower jaw; in the coal fish this is wanting. It is of a dark cerulean blue, and has from that circumstance obtained its English name. The eyes are large and protuberant, and the upper jaw, which in the common cod projects beyond the lower, in this species is the shortest of the two ‡.

The coal fish frequent the rocky and deep coasts all round this island, but are found in greatest plenty in the vicinity of the *Orkneys*, where their fry constitute a great

Belon de piscibus, p. 120.

[†] Colfisch Anglorum, Gesn. Gadus Carbonarius, Lin. Syst.

^{\$} Will. p. 168.

ges;

great part of the support of the poor. The young begin to appear on the coast of Yorkshire, in the month of July, in numbers that defy all computation: They are at that period only an inch and a half long; in August they are from three to five inches, and are taken in vast quantities with the angling rod; they are then reckoned a very delicate dish, but afterwards grow so coarse, that, by the time they are a year old, sew people eat them*. They are sold either fresh, or when salted, in the counties of York and Northumberland, at an inferior price to the outer species of cod.

The Hake to

This species has only two dorsal fins; grows from two to three feet in length, and is of a more slender shape than the common cod. Its form somewhat resembles that of the pike; whence it is called the sea pike by the French and Italians. The mouth is large, and surnished with a mixture of short and long teeth; and the palate, as is common to the fishes of this genus, is beset with sharp spines or teeth. The first dorsal fin has nine rays; the second, which extends almost to the tail, has forty; the pectoral fins are surnished each with twelve rays; and the ventral, which are situated before, are supported by seven cartila-

VOL. III.

^{*} Vide Schon. apud Will. p. 169.

[†] Afellus primus. Will. Gadus Merlucius, Lin. Syft.

Merlucius, Vide Gesner and Will,

ges; the tail is not forked, but each ray being equally produced, terminates in the same straight line.

The hake is found in abundance on many of our coasts; particularly those of *Ireland*, where there was formerly a stated fishery on the *Nymph Bank* off *Waterford*; immense quantities were caught there at the two seasons of their periodical appearance, June and September, when six men with hooks and lines frequently killed a thousand fish in one night; the produce of this fishery was salted up and exported to *Bilboa* in *Spain*; it has, however, been for many years upon the decline, owing to the fish deserting their wonted station *.

This dereliction of their accustomed haunts, is not peculiar to the fishes of this species; the haddock has in the same manner abandoned the coasts of Waterford, and the herrings and the basking sharks have displayed the fame caprice, in relinquishing their stations on several parts of the British shores. Naturalists have not yet given any plaufible account of this irregularity in the migration of fishes: In some instances it may be occasioned by the close pursuit of an unusual number of predatory fish, to avoid whose voracity they may be driven upon shores that they were formerly unaccustomed to frequent; a deficiency of the fmaller fith, that supplied them, may, in other instances, have forced them to abandon a residence, where they could no longer be supported . But the pernicious custom of trawling, is perhaps the most common cause of their abandoning the usual stations, because, by that means, not only a great part of their fpawn is demolished, which was lodged in the sand, but the worms and infects

[&]quot; Smith's Hift. of Waterford, p. 261.

infects which conflituted their food, are either destroyed by it, or driven into deeper water *.

There are three varieties of the hake found in our feas, befides that above described; the greater and leffer forked hake, and the three forked species: The last, Linnæus has omitted in the System of Nature; and the two former he has placed among the blennies. The bifurcated hake is about twelve inches in length; the colour a deep brown, excepting the colour of the lips, which are snow white; these surrounding a broad slat snout, give the animal a strange appearance, sufficiently characteristic of the species.

The Ling to

This species is longer than the river pike, being from four to fix feet; the fides and back in some are olive-coloured, in others cinercous; and the whole figure of the body nearly resembles that of the fish last described: the apper jaw projects beyond the lower, both being exasperated with several rows of short teeth; the palate is armed in the same manner, with parallel rows of small teeth, here and there interspersed with a sew of superior size, strength, and sharpaes; the mouth and tongue are large; from the corner of the lower jaw there hangs a single

A a 2 bark

^{*} Brit. Zool. Class iv. gen. 19.

[†] Afellus longus, Will. Gadus molva, Lin. Syft.

barb about two inches long; upon the back are two dorfal fins, the first consisting of sisteen, and the second of fixty-sive rays, extending like the anal fin, opposite nearly to the tail; the pectoral fins are each supported by sisteen cartilaginous rays, and the ventral by six; the three first projecting beyond the connecting membrane, terminate like spines *.

The ling abounds more or less upon the coasts all round the British isles; it is, however, most frequent near Scilly isles, and those on the west of Scotland and Ireland, where it has long formed a considerable branch of commerce, regulations being framed concerning it so early as the reign of Fdward I:I+. The sish is in perfection from the beginning of February till May; in June they deposit their spawn in the soft, muddy bottoms near the mouths of rivers; about this time the males separate from the semales, many of the former being caught by fishermen, without a single individual of the latter.

When the ling is in feafon, its liver is white, and abounds with an oil of an excellent quality and flavour; but as foon as the fish becomes out of feafon, the liver gradually assumes a red colour, resembling that of an ox, and it then produces no oil: The same change of colour of the liver is observeable in cod, and several other sishes, but not in a degree so remarkable. It would seem, that this oil which is lodged in the cellular membranes of sishes, returns into their blood, and supports them during the period of procreation; a season in which they pursue the work of generation with so much cagerness, that they neglect their food,

As

As the oil of this species can only be obtained while the animal is in season, so it can be extracted from the liver only by a flow fire; if a violent heat be applied in melting it, a very small quantity will be had from the fish, even in its greatest perfection. Such fish as are cured for exportation, must measure a certain length from the shoulder to the tail, otherwise they are not entitled to the bounty which parliament has granted for the encouragement of this trade; twenty-six inches is the length of a fizeable fish; those of inferior dimensions are called drizzles, and because incapable of procreation, they continue in season during the whole summer *.

There is a fresh water fish that somewhat resembles the ling, called the burbot, or eel-pout †: it abounds in the lake of Geneva, where it is called lota, and is sound also in several of the rivers in the north of England. The body is smooth, soft, and lubricated like that of the cel; the number and disposition of sins, resembles that of the ling: The season of generation in this species is December; at which period, though the body is small, one hundred and twenty-eight thousand ova have been taken from a single semale, each of which is capable of becoming a fish, after secundation by the male.

^{*} Brit. Zool. ubi fupra.

⁹ Will. p. 125.

The Five-Bearded Cod *.

THE colour of this is a deep olive brown; in its shape and flimy covering it fomewhat refembles the eel; it is, however, much shorter and thicker, especially towards the belly: There are two different species, the one with five beards, and the other uniformly with three; the former has four upon the upper jaw the latter only two. From the extremity of the lower jaw in both, there hangs a beard: The scales of both kinds are extremely minute; each has the same number of fins, and in the same position; they are diffinguished by the shape of the two dorfal fins; the first rifes a little way behind the head, where it is lodged in a deep furrow, and confifts of a number of short unconnected rays, or filaments; behind this furrow, in the middle of the back, rifes the fecond dorfal fin, confiderably higher than the first, and extending almost to the tail, which is rounded at the extremity; this fin is fupported by fifty-fix rays, and the anal one, opposite to it, by forty-seven. This species grows to the length of nineteen inches, and weighs upwards of two pounds.


The three-bearded cod is nearly of the same size with the other, of which Willoughby reckons it only a variety; it is, however, easily distinguishable, by having the upper part of the body variegated by a number of black spots upon a reddish ground. The Cornish sistemen

have

^{*} Mustela vulgaris, Rond. Sea-Loach, Will.



Plate XVII









have particular cant phrases, by which they imagine they charm these sishes, which they repeat in the same manner that the Sicilians do their mamassu di pajanu, &c. when they are in pursuit of the sword-sish *.

The Torsk.

We have already feen the fishes of this genus, possessing, fome three, others two dorsal fins; the fish now under observation is an inhabitant of the Orkney seas; and from having only one dorsal fin, forms a class distinct from both. It is called by the inhabitants of Orkney and Shetward, tusk or brismac; and is seldom found in a lower latitude than that of these islands, where it swarms in great abundance, and is either dried or barrelled up for exportation.

The torsk grows commonly to the length of twenty inches, and to the depth of four; the head is small, the upper jaw projecting beyond the lower, and both armed with a multitude of small teeth. The sish known in Sweden by the same name, is described by Linnaus with three dorsal sins; that of the Orkneys, however, has only one rising about six inches behind the snout, and extending nearly to the tail; opposite to it is the anal sin, commencing at the anus, and reaching the same length; the

rave

rays of each are but indistinctly perceived, on account of the thickness of the membrane that covers them.

From the extremity of the lower jaw there hangs a fingle heard; the colour of the head is dusky, that of the sides and back is yellow, and of the belly white; the pectoral fins are brown; the dorsal, caudal, and anal fins, are dusky, surrounded by a white margin.

SECTION V.

GENUS XXV .- The Blenny.

THE generic characters of this tribe, are, a short, blunt input, a fmooth body, covered with a glutinous fubstance, and compressed laterally. The teeth are slender, and the ventral fins fmall, supported only by two connected rays. The dorfal fin begins a little behind the head, and reaches the whole length of the back. The fithes of this genus are variously classed by naturalists, some of them being mingled with those of the last, while others are classed with the sea scorpions, and other heterogeneous tribes. Linnaus enumerates thirteen different species of the blenny, only four or five of which are known to frequent our coasts; the gattorugine, the crested, the smoothheaded, the fpotted, and viviparous blenny; all thefe haunt the rocky shores, and at low water are found under the stones among the tang. They are extremely active and vivacious; by means of their ventral fins they can creep among the rocks; and fome species can live out of the water during the length of a whole day.

The Gattorugine *.

The species has been sound on the coast of Wales; the body is without scales, and lubricated with a slimy substance. Towards the head it is thick, gradually diminishing towards the tail; the sides all along considerably compressed. It is above of a dusky hue, marked across with undulating lines; below of a pale ash colour, the ventral and pectoral sins partly orange. The rays of all the sins of this species project beyond their webs, and have a spiny appearancee. The eyes are situated almost upon the summit of the head, contiguous, and extremely protuberant; between them arises a kind of crest, which separates into sour branches, and which can be raised or depressed by the animal at pleasure. The cirri forming this crest are thick at the base, sharpening towards the top, and setaceous along the sides †.

Similar to the gattorugine, is the crefted blenny, a fish distinguished by the same crest-like fin upon the top of the head. Its residence is the same; for it is always found on the rocky shores, where it probably seeds on crabs and small shell-fish; the remains of those animals being found in its stomach ‡.


The smooth blenny resembles the crested, almost in every respect, except that of the smoothness of its fore-head, which is destitute of that erect sin, the use of which no naturalist has hitherto pointed out.

^{*} Gattorugine Vinetiis, Will. Blennius Gattorugine, Lin. Syft.
† Vid. Willough. p. 132. ‡ Brit. Zool.



Plate XVIII.





CRESTED BLENNY





VIVIPAROUS BLENNY

The Viviparous Blenny *.

THE viviparous blenny in its mode of generation, differs from the whole order of spinous fishes, in producing living young. In consequence of this, its habits must be confiderably different. Impregnation must take place within the body of the female, who, like most other viviparous animals, probably is diffinguished by parental affection for her brood, which generally confifts of two or three hundred. The feafon of her parturition is a little after the depth of winter; before impregnation by the male, the eggs are small, and of a whitish colour; after that period they are gradually increased in fize, and affume a red appearance. By and by they acquire an oblong shape, and two black spots begin to announce the rudiments of the head and eyes; after this the belly and viscera begin to appear: they are covered with a milky pellucid membrane, through which the intestines are feen: Last of all, the tail becomes discernible, like a small thread, bending towards the extremity.

During the whole time of gestation, the abdomen of the semale is greatly distended, not only by the young brood, but also by a white muddy substance, which probably supplies the nascent family with food. As the period of parturition approaches, this liquid gradually diminishes in quantity, till, upon the young attaining their full size.

Bb2

Mustela marina vivipara, Schon. Blennius viviparus, Lin. 879.

it almost wholly disappears, as if consumed by the numerous race which it has supported *.

When the fœtus have acquired nearly their complete fize, they creep from place to place in the uterus, as if attempting to difengage themselves from a state of consinement, which is now no longer necessary: after parturition, they are immediately capable of swimming and following the mother; and of supporting life by the same movements of the mouth and gills.

The young of the viviparous blenny, at their first appearance, are about two inches long; when full fized, they are nearly a foot. Their flesh is extremely coarse, and unsavory; it is ate chiefly by the poor, who sometimes take them in their pregnant state; and least their numerous young when dressed, should adhere to their knise or singers, they take the semale, before boiling, and strip her of her young, by passing the hand along the belly. Several of these fish are caught in the river E/k, in York. Solver, by throwing lines from the top of the bridge $\frac{1}{2}$.

^{*} Willough. p. 123.

[†] British Zoology, class 4. genus 20.

DIVISION III.—Thoracic Fishes.

SECTION VI.

GENUS XXVI - Cepola *.

We are now come to that division of the spinous tribes termed by naturalists the thoracic, from the position of their ventral sins, which are neither placed before the pectoral, nor behind them, but directly under them upon the thorax +. The first genus of this class, Linnæus terms the cepola, in the latter editions of his System of Nature, from a sish of that name caught in the Mediterranean. This animal has an extremely slender and tapering shape; the body being twelve inches in length, and scarcely one in thickness: It is of a slesh colour, and semi-transparent, so that the vertebræ of the back appear, and can easily be numbered.

The cepola has no feales; the fides are adorned with a lineal row of filvery fpots. The pectoral fins are finall, and their rays fo flender, that they are almost imperceptible. About an inch behind the head, rifes the dorsal fin, which is produced till it joins the tail, where it meets the anal fin, which is about thrice as broad; and begins

fo

^{*} Tonia prima, Rond. Cepola tonia, Lin. Syft.

[†] Vide Systema Nat. 156.

fo near the throat, that the anus is fituated almost summediately below the angle of the lower jaw. This fish is fold in the markets of Rome; but its fiesh, according to Rondeletius, is of a very indifferent quality*.

The only remaining fish of this genus is the red cepola, alled by the Genoese cavagiro †. The body of this species is more slender and compressed than that of the eel: The sides, back, and tail, are of a pale red; the belly white, and the whole body is destitute of scales. The dorsal sin begins immediately behind the head, extending along the back till it meets the anal sin opposite to it, at the extremity of the tail. In the larger sishes, these sins display a great brilliancy of colour; at the base they are of a beautiful yellow, and at the top a sine purple. The ventral and pectoral sins are placed exactly opposite to each other, upon the thorax.

^{*} De piscibus.

[†] Vide Will. p. 117.

GENUS XXVII .- Remora *.

The remora, or fucking fish of the English, is about eighteen inches in length, almost round. The skin is destitute of scales, smooth, and hard like leather. The head is oval, slattened, and broader than the body: The jaws are of unequal length, the lower projecting beyond the upper; both however are armed with small sharp immoveable teeth, covered by the lips †. The two ventral sins are united by a membrane; the pectoral are of a triangular shape, and two inches and an half in length.

The hind part of the head diffinguishes the remora from every other fish; it extends apparently two inches upon the back; it is quite flat, and striated with transverse lines, like the palate of other fish. It is wish this part that the animal adheres to the belly of the white shark so closely, that the latter when taken is found sticking fast to its skin ‡.

^{*} Remora, Rondel. Echineis Remora, Lin. Syst.

[†] Vide Guan. Hift. pifcium, p. 184.

De Terrey apud Willoughbium, p, 119.

GENUS XXVIII.—Choryphana.

This genus comprehends twelve species that are all exotics: their characters are a large bony head, flat, and floping towards the snout, which appears short and mutilated, as if cut through. The jaws are obtuse, of an equal length, and surnished with teeth; the palate is also frequently exasperated with small spine-like teeth. The back is convex and carinated, and the whole body of a uniform shape.

The Pompilus *.

This fish is said by Pliny, to accompany vessels for several hours when sailing on their voyage, and will not leave them even though threatened with destruction by the sailors. It is an inhabitant of the open sea, and is easily distinguished by a broad curved line, proceeding from the gills to the tail. Above this line the body is variegated with a number of spots; below it, towards the belly, the sides are marked with transverse doted lines. Above the eyes is a large spot of a gold colour. The

[•] Pompilus Rond. Idem. Plinii. † Vide Rond. apud Will. p. 214.

[†] Historia Animal,

fins are four, two pectoral and two ventral, befides a large dorsal and an anal fin, each extending to the tail. From its remote habitation in the ocean, this fish is but rarely caught; and even in the *Italian* markets, where it is fometimes fold, it has no appropriated name.

Another remarkable fish of this genus is the hippurus, also an inhabitant of the ocean, and never found upon our coasts. Rondeletius had an opportunity of examining this animal in Spain, where it is called lampugo, and appeared to him to be distinguished from the whole finny tribe by its superior beauty*. Immediately above the rostrum, there arises a large crest-like fin, which continues uninterrupted to the tail, where it meets with a similar sin upon the belly. The pectoral sins are rounded and broad, resembling ears; the ventral extend the half length of the body. The mouth is moderately large, armed in the jaws, palate, and tongue, with small sharp teeth. The eyes are large; the body covered with small scales, and of a pale blue colour.

The ancient naturalists believed, that this animal lay hid in a dormant state during winter, like the serpent; and that its young increased in size in a far more rapid manner than those of any other sish. The Spanish sishermen, after taking the young, shut them up in pools, where they pretend that a difference of their size is perceivable every day †. The other sishes of this genus are mostly peculiar to South America, and are only known by the barbarous names affixed to them by the natives.

De piscibns. † Aristotle Hist. Animal. lib. 5. cap. 10.


GENUS XXIX. The Goby.


THERE are eight species belonging to this family, that are enumerated by the Swedish naturalist. Their common characters are, the body straight, and gently compressed; the skin rough, with a thick coat of small imbricated and decidnous scales. The jaws of equal length, armed with small, sharp and immoveable teeth. The dorsal sins are two; the first small, consisting only of a few rays, the second broad and high; the pectoral sins are large, and rounded at the ends; while the ventral are united, and have the appearance of a funnel.

There are only two species of the goby known to frequent our coasts, the black and the spotted. The first grows to the size of six inches; the second only three, and is frequently caught in the shrimp nets upon our fandy shores*. Two or three other species frequent the Mediterranean; but as they are of no value, their habits and manners are but little known.

British Zoology.

Plate XIX .







GENUS XXX: The Bull-head:

This deformed tribe is distinguished by a large disproportioned head, much broader and thicker than the body; the visage is rendered staring and hideous, by two large eyes placed above; and the whole of this unshapely head is armed with spines. Some of this genus are inhabitants of the fresh water, while others are peculiar to the sea. There are six species enumerated by ichthy's ologists.

The River Goby *6

This small species seldom exceeds three inches and a half, and is very generally found in all our clear streams, lying at the bottom, either under a stone, or upon the gravel. It deposits its spawn in the sand, in a receptacle dug for the purpose; and its concern for its suture progeny is such, that with reluctance it leaves the spot from which it is to rise. The fresh water insects are its food; the pulex aquatilis being frequently sound in its stormach †.

The

Dottus Gobio, Lin. Syft. Bril-head, Miller's Thumb. Willough, British Zoology.

The back is of a yellow colour, variegated with three or four transverse patches of black. The mouth as well as the head, is disproportionably large: The pectoral fins are broad and rounded; the ventral extremely small, having only four rays. The anal, and two dorsal, are larger than usual in sishes of the same size. There is a species found in the lakes upon the continent, of still an inferior magnitude to our river bull head, and of a brighter colour.

The armed bull-head, or pogge, as it is called in the north of England, is an inhabitant of the fea, much larger than the species noticed, and formidable, as well as hideous in its appearance. It is near a foot in length; the head triangular, and armed with a number of sharp tubercles upon each side; the snout is slat, and turned up, armed also on the point with four strong curvated pikes; below the chin are a number of cirri, forming a copious beard, which adds to the threatening aspect of this animal. The body, from the head, a considerable way down, is of an octagonal shape; near the tail it becomes hexagonal, and is entirely covered with large bony scales, which give the body its angular form *.

Willough. page 137.

Father Lasher *.

This fingular animal is about half a foot in length, and in shape resembling the goby; the head and anterior parts being disproportionably large, the posterior stender as they approach the tail. It has no scales; but the sides are divided each by a rough lateral line. The head is blackish; the back variegated with pale and black patches, placed transversely: Below the lateral lines, the sides are yellow, and whiten as they approach the belly. The coverts of the gills and the crown of the head are beset with formidable spines, by the stroke of which this small sish can insist a painful wound †.

The whole aspect of this fish is threatening; and as it knows the use of its desensive weapons, it can increase the terror of its enemies, by swelling the opercula of the gills and the cheeks to an enormous size. It seems to have been long well known in the Mediterranean, though not for its peaceable and innoxious habits; for it is thus characterised by the poet:

Et capitis duro nociturus scorpius ichu.

The fcorpion threatening wounds with its bony head.

The

Cottus Scorpius, Lin. Syft, Father Lasher, Will,


⁺ Willough. p. 132.

The American species is exactly similar to our own, and is frequent in the Newfoundland seas, where it is called scotling. It abounds also on the coast of Greenland, where it is caught in deep water off these bold shores, and constitutes a very palatable dish to the natives.


‡ Brit. Zool. Class iv. Gen. 22.

Plate XX











SECTION VII.

GENUS XXXI .- The Dorce ..

This species is in all parts of an equal thickness; the body is extremely deep, and strongly compressed laterally; so that in shape it resembles the sounder; but swims creet, and not on the side. The head is stat laterally; the mouth immoderately large, being, together with the head, bigger than the body. The eyes correspond in size with the head; the pupil large and protuberant, and the irides yellow. It is sufficiently distinguished from every other sish, by the round black spot on each side, of about the size of a sixpence; a circumstance from which superstition has made the doree a rival to the haddock, for the honour of St. Peter's touch, when he took the tributemoney from its mouth, by leaving on its sides those incontestible proofs of the identity of the sish, the marks of that Apossle's singer and thumb †.

But if we are to believe the annals of ignorance and credulity, the doree may claim a ftill more ancient origin of the fpots upon its fides; for we have read that the gigantic St. Christopher, in wading through an arm of the fea, over which he carried our Saviour while a child, caught

^{*} Zeus Faber, Lin. Syft. Gallus marinus, Rondel.

British Zoology.

caught a fish of this kind en passant; and as an eternal memorial of the fact, left the impressions on its sides, to be transmitted for the inspection of all posterity *.

The hideous aspect of this fish long banished it from our table, till the celebrated comedian and epicure Quin introduced it. It required only his affertion to make it a delicious fish, and effectually to establish its reputation among the most delicate viands of the times. It is very common in the Mediterranean, the Bay of Biscay, and on the French coast; so when Ovid terms it rare +, it must have been owing to its delicacy, rather that its scarcity.

There is a species belonging to this tribe peculiar to the *Indian* seas, termed by *Willoughby*, faber Indicus, of a very remarkable appearance; the first ray of the dorsal fin extends far beyond the tip of the tail; that of the anal fin is nearly of the same length; while the two pectoral fins seem to consist only of a single ray, and extend almost to the tail ‡.

The Opah ||.

This fish belongs to the genus of dorees, although it far furpasses them in magnitude, weighing in some instances an hundred and forty pounds; measuring three or four feet in length, by two and a quarter in depth. It is so much compressed laterally, that its greatest thickness is only four inches. The colour of the opah is a vivid trans-

Willough, page 295, where Aldrov. afferts, that it is frequently hung up in the Italian churches, fo renowned for fable.

[†] Rarus Faber. Vide Halieut. ‡ Willough. Append. 3.

Opah, or King-fish, Phil, Trans. Opah, Brit. Zool.

transparent varnish of scarlet, burnished over with gold, and bespangled with oblong silver spots of different sizes. The breast consisted of a hard bone, resembling in shape the keel of a ship; and the sless had the taste and appearance of bees *.

The opah has only been four or five times caught on the British shores; and as no mention is made of it by Linnæus or Willoughby, is probably every where a rare fish. The fins, as well as the body, are of fine scarlet; and from the high encomiums lavished by those who have seen it, on its brilliant colours, we are led to suppose that it is one of the most beautiful of the tenants of the deep, and to regret its scarcity t.

Brit. Zoology. † Vide Sibbald, Hift. Scot. Tab. 6.

Vol. III.

Dd

GENUS XXXII .- The Scorpion *.

The animals of this tribe are found in our feas; but owing to the paucity of their numbers, they have no appropriated name in the English language. The species enumerated by Linneus are three. The lesser feorpion resembles a perch in its external figure; it is thick in proportion to its length, and loaded with an enormous head, rendered frightful by the strong spines with which it is beset.

The colour is a darkish dirty yellow, interspersed with brown spots. The skin is covered with extremely minute scales, resembling those of serpeuts. Above each eye is a small sin-like crest, as was observed in the blenny; and at the apertures of the nostrils there is a second pair. Along the back there is one large sin, which, from its lowness in the middle, has the appearance of two. The rays upon the first lobe are sharp and prickly, those upon the second are cartilaginous. The pectoral sins are broad, rounded, and prickly; each ray projecting beyond the membranes, and presenting a large spine. The ventral sins, which are exactly opposite to them, have also a spine terminating the first ray †.

The large fcorpion is about five times the fize of the above; the colour of a deep red, interspersed with dark spots; it is bearded below the under jaw, and has the opercula

²

Scorpana, Lin. Syst. Scorpana Salviani. † Willough. p. 331.

opercula of the gills armed at all points with strong spines. These weapons are of themselves sufficiently formidable to those employed in the capture of this sish; but they have been rendered still more frightful by a belief that they are supplied with venomous matter. Rondeletius mentions a boy who was miserably hurt by attempting to place one in his bosom: he is sully persuaded that the wound was poisonous, and for that reason mentions a prescription delivered by the ancients as its cure. He applied the liver of the sish, with an addition of a bruised lentiscus, to the wound, and thereby prevented those dreadful effects which were imagined to be the unavoidable consequences of a stroke from this animal ||-

|| Rondel, de piscib, art. Scorpius.

Dd2

GENUS XXXIII .- The Flounder.

This genus comprehends the numerous race of flat fish, which keep constantly on one fide, and reside at the bottom of the water, from wanting the swimming bladder. They make progress with one fide forward, and are, on that account, termed pleuronesses by the author of the System of Nature, who has enumerated seven different species belonging to this tribe. The generic characters of the slounder are strongly marked: The body is slat, and one side constantly of a different colour from the other. The head, as well as the rest of the body, is covered with small imbricated scales, and both eyes are placed upon the same side of it *.


The fishes of this genus are many of them excellent food, and found in great plenty in our seas. From their extreme voracity, they are not difficult to take with the hook and line. Sometimes they have been known to swallow the plummet at the end of the founding line, while the sailors were taking the depth of the water †.


[&]quot; Guar. Hift. Pifcium, p. 181.

[†] British Zool, Class iv. Gen. 34.



Plate XXI









The Holibut *.

This species is called in Scotland the turbot, though different from the fish known in England by that name. It is the largest fish of the slounder kind, and is sometimes sound on our shores three hundred pounds weight, while those in the seas of Iceland and Newfoundland greatly exceed even that size. In Greenland they are also caught of a prodigious bulk, with the hook and line, off these bold shores: There they are cut up into large slips, and dried for food to the inhabitants. The holibut is also common in the London market; but where there is such an abundant variety to choose, they are deemed a coarse unsavoury sish, excepting the part which adheres to the side sins, which is fat and luscious.

The colour above is an obscure green, bordering upon black; that of the belly a pure white. The scales are small, and the body all over free from spines; even the edges of the sins have no asperity from the projection of the rays. The eyes are upon the lest side of the mouth, and the right of the animal. The sin which runs along the back, begins above the eyes, and terminates about two inches from the tail; that upon the belly has its origin about seven inches from the point of the rostrum, and terminates in the same manner. There are two pectoral sins and six small ones on the belly †.

Pleuronectes Hyppoglossus, Lin. Syst. Hyppoglossus, Rondel.

[†] Vide Will. in Pifcem.


The Turbot *.

This species is excellent food, and grows to a great fize, weighing from twenty to thirty pounds. The skin is entirely destitute of scales, but is granulated, and has different asperities here and there dispersed over it. The whole upper part of the body, together with the head and fins, is cinereous, and thick set with a variety of black spots. The jaws are not surnished with a single row of teeth, as in the other congenerous sishes, but are exasperated with a vast number of small ones; as also the palate. The cyes are placed on the left side, not so near the edge of the back, nor so close to one another, as in the rest of the slounders. The dorsal fin takes its origin farther forward than usual, beginning near the upper part of the mouth, and extending till it nearly reaches the tail \dagger.

The turbot fishery is carried on to the greatest extent on the north coasts of England and of Holland. They are most successfully caught by the hook and line; the method practised by staked nets being very uncertain. The sishermen of Scarborough are most expert in their business. They go out in large cobles, with three men in each; and every sisher has three lines, surnished with two hundred and eighty hooks apiece. The lines, before shooting, are all three joined together, when they extend about three miles, and are fastened with buoys and anchors. They are drawn at every turn of the tide, the rapidity

^{*} Pleuronestes Maximus, Lin. Syft. Le Turbot, Belon.

[†] Willough. p. 94.





rapidity of the current preventing them at any other time *.

The bait most successfully used for all kinds of ground fish, is herring, which are caught at all times during winter and spring in the Channel. Next to herrings, the leffer lamprey is the most approved bait: The latter are frequently carried by land from Tadcaster, at great expence; for though the flounders of some kinds are extremely voracious, they are so remarkably delicate with regard to a baited hook, that if a piece of herring has been twelve hours out of the water, they will not touch it 十。

Of this genus there are no less than ten species known to frequent our shores: viz. the plaife, the common flounder, the dab, the smear dab, the sole, the smooth sole, the pearl, and the whiff: They are of various reputation as an article of food, and of every fize, from half a pound to three hundred.

^{*} Brit. Zoology. † Brit. Zool, Genus 23.


GENUS XXXIV .- Chatodon.


This genus confilts of a tribe of no less than twenty-three different kinds of fish, all of them entirely exotic. Their generic characters are, an oval body, covered with very minute imbricated scales; a small stat head, with a declining rostrum; a single dorsal sin, covered with scales, and rising high above the back; two small pectoral sins; and immediately opposite, upon the thorax, two sharp pointed ventral ones. The tail is distinct from the other sins, scaly, and in some bifurcated *. As the sishes of this tribe are in general inhabitants of a different hemisphere, we shall not tire the reader with a tedious repetition of barbarous names, where the history of the animals to which they belong is altogether unknown.

[&]quot; Guan. Hist. Pifc. p. 138.



Plate XXIII.









STRIPED WRASSE

SECTION VIII.

GENUS XXXV .- The Gilt-head.

The fishes of this tribe are for the most part inhabitants of the South Seas, where they often relieved the want of the adventurous Captain Cook and his hungry companions. The body is of an oval shape, carinated, and covered with very slender scales: The head is compressed, scaly, and of a moderate size; the jaws are furnished with teeth, which are covered by the lips, the lower projecting beyond the upper. There is but one dorsal, and one anal sin; the pectoral are placed near the belly, and greatly sharpened towards the end: the ventral sins are smaller, and placed somewhat behind them.

The Lunulated Gilt-bead +.

Ins species is an inhabitant of the Eritish coasts, where it haunts the bold, rocky, and deep shores. The whole genus to which it belongs feed chiefly upon oysters, and other shell-sish, which they comminute with their strong teeth before they enter the stomach. The apparatus of Vol. III.

[†] Aurata, Rondelet. Sparus Lunula, Lin. Syft.

the mouth is happily adapted for this purpose; for they are all furnished with flat back teeth, resembling the grinders of quadrupeds, and fulfilling the same office. Besides these teeth, and small sharp ones on the fore part of the jaw, the inner part of the mouth is lined with certain hard bones, which assist in the arduous work or grinding and massicating their hard food *.

They are a coarse fill, and in modern times held in no great esteem, though the caprice of the ancient Romans set a high value upon such as were sed with the oysters of the Lucrine lake, perhaps for no other reason than because they were sathionable eating at a certain period: They seem, from Martial's account, to have thought meanly of these fish in general †.

The lumulated gilt-head grows frequently to the weight of ten pounds: The form of the body is deep, refembling that of a bream; the back is fharp, and of a dufky green; between the eyes is a femilunar gold-coloured fpot; the fides are of the fame hue, but tinged with brown; at the farther angle of the coverts of the gills, there is frequently a black fpot, below it fometimes a purple one ‡. In this fpecies the teeth are covered with lips; and the back teeth are not fo flat as in some others of the genus. There is but a fingle dorfal fin, which stretches a confiderable way along the back, and has twenty four rays; the first eleven spinous, and the rest cartilaginous: The three former rays of the ventral fin are also spinous, and

^{*} Brit. Zoology.

[†] Non omnis laudem pretiumque meretur, Sed cui folus erit concha Lucrina cibus. Lib. xiii. Ep. 90.

Willough. page 307.

the others foft. The pectoral fins are long, terminating in an acute angle; the tail has feventeen rays, and is much forked. In the winter feafon, the gilt-head is caught in great abundance in the *Mediterranean*, and is frequently feen in all the fish-markets of *Italy*.




GENUS XXXVI .- The Wraffe.

Tue filhes of this tribe are distinguished by a flat and oval pody; finall imbricated and resplendent fine; a compressed head; double lips covering their long conic and blunt teeth. The coverts of the gills are fealy; the tail round; and one dorfal fin reaching the whole length of the back, having a flender kin extended beyond each ray; The anal fin is diffinguished by the same addition. The feari, fo famous among the ancients, are of this genus: they were deemed superior to the sturgeon *. Gefner quotes from Galen high encomiums on the falubrity of their flein. Ariflotle, Pliny, Oppian, and Ovid, all concur in a vulgar and unphilosophical notion that was anciently prevalent concerning the rumination of these animals: The two first of these authors do not indeed pronounce with confidence, or affert upon their own knowledge a fact fo extraordinary, but deliver it as a common opinion.


But whatever ideas the ancients entertained upon this fubject, it is so contrary to the analogy of nature, for any of this class of the animal kingdom to chew the cud, that it merits no credit till confirmed by indubitable experiments. After examining the teeth, throat, and intestines of the falmon, tearus, gilt-head, and mullet, and all those afthes that have been considered as ruminating, none of

the

Plate XXIV



GIBBOUS WRASS









the requisites for that process are found to belong to them. Their food also is of such a nature, that after the first comminution in the mouth, it has no farther occasion to be brought back there to undergo a second mastication. Pliny indeed afferts, that the scari cat grass and
weeds; which, if true, would give a shew of probability
to the rumination of these sibes; but unhappily the sact,
in most kinds of these sithes, seems as ill sounded as the
opinion which it is adduced to support †.

Below feems to have adopted the notion of Pliny, as far as the food of the fearus is concerned; for he afferts, that it is only found around the island of Crete, the sea in other parts not producing the marine plant upon which it subsists. Gesner, and several of the ancients, believed that this species, as well as some others, enjoyed intervals of seep in their subaqueous retreats; an affertion which receives no credit from the conformation of the eye; which being deprived both of eyelids and a nichating membrane, cannot possibly be shut ‡.

The Scarus T.

I ms animal is of a reddish livid colour, covered with broad transparent scales, and distinguished from every other sish, by certain prominent and transverse appendages on each side of the tail. The mouth is not very capacious, but stored with obtuse teeth; those upon the fore

[†] Plinii. Hist. Nat. ubi supra. ‡ Gesner de Piscib. p. 1019,

Scarus, Belonii. Labrus Cretenfis, Lin. Syft.

fore part refembling the human, and fitted, as is supposed ed, for cutting the subaqueous plants upon which it feeds. Along the back there is a single sin, exasperated by slender spines. The belly is surrolled with four sins: In this species it is uncommonly large, and silled with juicy sea-weeds. The intestines and liver are taken out, with all the sees, and made into a kind of sauce, with salt and vinegar, with which the slesh of the scarus is dressed and eaten. This is still a savourite repast among the modern Greeks; who, when they set in for a debauch, take large morsels of this dish, in order to give a more exquisite relish to their wine. The scari are caught in nots, into which there are introduced a number of green leaves of a certain shrub, of which these animals are fond.

The Wraffet.

THE usual residence of our wrasse is the deep rocky shores, where it is taken with a bait; though its ordinary food is small crustaceous sishes. Of all sish, this is most liable to vary in colour; in some, it is of a dirty red; in others, it is beautifully striped, especially about the head, with the richest hues of red, blue, and yellow. The lateral lines, opposite to the extremity of the dorsal sin, are incurvated: This sin is supported by twenty-six rays; the first sisteen spinous, and the remainder soft. The pectoral sins are large, and yellow coloured; each

of

^{*} Willough. p. 306.

⁺ Labrus Tinca, Lin. Syft. Turdus Vulgatiffimus, Will.

of the ventral has the first ray produced into a long spine; the three first rays of the anal sin are also spinous *.

The rostrum of the wrasse is protuberant, and bent upwards; and is farther lengthened by two thick sleshy lips, one of which projects upwards, while the other hangs down. The teeth are ferrated, the palate smooth, while the lower part of the mouth is full of prickly teeth.

The wraffe commonly weighs from four to five pounds: In its general figure it resembles a carp. Their slesh is neither palatable nor nutritive. By the French, as well as the English, it is called the old wife, a name of which it is not easy to assign the origin. There are seven or eight distinct species of the wrasse enumerated by Mr. Pennant among the British sides; these, however, are characterised by differences so minute, that a particular description of each would afford but little information in

[#] Willough. p. 320. † Brit. Zool, Class iv. Gen. 25.

GENUS XXXVII. Sciena.

The fishes of this genus are wholly exotic; and are characterised by an oval, compressed and carinated body: The scales are very minute, smooth, and towards the back and belly raised and loose. The jaws are of unequal length, surnished with a great number of sharp, incurvated teeth, that are covered by two thick lips. They have a single dorsal and anal sin, with a loose appendage beyond the rays: The ventral sins are placed on the thorax, a little behind the pectoral, and are sharpened at the point *. A great part of the sishes belonging to this genus are black, and obtained the names of sea-crows, ravens, and umbres, from the old naturalists. Some of them are peculiar to the Brazilian coasts, where they are distinguished by the original Indian names.

The Sea-Grow +.

I'ms species is of a black colour, in shape resembling a perch; the ventral and anal fins are black, as if dyed with ink; the back variegated with undulating lines, of dark

Guan. Hift. pifc. p. 140. † Coracinus, Rond. Umbra, Salviani.

dark brown and blue. The fnout is sharp; the mouth capacious, and inflead of teeth is roughened with fmall asperities: around the rostrum are various small apertures; these of the nostrils wide, and placed near the eyes. On the back are two fins, fo closely adjoining, that they appear fingle and contiguous. The umbre, a frecies belonging to this tribe, is caught in the Mediterranean, and fold in the markets of Rome, where it is ealled ombrind.

Vor III.

GENUS XXXVIII .- The Perch.

This genus comprehends no lefs than thirty-fix different species, very few of which frequent these seas, and their history is on that account but imperselly known. Only five kinds of the perch are found in the takes and on the coasts of Eritain; the river perch, the sea perch, the basile, the ruste, and the black perch.

Some species of this tribe have one, and others two dorfal sins; some have the tail bifurcated, which in others terminates in a straight line. The body is generally oblong, laterally compressed, and covered with hard scales. The jaws are of unequal length, large, and arcuated; the teeth are incurvated, and arranged in several rows. The eyes are large and protuberant, covered with a niclating membrane, and situated at the briak of the forchead †.

There animals are remarkably tenacions of life; fome of them, particularly the river perch, have been carried it ty miles among how, and have farvived the journey. Their fins are to prickly, that they are faid to defy the attacks of the pike: this, however, is only true with regard to the larger perches, if it can be credited at all; for there is no animal which the pike will more readily devour than a finall perch. From the eafe with which the river perch is taken and transported, it has become

P British Zool,

the most common inhabitant of our sish ponds, and affords a very wholesome and palatable food.

The River Perch *.

THE ancients were acquainted with this species; and among them it was deemed one of the first delicacies of the table.

Nec te, delicias menfarum perca, filebo Amnigenos inter pifces dignande marinis. Ausonius,

Rondeletius, and after him Gefner, blames the physicians in his time for ordering the river perch to their patients in febrile disorders, after a prescription of Galen, who meant the sea perch, a fish much lighter, as he alleges, and caser of digestion i. Experience, however, has shewn that this distinction is made without a discrence; both the sea and river kind being sound equally palatable and salubrious. In the time of Willoughby this prejudice against the river perch had been forgotten: He approves of the taste of Ausonius, in deeming the slesh of this animal a great delicacy.

The river perch is easily caught with common earth worms, or small frogs, for a bait; and is so voracious, that the angler, who falls in with a shoal of them, will sometimes kill the whole. This species feldom grows to a large fize, sew being sound above sive or six pounds,

Ff2 the

Perca Fluviatilis, Rond. Une perche de Riviere, Belon.

Y de Rond. apud Will. p. 292.

the greater number being far inferior even to that magnitude. The body is deep and oval shaped; the scales rough; the back much arched; the side line nearer it than the belly. A minute description of a sish so common would be unnecessary.

The Baffe *.

This fish is distinguished by an uncommon degree of voracity, and hence was termed a wolf (lupus) by Ovid+, a name generally adopted by succeeding writers. In the salt water pools of Italy, it sometimes attains to a prodigious size, and weighs sisteen pounds; the sless is extremely grateful to the tasle‡: In the lakes they are frequently sound by the sistermen frozen to death, as they suppose, but more probably suffocated by the exclusion of the air from the surface of the water; a circumstance from which Willoughby takes occasion to caution those who keep them in ponds to break the ice frequently during the continuance of frost §. This species inhabits indiscriminately lakes, rivers, and the sea; to the former, however, they probably, ascend from the sea, for they do not seem to breed in fresh water.

Upon the back there are two fins, both radiated with fpines: behind the anus another rifes and proceeds towards the tail, strengthened with fourteen rays; the three first spinous: The poctoral and ventral fins have each a mixture of prickly and cartilaginous rays. The scales

of

^{*} Lupus, Rondel. Perca Labraz, Lin. Syft.

[†] Halieut. p. 112. ‡ Rondelet. p. 268.

S Ichthyol., p. 271.

of this fish are of a middle fize, very thick fet, and adhering closely to the body.

There are varieties of this species; some variegated, with dark spots, and others of an unmixed colour. The ruste of the English belongs to this genus; it is the small, est of the tribe, and is found in our streams, where it frequents the deepest parts of the water, and is remarkably gregarious; large shoals being found in the same pool. It is of a much more slender form than the perchand seldom exceeds six inches in length.

SECTION II.

GENUS XXXIX. The Stickle-back.

I HIS genus is readily known by the formidable fpines which rife from the back of the greater part of the fish that compose it. There are eleven species enumerated in the fyflem of nature, three of which only are inhabitants of our rivers and shores; namely, that with three; that with ten, and that with fifteen spines upon the back. The former are feen in immense quantities in the fens of Lincolnshire, and the rivers that flow from them, At Spalding, they appear once every feven or eight years, in the Welland, where they mount the river in fuch vast columns, that they are used in manuring the land. An idea of their numbers may be formed, from a circumflance mentioned by Pennant, of a man earning four shillings a-day by taking and felling them to the farmer at an halfpenny a bushel *. The cause of these temporary migrations of the stickle-back, is supposed to arise from the vast quantities washed out of the fens by the floods, and collected in some deep hole, till overcharged with numbers, they are obliged periodically to attempt a change of habitation.

The body is oblong, compressed, and frequently covered with rough scales. The snout is long; and the mouth thick set with small granulated teeth. There is one dorsal, and one anal sin; the pectoral are small, and sharp pointed, situated upon the thorax, which is covered with scale or bony plates.

The Three-Spined Stickle-Back *.

THIS species is broad, and compressed laterally: it feldom exceeds two inches and a half in length; is of an olive green upon the back; the belly white: In fome the throat and belly are red. Before the douful fin there are three sharp spines which can be raised or depressed at pleasure; behind the anus there is one, fometimes two, fill ftronger, which are fixed to a bony plate. The tenspined stickle-back is of a still smaller size, and a more Hender form; the fides are finouth, not plated as in the preceding species. The fishers affert, that this animal is produced spontaneously, and from them tithes of various kinds: The foundation of this extravagant notion, is their observing them in ponds newly dug, where, in the course of a few years, different fish have been found. We have already affigned our reasons for these appearances, without having recourse to such an ill-sounded hypothesis +.

The Fifteen-Spined Stickle-back t.

SCHONFELDT first described this animal, which is an inhabitant only of the sea, and is much larger than any of the

^{*} Gasterosteus aculeatus, Lin. Syst. Stickle-back, or Sharpling, Will.

^{&#}x27; Vide Chap. i. fect. 4. Of the generation of fishes.


¹ Açulcatus marinus major, Will. Gafteroftous Spina hia, Lin. Syft.

two preceding. The body is flender, being only an inclithick, and nine long; towards the tail it is quadrangular, and extremely small: The whole skin is smooth; black upon the upper part, and filvery upon the lower. It has two pectoral fins, one dorfal, rising in a triangular form from the middle of the back; between which and the head there are fifteen unconnected spines, inclining a little towards the tail; on the middle of the belly there are two, at the anus one: The fins of the belly are wanting. The species inhabits the Baltie and German ocean, particularly the coasts of the Netherlands; it is called ersskrupper. Willoughby is midaken in supposing, that it is never found on the coasts of Britain*.

^{*} Ichthyolog: p. 3401



Plate XXV.



LOCHE

GENUS XL .- The Mackrel *.

This is one tribe of those shoals of migratory sishes, which annually visit our coast; and it is perhaps among the most celebrated of that class; both for its numbers and the delicacy of its food. They are gregatious in their habits; and while other sishes continue in our vicinity all the year round, and proul along the shore in small numbers, or solitary, these in a compact band visit their accustomed haunts at stated seasons with so much regularity, that you can pronounce with certainty the time of their appearance.

What impels the mackrel and other migratory fishes to undertake such distant voyages, what directs their course, and what supports them upon their passage, are subjects on which naturalists have indulged in all the licence of conjecture, without being able to form any satisfactory determination. Some have imagined, that we owe the visits of the whiting, the haddock, and the mackrel, rather to their sears than their appetite; and that they are driven upon our coasts in endeavouring to avoid the pursuit of their destroyers. It is, however, more probable, that they approach the shore in quest of food, which is found in greater plenty there than in the depths of the seat.

Vol. III.

Gg

Tt

^{*} Scomba, Rondel. Scomba, Scomba, Lin. Syft.

[†] Goldsmith's Nat. Hist. vol. vi. p. 326.

It is to be regretted, that the mackrel, though perhaps the most palatable of all the gregarious sisses, is least useful, owing to its extreme tenderness, and inaptitude for carriage: It is capable, indeed, of being preserved by pickling and salting; a method, which we are surprised to find is only practised in *Cornwall*, where this food proves a great relieve to the poor during winter *.

The ancient Romans were just to the merits of this fish; for among them it was held in high esteem, especially on account of the garum, a celebrated pickle, that gave a high relish to their sauces, besides being medicinally useful. According to Belon, this precious pickle is still in sashion at Constantinople. Formerly it was prepared from different kinds of sishes; but that procured from the mackrel was deemed preferable: The best was made at Carthagena, vast quantities of mackrel being taken near an adjacent isle, called from that circumstance Scombraria 1.

The mackrel is eafily taken, by a variety of baits; but the capture always fucceeds best during a gentle gale of wind: a coloured feather, or a piece of scarlet cloth, is often found sufficient to allure these dim-sighted animals to destruction. In spring their eyes are covered with a white silm, which renders them half blind, till it is cast some time in the summer.

This fish is probably found in all the European seas, being universally known in that quarter of the globe. It feldom exceeds two pounds weight in the common species, and its body is of a very elegant form, and beautifully variegated

^{*} Borlases Cornwall, p. 269.

[†] Plinii, Lib. xxxi. cap. 8.

variegated with the finest hues of green, blue, and silver*. The common mackrel is from a foot to a foot and a half; the body thick, round, and sleshy, but tapering to a slender point as it approaches the tail, which is bifurcated. The scales are so small as to be hardly perceptible; behind the anal and second dorsal sin there are several protuberances or spurious sins, both above and below †: These may be deemed the distinguishing marks of this animal, which is too well known to require a more particular description.

The Tunny t.

This species is well known in the Mediterranean, which they annually enter from the Atlantic ocean; and are taken by the inhabitants of the coasts of Spain and Italy as they advance. They were supposed by the ancients to breed in the Levant, and Palus Mætis. Some of them grow to an immense size, weighing above four hundred pounds; a magnitude which seems altogether incompatible with the short life which the ancients allotted to these animals: Both Pliny and Aristotle affert, that two years is the utmost period of the life of a tunny §.

A very extensive tunny fishery was carried on by the ancients in the Mediterranean; particular stations being G g 2 fixed

British Zoology, gen. 29. † Willough. p. 181,

⁷ Thunnus, Rondel. Scomber Thunnus, Lyn. Syft.

⁷ Hist. Anim. Lib. ii. cap. 13.

fixed upon for that purpose, near some promontory, where a man was placed to give notice of the motions of the sish, and warn the sishermen below. It is from one of these precipices, that the lover in Theocritus threatens to leap, with a view of softening the rigour of his mistres's cruelty. The manner of taking the tunny in nets, as it is at present practised, is perhaps similar to that of the ancients; many of the sishing stations are exactly the same. The trade is still carried on to a great extent: when taken, the sishes are cut into pieces and salted up in barrels, in which state they are exported to all the neighbouring countries. The abdomen is reckoned by far the most delicate part of the sish: it is cut separately, and sold to the nobility and wealthy citizens of Kome, under the name of taremtello.

The tunny frequents the English coasts, but not in the fame numerous shoals that are seen in the Mediterraveans. It is also pretty common in the locks on the western coasts of Scotland, where it comes in pursuit of herring, and often during night strikes into the nets, to the great damage of their owners. In the morning, when the nets are drawn by the sishers, the tunny watches near them, for any sish that may chance to drop out. The sishermen avail themselves of this avidity for prey in the tunny, and baiting a hook with a herring, they seldom sail to take him.

The flesh of the tunny, when recently taken and cut up, has the appearance of raw beef; but after boiling, it turns pale and tastes like falmon. They are about eight feet

These were termed Θυγιοςκοπεία.
 Vide Salvian. apud Will.

[†] Vide Idyllia. § British Zool.

feet in length, the body round, thick, and fleshy; but growing slender towards the tail, where it has an angular shape. The two sins on he back are each strengthened by sourteen strong spines; those of the first are capable of being depressed, and folding so closely into a sulcus in the back, that they become imperceptible. Behind the anal and second dorsal sin, there are a number of small spurious ones both above and below.

GENUS XLI .- The Surmullet.

This genus comprehends the three species of mullets; all sishes of the most delicate kind: some of them were in such request among the Romans, that they were bought by the private citizens for their weight of silver. It was in the purchase of this sish, above all others, that the wealthy senators showed their extravagance and sensuality. Juvenal and Pliny afford ample testimory of the luxury of their age, which, happily for mankind, has never been equalled in any other period of human society. An instance of sixty-sour pounds being said for a sish of three pounds weight, is recorded by Pliny.

But the manner of dreffing and eating these fish, still more than the extravagance of the price, shewed the epicurism of the ancient entertainments. The mullet was not reckoned worth a tarthing if it did not die in the hand of your guest. There was water kept in the eating room into which the living animal was put, and from thence conveyed immediately to the stew, where it was dressed, which was also in the same apartment, and under the table; from thence the fish was placed upon it.

It

^{*} Mullum fex millibus emit, Æquantem fanc paribus fishertia libris. This fum is equal to 481. 8s. 9d. Juv. Sat. xv.

[†] That of Asinius Celer, a man of consular dignity, Vide Lib. xx,

It was customary to put the mullets into glass vases, that the company might be entertained with the various changes of their rich colours as they lay expiring *. It was Apicius, that prince of gluttons†, that first hit upon the ingenious invention of suffocating them in the exquisite Carthaginian pickle, and afterwards procuring a rich sauce from their livers.

The Striped Surmullet \$.

There are three species of this genus, the red and the striped, which are sometimes sound in the British seas, and the king of the surmullets, a sish peculiar to the Mediterranean, and destitute of the two cirri which hang from the chin of the other species. The striped surmullet is about sourteen inches in length, and weighs about two pounds and an half. The body is thick; the snout blunt, and from the lower side of it hangs two beards, about an inch and an half long. The sins are of a beautiful red, and yellow; the colours of the body vary in their hues, into a thousand different shades, while the animal is dying.

^{*} Vitriis ollis inclusi offeruntur, et observatur morientium color, quent multas mutationes luctante spiritu vertit. Seneca, Nat. Quæst. lib. 3.

[†] Nepotum omnium altissimus gurges. Pliny, lib. x. c. 48.

[!] Mullus major, Will. Mullus Surmuletus, Lin. Syft.

GENUS XLII .- The Gurnard.

The distinctive characters of this tribe, called by Lineaus trigla, are, a scaly body, of an uniform shape, compressed laterally, and attenuated towards the tail. The head broader than the body; and sloping towards the snout, where it is armed with spines; the upper jaw divided, and extending beyond the lower. The eyes are far from the rostrum, near the top of the head large and prominent, particularly the upper margin of the orbits. The two dorsal sins are unequal; the first short, high, and aculeated; the second, long, sloping, and radiated. The ventral and pectoral sins are uncommonly large; and from their base there hang three loose and slender appendages*.

To this genus belongs the flying fish of the failors, or trigla volans of Linnaus, remarkable for the length of the pectoral fins, which it fometimes uses for the purpose of flying. There are two species of the gurnard that participate in this manner the powers of the feathered race; they are both found only in warm latitudes, and nature has probably vouchfased them this endowment, to enable them to escape from their numerous pursuers below. It is a curious spectacle in the intratropical regions to behold a thousand of these fishes, all darting

darting from their native element; and while they endeavour to avoid the purfuit of the dolphin below, falling into the power of enemies still more dextrous and inveterate in the aereal element. The slights of these ambiguous animals are but short; for as soon as their wings dry, they drop back into the subjacent waves without a possibility of rising till they are again wetted there: It is from the sudden drying of their wings, that they often fall upon the decks of vessels, particularly those that double the Cape of Good Hope. These sishes are seen in the Mediterranean, but seldom venture farther from the line; none of those that frequent the British seas are capable of rising from their native element *.

The Gray Gurnard +.

The colour of this fish, upon the back, is a dirty green, variegated sometimes with black, but more frequently with clay-coloured spots: Below the lateral line, which is remarkably rough, the colours are more diluted, and the white spots more frequent, till you approach the belly, which is silvery. The head is large, covered with bony plates, of which the highest that covers the top of the head, runs backwards, and terminates in two sharp horns or spines. The snout, and upper part of the eyes, are also fortissed in the same manner by spines. The Vol. III.

^{*} Vide Willough. et Marcgr. de piscib. Bray.

[†] Gurnardus Grifeus, Will. Trigla Gurnardus, Lin. Syft.

jaws, and the whole inner part of the mouth, are covered with very small teeth. The eyes are large; the irides of a filver colour, with a tincture of brown: The nostrils are fmall, and placed on the fides of the rostrum: On the extremity of the gill covers, is a thick, fharp, and long fpine. The first dorsal fin confists of eight spiny rays; the fides of the three first tuberculated: The second has nineteen rays, and is lodged, like the former, in a groove, rough on each fide. The pectoral fins are long and tranfparent, and supported by ten rays; from their base hang three beards*. The British fishes of this genus are, the tub fish, the feraphine, the streaked, the gray, and red gurnard, and the piper; the last, Rondeletius has termed the lyre, on account of its bifurcated roftrum, which fomewhat refembles that instrument among the ancients. The whole of this genus is excellent food.

British Zoology, gen. 31.

DIVISION IV .- Abdominal Fishes.

SECTION X.

GENUS XLIII .- The Loche.

We are now come to the last division of the spinous sistes, termed by naturalists, the abdominal, from the situation of their ventral sins, which are placed far behind the pectoral in the abdomen. The loche, or cobitis of Linnæus, which is placed at the head of this tribe, is perhaps the most unshapely of the whole race. The body is compressed laterally, and nearly of the same depth from the head to the tail: It is destitute of that symmetry and elegant tapering form, which is characteristic of the greater part of sishes. The scales are deciduous, and so small as scarcely to be perceived. The back is marbled with brown spots, upon a paler ground of the same colour; the belly, a dirty white ...

Linneus has five species, that are peculiar to our rivulets. The loche seldom attains to the fize of four inches; and is rendered hideous by a beard that projects all round the snout. Forbidding, however, as this animal may seem, we are told our sportsmen sometimes swallow it down alive in a glass of wine †.

Hh 2

The

The body is smooth and slippery, and almost of equal thickness; the two dorfal and caudal fins are marked with small spots, the latter somewhat rounded, and barred with stripes of black. This disgusting fish, which constantly crawls at the bottom of our rivers, is by some reckoned a delicate food, when made into a kind of sauce. They generate commonly in the month of April, among the long aquatic weeds about the brink of the rivers †.

[†] Willough. page 265,

GENUS XLIV. Amia.

Or this genus there is only one species, distinguished by the baldness of the forehead, which appears as if excoriated. The body is stender, a little compressed, and covered with imbricated and slexile scales; the head is bony, rough, and slattened; the jaws of equal length, and surnished with a number of sharp erect teeth; the tongue is obtuse; the palate armed with two teeth; near the eyes are two apertures of the nostrils; and not far from them are two beards.

There is only one dorfal fin rifing from the middle of the back, radiated and floping; the ventral and pectoral fins are small; the former placed behind the equilibrium of the fish, in the abdomen. The tail is rounded, supported with rays, and short ||.

|| Guan. Gen. Piscium, p. 208.

GENUS XLV. Silurus *.

A VARIETY of exotic fishes are contained in this tribe; that which we select for description is found in the Danube and Vistula, sometimes in the Rhine, and the lakes of Germany. It grows to the length sometimes of sixteen seet, and weighs an hundred and sifty pounds. At Vienna, Willoughby observed them of this prodigious bulk †. The belly seems as if worn by continual rubbing on the ground, whence it is probable that the silurus frequents the bottom of the water. It is extremely voracious, and destructive to other fish wherever it appears: by many it is commended as an article of food, when skilfully dressed.

This animal refembles the cel in colour, and that vifcous substance with which the body is covered. The belly and sides are variegated with large spots of black and white, and the whole is destitute of scales. The head is broad, and much flattened. When the jaws are open, the gape of this sish is tremenduous; for it extends almost the whole length of the head. Before the eyes there are two antennæ, resembling the scelers of insects, and probably destined to serve the same purpose. From the under lip, there hang four similar seelers, which the animal is said to cast annually ‡.

The

^{*} Silurus Rondel. The Sheat-fish, Will.

f Ichthyology, page 128.

† Vide Gesnerum apud Will.

The dorsal fin is small, having only three rays; from the anus to the middle of the tail, there is a very long fin, supported by twenty rays; the pectoral fins have each a strong spine projecting from the first ray, which the animal probably uses as an instrument of defence.

GENUS XLVI .- The Touthis.

The two species which Linnens has arranged under this genus are both exotic filles. The generic characters are, a cylindrical body, covered with tuberculated scales, which give the animal somewhat of an angular shape. These scales are thinly set, scarcely imbricated, and are extremely rough to the touch. The head is broad, slattened horizontally, and sloping towards the shout, which is obtuse. The mouth is destitute of teeth; the eyes distant from the shout, placed almost on the top of the head, and covered with an entire nichating membrane. On the inner side of the lips are innumerable cirri, which serve the animal instead of teeth *.

A Bravilian fish, termed by the natives guacari, belongs to this genus. The fins are in general supported by very strong spines. The tail, which in the lower division is about three inches in length, and two in the upper, is supported by very strong spines. The whole head is covered with a rough kind of shell; the scales of the body triangular, and arranged in rows, forming a strong coat of mail, well adapted for defence †. The whole body of this sish is of a sasron colour, diluted towards the belly, but deep upon the sides and back; over this ground there is dispersed an innumerable quantity of small brown spots, resembling mustard seeds. The native Brazilians eat this sish; but after the bulky tegument is subtracted, very little sless remains, and that of no exquisite quality.


^{*} Guan. Hist. Piscium, p. 188.

[†] Willough. page 277.









SALMON TROUT



TROUT



GENUS XLVII. - Salmon.

This family contains feventeen species, many of which are perfectly familiar to the inhabitants of this island, being found in all its lakes, rivers, and shores. The most remarkable for beauty and utility is the common salmon, whose external figure may vie in elegance with that of any fish that swims, and whose shell known over all Europe for its agreeable take and nutritive quality.

A minute description of the parts of the common salmon would be superfluous. It is found over all the north of Europe and Asia, from Britain to Kamscoatka*; but does not venture into the warm or intratropical latitudes. It is frequent as far north as Greenland, and is seen in some of the rivers of France; but none have been caught so far south as the Mediterranean.

The falmon is an ocean fish, and therefore is only found in those rivers and lakes that communicate with the sea. As the ancient Grecians never extended their empire to the ocean, they were unacquainted with this excellent fish; and hence their language has no name to distinguish it ‡. The extent of the Reman territory gave them many opportunities of being acquainted with the salmon; and in their language it has a name from which ours is borrowed; the word seems to express its bounding or leaping [].

Vot. III.

Ιi

Among

^{*} Hift. of Kamsch. p. 143.

[†] Rondel, Fluvial, 167.

f Gefner, apud Will. p. 189.

Among feveral nations the falmon has different names, according to the age of the fish. The ancients diffinguished them into falmones, fariones, and falares, according to their age *. In Ingland, the inhabitants of York-fibire call those of the first year, smelts; of the second, sprods; of the third, mosts; of the fourth, fork-tails; and of the fifth, half sish; for it is not till the fixth year that this animal attains its full size.

There are falmon-fitheries in Iceland, Norway, and Labland+; but no where are they carried on to greater extent than in Britain and Ireland; the fifhery of Colvaine in the latter kingdom being an inexhaustible fund of provision and of wealth.

The long migrations of the falmon at the fpawning feason is, perhaps, the most extraordinary part of its history. It annually mounts rivers, sometimes to the distance of several hundred miles, till it sinds a place in their gravelly bed, sit for depositing the seeds of a future family. There are hardly any obstructions which it will not overcome, to attain this purpose. It easily makes its way against the most rapid current, and will leap up caracasts of an association height. In endeavouring to surmount these perpendicular falls, it often fails in the attempt, and tumbles back into the water, and thus is in some places caught in baskets placed below the current.

The manner of depositing their spawn, after having personned these ardnous journies, is curious. When a proper place is selected for the purpose, the male and se-

ž malė

^{*} Teque inter geminas species neutrumq. & utrumque, Qui nec dum Salmo, nec jam Salar, ambiguusque; Aniborum, medio Fario, intercepte sub zvo. Ausón;

⁺ Hackluyt, Voy. I. 416.

made unite in digging a pit of eighteen inches deep, into which the former ejects his smelt, and the latter her spawn: The place is carefully covered up with their tails, which at that period are worn bare. In this state, the spawn lies buried from December till Spring, if not disturbed by violent floods; while, in the mean time, the parent sishes hasten back to purify and strengthen themfelves in the sea; for, after spawning, they are lean and slabby. It is when they first leave the sea that they are plump and vigorous; and such as are in that condition have a number of insects adhering to them above the gills *.

About the latter end of March, the ova begin to exclude the young falmon, which are then refuscitated, and fwarm in numbers, of which it is difficult to form any idea. They are then termed fry in Scotland, and fmelts in the fouth of the island; and they remain in the fresh water till the arrival of a slood, by which they are swept down into the sea. In June, the young saimon begin to appear again in the rivers, in dropping single sish: there they increase with amazing rapidity in size; for in June, when they appear first in the rivers, they are only about a foot or sourteen inches in length; and in August some are seen eight and nine pounds.

The quick growth of the fal non is by no means occasioned by the voracity of these sich; for all the sithermen agree, that no food is ever found in their stomach. Perhaps this may be owing to a temporary neglect of food about the season of spawning or secundation, as is the case with the seasons. It is well known that the

I i 2 angler

[?] The lernææ Salmoneæ, Lin. Syft.

angler takes them faccefsfully with the fly, the worm, or fmall fills, an evident proof that there are the animals on which they feed.

At Berwick on the river Tweed, the capture of the falmon is immense during the month of July. At that senson, they are salted, and barreiled up for exportation: Even the immense consumption of London not being able to keep pace with the vast leads that are daily taken.

Upon the Tweed there are above forty different filheries, extending about fourteen miles up the river, which are rented at the fum of 5400 l. annually: The expence of fishing-tackle, servants wages, and other requisites, amount nearly to an equal sum; so that upwards of 10,400 l. must be obtained by the different lesses, before they can clear their expences; and hence, 208,000 salmon must be caught there, upon an average, every year, before any profit can accrue to the tenants.

The fishing feason commences on the 30th November, and ends on Michaelmas day. The corporation of Berwick are conservators of the fishery, part of which belongs to that town. The river being the boundary of the two nations, the whole north fide is Scotch property*.

The rivers both of Scotland and Ireland abound with these sish; and a pretty extensive capture is made in some of them. The mode of preserving them is nearly the same with that practised in Berwick, and the markets are similar; for in these kingdoms they begin to export during the summer months, after the London market is supplied. Very severe laws were enacted in Scotland for the preservation of salmon. So late as the reign of

'James

^{*} Vide British Zoology, Gen. xxxiii.

James IV. killing this fish was made capital after the third offence; before that period, the culprit might redeem his life *.

Various causes have of late operated to the prejudice of the British trade in this important article. The great increase of the Newfoundland sithery has reduced the price of salmon in the foreign markets, while the gradual increase of knowledge throughout Europe has led to a much greater relaxation in the discipline of the Romish church, and particularly in the article of abstinence; a regimen which is daily losing credit as a spiritual restorative. And to these circumstances we may also add, the great improvement in agriculture that has every where taken place in these kingdoms within the last sifty years; which, by the frequent application of lime upon the foil, has probably greatly diminished the number of salmong as well as of other river sishes.

The Sea-tront to

This species is larger than the common, and disters from it in its habits and colour, as well as in fize. It annually migrates like the salmon, from the sea; and after having deposited its spawn, it again returns to its savourite refidence in the ocean. The shape is more thick than that of the river-trout, and its weight sometimes above three pounds.

^{*} Vide Regiam Majest. Rob. III. cap. 7.

[†] Salmo Trutta, Lin. Syst. Trutta Salmonata, Will,

pounds. It differs from the falmon, in having the tail terminating in a flraight line. The head is thick, fmosth, and dufty; the back is of the fame colour, but grows fainter towards the fides, which are marked with large irregular fpots of black. The field, when boiled, is paler than that of the falmon, but grateful to the tafte.*

Another species, or perhaps variety, of the trout, is the samlet, the so-allest of the kind, and by some accounted the fry of the salmon at a particular stage of their growth: but as this species remains in the rivers after the departure of the young salmon, and is of a much smaller size than they are after their return, the inference is obvious, that it must belong to a different tribe. It nearly resembles the river-trout; but may be distinguished by a broader shape; by its superior whiteness, and by the shape of its tail, which is more sorked; and, lastly, by the smallness of its size, which is much inscrior to that of the trout †

The River-Trout 1.

A HE common trout is a fish universally known in this island, for the delicacy of its slesh, for the sport it affords to the angler, and for the superior beauty of its colours. The common species it is unnecessary to describe; but there are some varieties not unworthy of notice. The colours of the trout vary greatly in different waters: lu Llyndivi,

Willough. † Brit. Zoology, Gen. 33. ‡ Salmo Fario, Lin. Syft. A Trout, Will.

Myndivi, a take of South Wales, are trouts marked with red and black fpots, as large as fixpences. In Loch Neagh in Ireland, and Hulfe Water in Cumberland, there are trouts taken which weigh about thirty pounds. There are probably of the fame species with the celebrated trout of the lake of Geneva. The fize alone of these fithes can recommend them; for in take and flavour they are all inferior to our common river-trout.

It is remarkable that neither the Greek nor Roman authors make any mention of this fish, and that a species of such delicacy should never have become a fashionable dish. In the Italian rivers the trout abounds, and in all probability did so at the time when the Romans were transporting oysters from Sandwich*, and the livers of the sear from the coast of Atrica +. But it was perhaps owing to its proximity of situation, and to the ease with which it might be procured, that the trout was despited by the votaries of Epicurus, who scrupled not to lavish a large fortune on a dish of slammingo's tongues, and other insipid rarities ‡.

It is the veracity of the trout that leads the way to its destruction, and makes it a prey to the inventions of the angler. The passion for angling is so great, that in the neighbourhood of London, ten pounds a year is the sum paid for the liberty of shing in some of the adjacent rivers.

But there is a finaller fish in some parts more plentiful than the trout, which affords superior diversion to the sportsman, and that is the parr, a fish of remarkable beauty, and elegant proportions. This beautiful species

is

^{*} Juven. Sat. iv. 141.

[†] Suctonius in Vita Vitellii.

Martial, Lib. xii. Epig. 71.

is a well known inhabitant of the greatest part of those pure and rapid streams which descend from the mountains in the *Highlands* of *Scotland*, where we have witnessed above twelve dezen taken with a single red and line in a few hours.

The Charres:

This species is very properly denominated the Lipino charr by Liminus; for its constant residence is in the lakes of the high and mountainous parts of Europe. A few are found in some of the lakes in Wales, and in Lock Inch in Seculard; from which last it is said to migrate into the Spey to spawn: Seldom, however, does this species venture into any running stream; its principal resort is in the cold lakes of the Lapland Lips, where it is fed by the innumerable larves of gnats that insess those dreary regions. When the Laplanders migrate to the distant lakes during summer, they find a ready and luxurious repast in these sithes; which to them are extremely palatable without any sauce: accustomed to temperance, and exercise, these hardy natives are independent of the inventions of epicurism;

The largest and most beautiful charrs are found in the lake of Winander-Mer, in Westmoreland, where there are three

^{*} Salmo Alpinus, Lin. Syft. Umbla minor, Gefner.

⁺ British Zoclogy.

ş

Plate XXVII.









three species, the red, and the gilt, and the case-charr. These kinds are nearly similar in their external appearance; but the time and manner of their spawning are so different, as to afford room for their specification into three distinct samilies. The case-charr spawns about Michaelmas, and chiefly in a river that runs into the lake called the Brathy. The spawning season of the gilt charr is from the beginning of January to the end of March: they never ascend the river; but make choice of a springy part of the lake, where the bottom is smooth and sandy *.

The figure of the charr is more slender and lengthened than that of the trout: The colour of the back is an olive green, variegated with spots of a dusky white, and others of a dull yellow. The belly is of a pale red, in the female greatly diluted, and approaching to white. The whole body is covered with very minute scales † The manner of taking these fishes is with nets or trammels, as they are called, which are furnished with bait to allure the fish, and left set for several days, till they are known to enter them.

The Gwiniad &.

This fifth feems to be an intermediate species between the trout and the herring; like the latter, it suddenly Vol. III. K k dies

^{*} Brit. Zool. † Willough. p. 196.

[§] Salmo Lavaretus, Lin. Syft. Lavaretus Allobrogum, Will.

dies when taken out of the water. In length, it is nearly a foot, laterally compressed, has two fins upon the back; the first about three inches from the snout, and supported by thirteen rays; the second, which is placed near the tail, is thick, fat, and altogether without rays. It is this spurious sin which Willoughby assumes as the generic mark of all the sishes of the trout kind *.

This fish, like the charr, is of an alpine nature, and inhabits the lakes of the mountainous parts of Europe; fuch as, Switzerland, Savoy, Norway, Sweden, Lapland, and Scotland: Into the last mentioned country, it is said to have been introduced from France by the unfortunate Queen Mary; a circumstance rendered probable by the French name vangis, given it by the inhabitants in the vicinity of Lockmaken. The British name of gwiniad is bestowed upon it from the whiteness of its scales. The gwiniad is a gregarious sish; and approaches the shores in vast shoals in summer and spring, which are the seasons in which it is taken: In some of the German lakes it is caught in vast abundance, and salted up for sale in the neighbouring towns. The slesh is insipid, and will not preserve for any time without salt.

There are twenty-nine different species of fishes belongeing to this genus, of which nine species, with their varieties, are all that is known in *Britain*.

Vide Ichthyol. p. 1833



Plate XXVIII.






SEA PIKE



SAURY



ATHERINE



ARGENTINE

SECTION XI:

GENUS XLVIII .- The Pike.

This tribe contains in it four species of British fishes; namely, the common pike, the sea pike, the gar, and the saury: They are all distinguished, by having only one dorsal sin near the tail; by a long, slender body, laterally compressed; and by the length of the lower jaw, which projects considerably beyond the upper. In general, the snout of this samily is long, and slattened horizontally; the scales are small, bony, and deciduous *.

The most remarkable part of the manners of these sistheir voracity. Almost every kind of animal that is not two large for its swallow, is indiscriminately devoured by the common pike: even those of its own species are sacrificed to its gluttony, as often as other food is wanting to fatiate its ungovernable appetite. Two instances are mentioned of its being choaked, by attempting to swallow animals too large for its power of deglutition: In the one, the animal expired in endeavouring to devour a pike nearly of its own size; the other happened at the canal at Trentham, where the pike feized the head of a swan, as she was feeding under water, and gorged so much of it as killed them both †.

K k 2 All

[·] Guan. Hift. pifc. gen. 43.

All writers agree in afcribing uncommon voracity to these animals, and mention sacts in support of
it so extraordinary, as would seem incredible, were
they not in a great measure consirmed by daily observation. The peculiar structure of the jaws, which
are loosely connected together, and have on each side
an additional bone, like those of the viper, while it
savours their voracious instincts, proves often their ruin,
by prompting them to swallow morsels too large, even
for that uncommon distension of which they are capable.

The digestive powers of the pike are as remarkable as its voracity: After swallowing a fish of a size but little inferior to its own, those parts that have entered into the stomach are dissolved with amazing rapidity; while those in the mouth and throat, which are yet entire, make a constant progress downwards, as the process of digestion makes way for them. Hence a fish, which is too large to be swallowed entire, appears for a while with its tail hanging from the mouth of the pike; but soon after totally disappears, and is dissolved by the stomach almost as quickly as it enters there **.

From this extraordinary voracity of the pike, aided by fuch uncommon powers of digeftion, he is by far the greatest tyrant of the fresh water. He is said to contend with the otter for his prey, and sometimes to force it from his mouth. The angler, in drawing a trout, has been known at once to lose his line and his prize, by the unexpected attack of one of these plunderers. When we consider the numbers and voracity of these animals, it may appear strange that any of the inhabitants of the

fresh water should escape their devastations. Young geese and ducklings, when they first venture into the ponds, are often destroyed by them: and all the smaller sishes shew the same terror at the appearance of the pike, as the little birds do at the sight of the hawk or the owl*.

The devastations committed by the pike are considerably increased, by the great longevity of that animal: If the accounts of naturalists can be credited, the period of its existence far exceeds that of every other sish, not excepting those of the cetaceous kinds. The Polish naturalist Reaccynski mentions one that reached its ninetieth year; and Gesner gives a print of a brazen ring, that had been assixed to one that was caught near Hailbrun, in the year 1497. On it were inscribed these words in Greek characters; I am the fish that was first of all put into this lake by the bands of the governor of the universe, Frederic II. in the 5th of October 1230. According to this account, the sish must have been no less than two hundred and sixty-seven years of age, a fact too extraordinary to be received on the evidence adduced for it.

The generation of the pike, from its being found in ponds, where none were ever introduced, has been supposed as extraordinary as its longevity. Nothing, however, seems more easy than to account for these facts, on the well known principles of the generation of sishes: If a heron hath devoured their ova, and afterwards excreted them while sishing on one of these ponds, it is highly probable, that they may be produced from this original, in the same way that the seeds of plants are known to be

^{*} British Zoology, Gen. 32.

[†] Hift. Nat. Pol. p. 152.

Vide Icones piscium, p. 316.

differinated *. When the female pike is about to fpawh, the is faid to withdraw as much as possible from the common haunts of these sishes, that she may conceal her brood from the depredations of the rest, it being a well known sact, that the male of many species pursues the female when about to deposit her ova, and on the first opportunity devours them.

When the pike is in feafon, its colours are a very brilliant mixture of green, with bright yellow fpots; the gills are then of a vivid and full red; when out of feafon, all these colours decay; the green becomes gray, and the vellow fpots grow pale. It is only when in feafon that their flesh is eatable: it is then white, firm and palatable food. Gefner relates a most extraordinary method of expoling them to fale in England, which feems as little entitled to credit as his account of their longevity. He heard, he alleges, from an eye witness, that the animal's breast was cut up about two inches, to shew its degree of fatness, and if no purchaser offered, the wound was infantly fowed up, and the fish again committed to the pond, where, by being rubbed with the glutinous matter on the body of the tench, it was from cured +. In fome parts of England the pike was formerly fed in small perforated boxes of wood, fixed by chains to the banks of the river. Such floating chefts Willoughby declares he has frequently feen in a river near Canterbury.

The feason of generation among the pikes is in March or April, according to the heat of the weather: their fertility is extraordinary; an hundred and forty-eight thousand

^{*} Willough, ubi fupra.

[†] Vide Gefner apud Will. p. 238.

thousand ova being taken from a fingle fish caught in the Rhine, which did not weigh above nine pounds *.

This species of fish is common in the lakes, rivers, and ponds, over the whole of the north of Europe: It is, however, most numerous in Lapland, where pikes are said to be taken sometimes eight feet in length; and their expertation for sale forms a considerable article of commerce. The pike abounds in most of the lakes and rivers of Britain, and in such numbers as renders the common account of this sish being introduced into England in the reign of Henry VIII. extremely improbable †.

The American pike, or Efox offcus of Linnaus, feems far more abundant in the lakes of Canada than ours, and forms there a most valuable article of food. It grows commonly to the fize of four or five feet; it is covered all over with bony rhomboidal scales, which render it almost invulnerable; but is, notwithstanding, taken by the Indians in such quantities, that it forms no inconsiderable portion of their sustenance.

^{*} Leonard. Baltnerus apud Will.

[†] In the Lincolnshire fens, numbers are taken in semicircular baskets, open above and below. The manner is, by letting down the basket from the end of a fen-boat, and as soon as the fish is sound to have entered, it is immediately taken up into the boat.

The Gar *.

HIS fish is commonly known by the name of the feaneedle: it makes its appearance in fummer, a little before the mackrel, a fish it resembles in taste as well as in the spurious fins near the tail. It is a long slender fish. flattened a little towards the belly, and quadrangular as it approaches the tail. The head is flat, projecting forward into a very long sharp snout, which is characteristic of this species. The fides and belly are of a bright filvery colour; the back is green, marked along the middle with a dark purple line; the fides are also each distinguished by a line running from the gills to the tail. The lower jaw projects a great way beyond the upper, and terminates in a certain foft fubstance: on both fides it is provided with teeth, like the common pike; and the upper jaw is moveable as in the crocodile; a circumstance from which the gar is distinguished from all other fishes, as well as by the superior richness and vivacity of its colours while in the water.

To this genus is commonly referred the faury, a fifth not enumerated by Linnaus, but described by Rondeletius as an inhabitant of the Mediterranean. Above and below the tail there are a number of spurious sins, resembling

^{*} Acus vulgaris, Aldrovand. Esox Belone, Lin. Syst.

[†] Pisces, 232.

bling those of the mackrel. The jaws are produced like those of the garr, but are of equal length, and terminate in a very sharp point. The upper mandible is slightly incurvated, and the length of each is one inch. The eyes are large; the body anguilliform, but towards the tail it grows suddenly small, and tapers off into a very slender point*. Some years ago, a shoal of these sishes appeared in the Frith of Forth, and great numbers were cast ashore upon the lands at Leith, where the inhabitants called them Egyptian herrings.

Pritish Zoology, genus 32.

Voi. III.

Ii I

GENUS XLIX .- Elops.

THE faurus is the only species belonging to this genus; it is found in Carolina and in the West Indian seas; and in its external form nearly resembles the pike. The mouth has the same extraordinary gape; while the upper, as well as the lower jaw, is armed with teeth. The ventral and dorsal fins are somewhat larger than those of the common pike, and also placed nearer the head. The anal fin is hard by the tail, which is bifurcated.

The head of the faurus is large, and flattened horizontally; the body is flender, retembling a lance, and covered with large angular feales. The eyes are large, orbicular, and placed at a distance from each other; they are half covered with a nichating membrane. What particularly diffinguishes it, is a bony feale at the origin of the caudal fin, both above and below †.

^{*} Elops Saurus, Lin. Syft. Saurus, Slean. Jam,

[†] Guan, Hift, piscium,

GENUS L .- The Argentine *.

Or this genus there are two species; the one peculiar to the American seas, termed by Linnæus the Argentina Carolina; and the other found in the European and sometimes in the British seas. The European argentine is most frequently seen in the Mediterranean: Willoughby, who examined one at Rome, says, that the outside of the air bladder confils of a soliaceous silvery skin, which was made use of in the manufasture of artificial pearlt.

The fnout of this fmall fish is sharp and prominent; the body round and tapering, seldom exceeding two inches in length. The back and sides, as far as the lateral lines, are a pale ash colour, mixed with green: The belly, and parts below these lines, are exactly coloured as if covered with silver leas. The mouth is small, without teeth, excepting some asperities resembling them upon the roof of the mouth, and the tip of the tongue. The eyes are large; the irides silvery; and on the forehead is a patch of dark purple. The dorsal sin is placed near the equilibrium of the fish, and supported with ten soft rays. There are two pectoral, two ventral, and an anal sin, near to the tail, which is bifurcated.

^{*} Argentina Sphyræna, Lin. Syft. Pifciculus Romæ, Argentina dictus, Will. † Willough. 229.

GENUS LI .- 7 be Atherine *.


This tribe Linnaus has divided into two species; both distinguished by the slatness of the upper jaw, by a filvery stripe along the sides, and by having six branchiossegous rays. The atherine best known on our coasts, is very common near Southampton, where it is called a smelt. It is in season from March to the beginning of June, at which period it begins to spawn. It never deserts the place; but may be taken there at every season, excepting in hard frost.

The atherine is from three to four inches in length; and the body is femi-transparent, except where the spine, or the viscera intercept the rays of light. The shape of the body is oblong, and slender; the back straight; and the belly a little prominent. The belly is silvery; the back variegated with blackish coloured spots, which become more discernable after the scales are stripped off. The mouth is large; and its aperture, when the jaws are shut, seems to look upwards. The lips hang down: both jaws, as well as the palate, are beset with small teeth; and upon the back are two dorsal sins.

^{*} Atherina Hespetus, Lin. Syst. Pisciculus Anguella Venetiis dictus, Will.



Plate XXIX.



GENUS LII .- The Mullet *.

Rondeletius has described five different kinds of seamulets, which do not seem to be any thing more than varieties of the same species: Linneus enumerates only two; the sea-mullet, and the white. The former is taken both off the Luglish and Scotch coasts. It is covered all over with large scales; the teeth are all placed upon the tongue and palate. The back, which is of a dirty green, is slightly incurvated; while the belly projects more considerably, and is of a pure white. The upper part of the siss curiously variegated with white and black lines, proceeding in a parallel direction, from the head to the tail: The scales are also arranged in rows parallel to each other; and in the same direction. There are two dorsal sins; the sirst supported by spinous, and the second by soft rays; the tail is bisurcated.

The residence of the mullet is the sandy shoals and bays, into which there is an inlet of fresh water: In such situations, they are sound all round the shores of Great Britain, where they leave the sand all marked with round holes, the traces of their digging, for they keep constantly rooting like hogs among the mud. When surrounded by a net, the whole shoal frequently escapes by leaping over it; one attempting this mode of escape, and succeeding, is immediately followed by all the rest ‡.

The

Mugil Cephalus, Lin. Syst. Mullet, Willough.

[†] Willough. p. 274. † Oppiani. Halieut.

The mullets are in great plenty in the Mediterranean, and along the fouthern coasts of France. In a certain lake near Martegues, in the fouth of France, there are vast shoals which enter there during spring, the season of copulation among these sishes; after impregnation they return to the sea, but are intercepted by weres of reeds.

Of the milts and roes of the mullet, the famous Botargo of the Italians is made up. The materials are taken out entire, and for a few hours covered up with falt; afterwards they are preffed between two boards, dried in the fun for about a fortnight, when they are fit for use. This composition is said to brace a weak stomach, and to give an exquisite relish to wine, when ate before drinking.

The food of the mullet is mud, or fea-weed; it never attempts to devour any fish. The slesh of this animal is palatable, though at present not a fashionable dish; its slavour greatly depends on the ground where it is fed; if among mud, it constantly tastes and smells of that kind of food.

According to Gefner*, the mullets are the most salacious of all animals; the male pursuing the semale even after she is captured, till drawn ashore. At Tarentum, a male, pursued his semale that had been caught with the book and line, till he received several strokes with a spear, and was at last taken and drawn into the boat: By such an ungovernable desire are these animals impelled to sulfil the purpose of nature.

At Athens, and afterwards at Rome, when an unfortunate galant was catched in the company of his miftrefs, the mullet is faid to have been employed by the enraged hufband husband as an instrument of a shocking punishment *. Juvenal and Horace both allude to it in mentioning the various modes of revenge which rage distated to the injured spouse †.

----Necat hic ferro, fecat ille cruentis Verberibus; quosdam mæchos et mugilis intrats

^{*} Vide Caufabon, in Athen.

[†] Juven. Sat. x. 316. Hor. Sat. ii, lib. i. 132,

SECTION XII.

GENUS LIII .- The Flying-Fift *.

This genus comprehends two kinds of flying-fifth, the volitans and evolans of Linneus; the former is the most common kind of the winged inhabitants of the fea-Hardly any inflance occurs of either kind being found in the British feas +: They are frequent in the Mediterranean, and warmer latitudes in the Atlantic, where they are continually hunted by the Dorados while they continue in their own element; and, as often as they endeavour to escape them by ascending to the air, they become the victims of the gull or the albatrofs; or are by them again forced back into the mouth of their purfuers below, who have still their eyes fixed upon them, and keep pace with their flight. Thus harraffed by inveterate enemies in both elements, they often fall down upon the decks of thips as they pais, and feem to yield themselves up to man as the more merciful destroyer.

In fize, the flying fifth nearly refembles the mackrel, only of a longer and more flender form: if the wings are taken away, it bears a flrong fimilarity to the mullet:

The

^{*} Exocætus Volitans, Lin. Syst. Mugil Alatus, Rond.

[†] Vide Pennant's Brit. Zool. gen. 38.

The icaies are of the fame large fize; and the fame dark green covers the back: the head is depressed; the snout sharp; the mouth small, and destitute of teeth. There is only one dorsal sin, about a fourth of the sish's length from the tail. The two pectoral fins, which are the instruments of slight, are almost as long as the body, and supported by sourteen strong branchy and jointed rays.*.

The ancients were acquainted with the flying-fish, and gave it the name of fea-fwallow +; and all the different writers have taken notice of its faculty of mounting into the air by means of wings.

* Willough.

† Hirundo Plinii; l. is.

Vol. III.

Mm

GENUS LIV .- Polynemus.

Ims tribe contains three species, all of which are race exotics, which never visit our seas. Their bodies are oblong, thick, and covered with scales; as is also the head, which slopes off into an obtuse snout. The jaws are of equal length when open; but when shut, the lower is longest; both are surnished with small, thick and immoveable teeth. The tongue is narrow and smooth; but the palate is exasperated with a number of sharp tubercles. The eyes are of a middle size, placed at a great distance from each other, and covered with the common tegument of the head; before them appear the two large apertures of the nose.

The fides are compressed, and the abdomen carinated 3 each of the lateral lines are smooth, and bend in a direction parallel to the back. There are two dorsal fins; the first spinous, and placed about the fish's equilibrium; and the second, which is larger, behind it. The pectoral fins are long and narrow; the ventral are placed in the abdomen; the tail is large, broad, and bifurcated.

GENUS LV .- Mormynis *.

The fishes of this genus are only two, and both strangers to our shores: Their generic characters, an oval body compressed at the sides, and narrow towards the tail: The snout protuberant; and the opening of the mouth very small and rough, with a few setaceous teeth placed on the extremity of the jaws. The back has a large swell, and from it rises one dorsal sin; the pectoral sins are small, and situated near the belly; the anal sin is behind the equilibrium, and that of the tail bisurcated.

M m 2

^{*} Le Mormyre, Guan. Mormyrus, Lin. Syst.

f Guani Hist, pisc. gen. 50.

GENUS LVI .- The Herring *.

Many of the fishes of this genus are well known. Some of them, such as the herring and pilchard, constitute a larger portion of the food of the human race, than perhaps any fish whatever. The herring sishery, which forms to considerable a branch of commerce to the English, Dutch, and other nations of the north of Europe, is a modern invention. Those myriads of fish which annually teem from the Artic Seas, and which are the objects of this sishery, are supposed to have been altogether unknown to the ancients; for there is no word which seems to have been appropriated to this most numerous species.

The external characters of the herring are fo well known, that all description of them would be supersuous. The grand winter residence of this sish is within the artic circle, where insect food abounds in a still greater degree than in warm climes. The particular food, however, of these sithes has not yet been perfectly ascertained: The intestines are filled commonly with a blackish mud, apparently too much digested to discover exactly its nature; it is supposed to be a crustaceous insect;

From the Artic Seas, the herrings annually migrate along the shores of America, as far as Carolina; along those

^{*} Clupea Harengus, Lin. Syft. Herring, Will.

¹ Onifeus Marinus, Lin. † Catefby Carolina, II. 33.

those of Europe, as far as the north of France; and on the east of Asia, they are found on the shores of Kams-chatka. The great army that annually issues from the north, separates into several divisions: The first makes its appearance off the Shetland Isles in the months of April and May; but these are only the harbingers of a far more numerous body, that follows in June. The appearance of these shoals is always announced by the gulls, gannets, and other rapacious birds, that continually hover above them; but when the great body approaches, about the beginning of Harvest, its breadth and depth alter the appearance of the ocean, which sparkles with various colours, like a bed of precious stones, by reslecting the rays of the sun from the scales and sins.

When the herrings first quit the regions of the Frozen Sea, they are divided into different columns, of five or fix miles in length, by three or four in breadth; and in their progress fouthward, the first obstacle in their way is the Shetland Islands, by which they are separated into two grand divisions; the one advancing along the whole British coast, filling every bay and creek, till it reaches the Channel, after which it gradually thins, till it disappears. The other great wing makes a similar circuit round the west coast, till it reaches the north of Ireland, where it is again subdivided; part entering the Irish Sea, and part scattering along the west shores of Ireland, till it disappears about the entrance of St. George's Channel.

Several stationary sisseries are established on the west of Scotland and Ireland; but the herrings are by no means uniform in resorting to the same loch or bay annually. They frequent a certain space for a number of years, and then capriciously desert it for perhaps as many more. On the coast of Wales, and among the Hebrides,

they have at different times occupied and deferted their feveral stations, without any apparent cause. But although this stupenduous gift of the munificent Author of Nature, is at times partially distributed, it is never totally withdrawn. The same instinct invariably operates; and if one part is deprived of its effects, another teems with plenty, and relieves the necessity of such as are less liberally supplied: Thus, thousands of the poor are annually supported, and rendered happy, by that instinct which the Almighty hath originally impressed upon this useful part of his creatures *.

But this appetite for migration, which brings the herzing annually to our shores, serves also purposes of the greatest importance in the economy of these animals: It leads them to the shallow and tepid waters of the temperate zone, to deposit their spawn, where it is matured and vivished with greater certainty than in the midst of the frozen ocean.

The multiplication of their kind, is, therefore, perhaps the primary purpose of nature in the migration of the herring. It is not from a defect of food that they abandon regularly their northern retreats; for it is immediately after leaving these, that they are most plump: before their return, they are greatly reduced by spawning, and are then thin and miserable. The time of spawning is from the beginning of winter till January, when they almost totally disappear from our coasts, or are taken in simall straggling parties by the sishermen, for the purpose of bait. It is probable, that at the time they abandon our feas, they again repair to the north, to restore their yigour,

^{*} Brit. Zoology, Gen. 39.

vigour, and, by impregnation, to replace that immense waste, which the exertions of man, and the rapacity of fishes, has made of their species.

The spawn, after being discharged by the parent sishes, continues to float on the waves for a confiderable part of the Spring. In the beginning of Summer, the young enter upon life, and in June and July are feen in vaft numbers approaching the shores. They are then about two inches in length, and most probably feed upon infects, as thousands have been caught a little after that period with the common trout fly. Along the Yorkshire coast, these young fish are called file; upon that of Scotland, they obtain the general name of fry, which feems to be applied indifcriminately to the young of all fish. During winter, it is probable that the young herring retire to the north, along with the old. On their appearance next fummer, although the shoals are made up of fish of very different fizes, it does not appear that they are then distinguishable from the fish of greater age.

The Dutch, who have fet the example to the other nations in almost every thing relating to commercial industry, first commenced the herring fishery in 1164*; and continued in the exclusive possession of it for several centuries. At length the English, roused by their gains, and jealous of that naval power of which it was the grand source, endeavoured to participate in this lucrative commerce, and for that purpose sought many well disputed battles. The superior industry and experience of the Dutch, always have, and probably long will, secure them the greatest share of this trade. The most considerable stations of the British sisteries are off the Sketland and

Western

Western Isles, and off the coast of Norfolk; in all these stations the Dutch ships are ever ready to take their share.

One William Benkelen of Biervlet, near Sluys, was the inventor of the useful expedient of pickling herring; and it is from him the operation has its name, both in the English and German languages. This great benefactor of mankind died in 1397; and his memory was held in such veneration by the Emperor Charles V. that he paid a visit to his tomb. The Dutch, it is said, are remarkably fond of the herring in its pickled state: A premium is given by the state to the first bush that arrives loaded with a cargo; which fells them for a very high price. The whole inhabitants welcome the arrival with shouts of exultation; and the same joy seems painted on every face, which the Egyptians display on the first overslowing of the Nile*.

The Pilchard t.

The pilehard, in some respects, resembles the herring; it is remarkable for similar magnations, and stated returns from the north. As this fish is far from being so general as the herring, appearing only on the coast of Garawall for a short time in the end of Summer, we shall give a short description of its external figure.

Its

Oil

Its ordinary fize does not exceed nine inches; the body is a little thicker in proportion than that of the herring, and the belly lefs fharp: The scales are very large, thin, and rounded; the back is bluth; the sides and belly silvery; the head is compressed, and the mouth without teeth. The upper angle of each of the gitls is marked with a large black spot; some have sour or sive such spots disposed in a row near the tail; the lower edge of the belly is ferrated, owing to the row of large clypiform scales that runs along it.

The pilchard annually appears in large shoals off the Cornist coasts, from whence it is regularly exported in great quantities into the Mediterranean. The fishing season is from June till the beginning of Winter; a few sometimes are found hovering about till Christmas. There are spies, or huere, as they are provincially called, stationed on the cliffs that project into the sea, whose business it is to watch the progress of the shoals, and to give notice to the boats below of the measures they are to take, by means of signals previously agreed upon.

By an act of James I. these liners are empowered to go over the grounds of others, without being liable to damages *. By their motions the nets are both shooted and drawn; and when they have been successful in inclosing part of a shoal, an hundred thousand are sometimes taken at a single draught. In October 1767, there were at one time inclosed in the Ray of St. Iros, seven thousand hogsbeads, or two hundred and forty-sive millions of sish. The average amount of the export of these sists to be 29,795 hids, annually; which, including the bounty paid by government, and the price of the

Nn

Vol. III.

oil extracted from them, are fold for about forty-nine thousand pounds.

But the great benefit arising from the pilchard sidnery, is the employment it assords to a number of seamen, which are thus trained up for the desence of the nation. A variety of hands, too, are employed on shore, in the disterent operations of salting, pressing, washing, and cleaning the sish, as well as the trades people, who depend upon the construction and sale of boats, nets, ropes, and casks *.

The Anchovy to

This species is taken in the Mediterranean, and export, ed, pickled, to the different nations of Europe. Before falting, the head and viscera are taken away; and in this state they are ate raw, with vinegar and oil. They are supposed to give wine an excellent relish; but probably their chief merit consists in bracing the stomach, after being relaxed by excessive drinking.

Near a century ago, the anchovy was found at the mouth of the river $Dee \ddagger$; and fince that time has hardly ever been deemed an inhabitant of our feas. The length of this species is from four to fix inches; the body slender, but thicker in proportion than that of the herring: The

eyes

^{*} Borelase's Cornwall.

⁴ Clupea encraficolus, Lin. Syft. Anchovy, Willough.

Ray's Letters, 47.

eyescare large; the irides white, with a cast of yellow; the under jaw much shorter than the upper; the teeth small; a row in each jaw, and another on the middle of the tongue. The scales are large and deciduous; the back green, and semipellucid; the sides and belly solvery and opaque; the edge of the belly smooth, and the tail forked.

The genus of herrings, according to Linneus, contains eleven species; an ong these, however, the pilchard is not enumerated; although, from its history, we have seen that it is one of the most common, and perhaps the most numerous, fish upon our coasts.

The fprat is also a very common species belonging to this genus, which Willoughby and Ray have supposed to be the fry of the herring. It nearly resembles the anchovy in fize and shape; but from the time of its appearance on our coasts, it seems to be a distinct race from the herring. These fish come into the river Thames in the beginning of November, and continue there tilt March; a season when the herring have long retired into the North Sea. During the whole of the Winter season, they constitute a large portion of the food of the citizens of London. At Gravesend and Narmouth, they are cured like red herring, and sometimes pickled like anchovies, from which they differ but little in their slavour.

The Shad *.

Thus is supposed to be the thriffa of Ariffetle and Oppian, and the alaufa of Arfonius; but the descriptions which the ancients gave of their species were so vague and general, that we can feldom aftertain the enact species they had in view. The flud, in the flape of the body and head, bears a firong refemblance to the herring; the former, however, is a deeper fish, and more compressed laterally; it is also confiderably larger, weighing from four to eight pounds. In colour, this species resembles the pilchard; the back being of a blackish blue, and the fides filvery. On each fide, near the gills, there is a Targe black fpot; and behind it, in the same straight line, there is a row of five or fix others, fmaller in fize. The lateral lines are obscurely discerned; the scales are large and decidnous, their margins punctuated with finall black dots; the belly is carinated, and firengly ferrated by the edges of forty scales, which proceed in a row, from the guils to the anus. The eyes are large, covered as far as the irides with a loofe membrane; the upper jaw is fomewhat shorter than the lower, and exasperated on the margin with very minute teeth; the other parts of the mouth are fmooth. The dorfal fin is fituated near the centre of the fish, supported by eighteen cartilaginous rays, of which the middle are longest; the ventral, pectoral.

^{*} Clupea Alofa, Lin. Syft. The Shad, or mother of herring, Will.

toral, and anal fins are fmall in proportion to the fish's fize; and the last is placed very near the tail, which is greatly forked *.

The Severn produces the shads in higher perfection than any other British river. They appear there in April or May, according to the warmth of the Spring; and after two months they disappear, and are succeeded by other varieties of the same sish. About Gloncester, the shad is esteemed a great delicacy; it is there caught in nets, and sold at a price as high as that of salmon It is from thence they are sent to the London market, where they are distinguished from the shad of the Thames, by the French name alose.

It is not ascertained where the shads spawn: at the time of their mounting the river, they are in sull roe; but none are caught on their return to the sea, after shedding their spawn. The bleak, which are caught in the Severn in the months of July and August, are erroneously supposed to be the young of this species. The shad is found at certain seasons in the river Nile; it is there, as well as in England, a sish of passage; it is also seen in the Mediterranean near Smyrna, and on the coast of Egypt, in the neighbourhood of Rosetto.

This species was reckoned very mean food by the ancients; Ansonius characterises it as the food of the poor §. Those of the Thames agree to his description, being a very insipid and coarse sish. Jovius gives a similar account of this animal as an article of food; but afferts that it improves after ascending the Tiber ||. In the

^{*} Willough. page 227. † Belon, 307. ‡ Belonii Itiner, 98.

[§] Stridentesque socis obsonia plebis Alausas; Mosella, 138.

De Pifcib, Roman,

the Severn there is a variety caught near Glocefler, called the twaite, which is equally differenced with the Thames fhad. These varieties are distinguished by their fize, the twaite being nearly one half smaller than the true shad; and has always fewer black spots behind the gills *.

^{*} British Zoology, Gen. 39:

SECTION XIII.

GENUS LVII .- The Carp.

LINNEUS makes this genus confift of no less than thirty-one different species; analogous indeed in their external characters, but differing considerably in their habits and economy. They are all of an oblong and oval shape; the body laterally compressed, and covered with large imbricated and deciduous scales. The mouth is small, and destitute of teeth; hence they have been denominated leather mouthed sishes. They are all distinguished by three branchiostegous rays, and one dorsal fin rising from the middle of the back, which is convex, and somewhat carinated.

The nostrils are covered with valves, by which they are alternately opened and shut; and the greater part of the genus have cirri, or beards, round the mouth. Some of these sish were brought to England from the south of Europe, and are now naturalised in our fresh-water ponds.

The Garp *.

THIS species was introduced, according to Fuller, into England in the year 1514, by Leonard Mascal+; but before that time, we read a description of it in the book of St. Albans, printed in 1496. It is by means of transportation that they have been naturalised in most countries of Europe, as is the case in Sweden, England, and France. They have not yet found their way into Russia, though Polish Prussia, which is probably their native country, is at no great distance: There they abound in all the rivers and lakes, and are taken of such vast size, that they have become a valuable article of commerce. The merchants purchase them from the lakes and rivers belonging to the Polish Noblesse, and transport them in well-boats to Sweden and Russia. In imitation of them, Some of the English gentry have begun to turn their fishponds to account, by regularly felling their carp t.

The carp is, perhaps, the most remarkable among the spinous sistes for longevity: An instance is recorded of one in the palatinate of Germany, that lived an hundred years in the fosse surrounding a fort ||. Their size in some countries is said to correspond to this extraordinary longevity; several being taken in the Lacus Larius, of two hundred


^{*} Cyprinus Carpio, Lin. Syst. La Carpe, Belon.

F British Worthies, Suffex, 113. . . Brit. Zool. gen. 40.

A Gesner de piscib. 312.

Plate XXX







CHUB



hundred pounds weight; while those of the Dneister sometimes arrive at the enormous length of five feet.

Of all the finny tribe, the carp is most tenacious of life: when placed in a net wrapped in wet moss, and hung in damp cellars, it will subsist for a fortnight †. This fact is so well attested by experiment, that a method of fattening this sish has been adopted, which is founded upon it. The net is from time to time disped in water, and the sish crammed with wheat-bread soaked in milk; and by this management, it becomes in a short time nor only much fatter, but of a far superior slavour to those sed in the pond ‡.

The fishes of this species being in some measure domesticated, their manners and economy have been more narrowly examined than most of the other tenants of the water; and the ascertaining their secundity has been the result of these observations: The carp is prodigiously prolific; its belly being almost at all seasons distended with a roe, which sometimes increases before parturition, to the full size of the fish who carries it; and when weighed opposite to it, has often been known to preponderate. Upwards of two hundred thousand ova have been numbered in one roe ||; and, if we may credit Arijtotle, this immense off pring is produced sive or six times every year s. Such is the kind attention bestowed by nature in preserving this useful class of animals, for the support of other portions of her innumerable family!

The carp, like most other species, frequent the shallow water at the season of spawning; when a dozen of males Vol. III.

^{*} Jovius de pife. Rom. 131. † Hift. Nat. Pol. 142. / † Derham's Physo. Theol. Ed. 9. p. 7. || Phil. Transact. 1767.

⁵ Hift, Animal

are feen pursuing one female as she emits her ova; and upon these they eject their smelts: by which operation naturalist suppose that impregnation is effected. In this uncertain process, however, many of her ova must be distipated in the water, without coming into contact with the sperm of the male; and hence that sertility is abridged, which would otherwise soon overstock the whole waters upon the globe.

Among all domiflicated animals, the purity of their manners is vitiated; and mixed races are produced by an unnatural connection between different species: This is examplifyed by the horse and the ass, by the different species of dogs, and by many of our domestic poultry; the carp and tench, in their consinement in ponds, are also said to mingle their spawn, and to transmit an ambiguous race to posserity: In similar circumstances, the carp and bream, are reported to "overstep the modesty of Nature," and to violate those laws which she has so generally established between kindred tribes.

The flesh of this species is rather insipid and soft, but varies greatly, according to the place where the fish is reared, and the nourishment with which it is fed: Those in stagnated water are seldom sound so well stavoured, as they that are taken from a pure and rapid stream. Of the roe of this sish a caviare is made for the Jews, who abominate this, and every kind of the sturgeon.

The carp is extremely shy in taking a bait, and, for this reason, the more common method of taking it in rivers is with the net. Its cunning is displayed by the various methods it practises to elude this instrument of destruction: Sometimes it leaps over the net; at others, immerses itself so deeply in the mud, that it is drawn over it: At the spawning season, all this artistice seems to forsake the animal, and it becomes then so simple, that it suffers itself to be tickled, handled, and caught, by any person that will attempt it *.

This fish, when young, is of a dusky colour, growing yellow as it increases in years: The body is of a thick thape; the feales remarkably large; and when the fifth is in feason, of a fine gilded hue. The mouth is furrounded by fat, fleshy lips, of a yellow colour; at the angle of the mouth there is a fingle beard, of a yellow colour, and above it another, fhorter, black, and lefs observable. The carp has no teeth; their office is supplied by two fmooth triangular bones, above and below, which are placed at the entrance of the throat, and by acting against each other, comminute the food before it passes downward. The dorsal fin extends far back, almost reaching the tail, which is bifurcated. Opposite to it is the anal fin, the third ray of which, as well as that of the dorsal, is strong, and armed with sharp teeth, that point downwards.

The Barbel +.

This species is longer and more tapering in its shape than the perch; being from a foot to eighteen inches in length: In the Nile, it sometimes weighs twenty pounds.

O o 2 The

British Zoology, gen. 40.

[†] Barbus, Rondel. Cyprinus Barbus, Lin. Syft,

The colour of the back is a pale olive; the belly is filvery, and proceeds without protuberance, nearly in a
firsight line; so that, when tenching the bottom, the mouth
at the same time reaches the ground. The head is sloping,
the snout somewhat pointed; and on the summit of the
rostrum there are two beards; other two spring from
the sides of the mouth; a circumstance from which this
sish derives its name almost in every language. From
the middle of the back, which is a little arched, there
rises one dorld sin with ten rays, of which the first is the
largest. The task is bifurcated, and between it and the
anus there vises another sin supported by seven rays.

The flem of the barber feems to have been held in no offication among the ancients: Aufonius alone mentions it, and, without any commendation, excepting that it improved by age +. It frequents the still and deep parts of the British rivers, where it lives in fociety, and roots like a hog among the fandy banks. The Danube is famous for the refort of this fish; it is there found in the caverns of the rocks, and holes under the banks, in numbers that defy all computation. The peafants take them with their hands, in quantities sussicient to load a waggon t. Those in our rivers are characterised by the same tameness and insipidity; they are often taken with the hand, by diving: In fummer they move about in quest of food; but ouring autumn and winter, they confine themselves to the deepest holes. The poorer peasants alone, and that from hard necessity, are constrained to eat them boiled with a piece of bacon. They are deemed the coarfest of all frella

^{*} Willough, p. 259.

† ——— Tibi contigit uni
Spirantûm ex numero non inlaudata fenectus.

† Albertus.

Welh water fish: their roes are even held poisonous; for they affect those who unwarily eat them, with a nausea, vomiting, and purging *.

The Tench +.

THE tench is unnoticed in the writings of all the aucients before Aufonius; and by him it is mentioned in tuch terms of difrespect, as shews the capriciousness of taste;

Quis non et virides vulgi folatia Tincas Norit ?

Willoughly mentions the infalubility of this fifth, on account of which, it was unanimously condemned by the medical people of his time; he denies, however, that it is ungrateful to the take; potterity have been of the fame opinion, and have gratified their palates with it without feeling any bad effects.

The refinence of the tench is either in stagnated or running water; although it is taken in the Tiber, a pretty rapid stream, it seems to have migrated there from the adjoining takes. There is faid to exist a wender-

ful

^{*} British Zoology, gen. 40.

[†] La Tanche Belon. Cyprinus pinna ani radiis 25. Lin. Syst.

[‡] It is now naturalised in England, where it has been imported from its native lakes in the fouth of Europe. We have already noticed with regret, that the ingenuity of man has made so little progress in the domestication of sishes; this and the carp being the only species of near five hundred, that have been brought into that state.

ful friendship between this animal and the pike *, a sish which, though it devours every other, will not offer violence to the tench, on account of its healing quality. It has hence been called the physician of the sish; the slime of its skin being of a healing nature, an inhabitant of the water is no sooner wounded, than the physician is at hand with a powerful styptic:

Close to his scales the kind physician glides, And sweats a healing balfam from his sides †.

The Jews practifing physic at Rome, are said to apply the half of a tench, after being cut up, to the soles of the scet, when their patients are seized with violent sceners; we believe, however, that the medical virtues of this sish, either while dead or alive, never existed but in the imaginations of the ignorant and credulous.

The tench spawns in spring, and in the beginning of summer; it is superior in sertility to the carp, near sour hundred thousand ova having been numbered in a single roe. The young fry are remarkably quick in their growth; a circumstance the more credible, because they devour mucus and every kind of silth with avidity. The most proper bait for them is small earth worms: according to the present sashion of eating, they are deemed wholesome and delicious food.

The tench is thick and short in proportion to its length; the scales are very small, and covered with slime: The colour of the back is dusky; the dorsal and ventral sins of the same colour: The head, sides, and belly, are of a greenish cast, most beautifully mixed with gold, which


is

^{*} Salvianus apud Will.

^{*} Diaper's pifcat. Eclogue, ii.



Plate XXXI.





WHITE BAIT







MINOW

is in its greatest splendor when the fish is in its highest season *.

The Gudgeon t.

The gudgeon, according to Willoughby, is preferred by the English to most of the river sish; it is taken in gentle streams, and is generally of a small size, measuring only about six inches: In the Kennet and Cole, they are of superior bulk to those of every other place, and sometimes weigh half a pound ‡. They are assembled by raking the bed of the river, which makes them crowd in shoals to the spot, expecting food from this disturbance.

This species is mentioned by Aristotle ||, and is therefore probably an inhabitant of the rivers of Greece: We know from the accounts of travellers, that it is found in Syrias; and thus exhibits a remarkable instance of the capacity of fishes to accommodate themselves to different climates. River water must always correspond with the temperature of the country through which it slows; while that of the sea is of a much more equal heat in different latitudes; and hence we find sea fish travelling into a greater variety of latitudes than those of fresh water.

There

British Zoology, gen. 40.

[†] Gobio Fluviatilis, Rond. Cyprinis pinna ani radiis 2. Lin. Syst:

[‡] Brit. Zoology, ubi fupra. | Lib. viii. cap. 19.

[§] Dr. Ruffel, p. 75.

There are two species of the gudgeon, a larger and a fmaller; the former is an inhabitant of our rivers, the latter of those of Germany, where it is called wapper. The British gudgeon upon the back is dusky; the belly of a dirty white: It is distinguished by nine or ten large spots of a blackish cotour, upon the lateral lines; others of a smaller fize, being irregularly scattered over the back and fins.

At the angle on each fide of the mouth, there is a finall beard of a quarter of an inch; near the eyes are two large apertures of the note: Neither jaw is furnished with teeth; but at the entrance of the throat there are two triangular bones, that perform the office of grinders, refembling those of the carp. There is a fingle dorsal fin, with nine rays; the pectoral have each sourteen, and the yentral eight or nine; the tail is bifurcated.

The Bream *.

THE bream is one of the fresh water sish, in no high reputation; its slich is fort and insight. It is of a rhomboida! shape, and in the lakes of Italy is found sometimes three feet in length, by two in depth. The back is greatly arched, and the sides strongly compressed: The lateral lines which proceed from the gills, are bent downwards,

Cyprinus Brama, Lin. Syft. La Bremme, Belon.

tive

curvated like a bow, and nearer to the belly than the back of the fish: above, the bream is a blackish green, the fides and belly white, with a gilded line when the fish is in high season *.

Stagnated water, or rivers gently gliding along, are the refidence of this species; which there feeds upon herbs, mud, and clay. Its feafon of spawning is in the month of May; at which period the males are marked upon the head with a number of white tubercles; an incident which Pliny takes notice of as befalling the fishes of the Lago di Como in Italy +.

Similar to the bream is the rud, a fish frequent in the Rhine, the fens of Holdernesse, and in the Charwell, near Oxford. It is about two pounds weight; and in feafon, almost the whole year round, except in the month of April, when it fpawns. At this feason, the head of the male is rough with white tubercles, refembling those of the preceding species. The rud is reckoned a superior dish to any of this genus.

The rud feems to be an intermediate species between the carp and the bream; it is broader than the former, and deeper than the latter fish. The colour is brown, changing into yellow; the fins are of a reddish hue, and the whole body is covered with very large scales. The opercula of the gills are, for the most part, marked with a blood-coloured fpot.

In many of the fish ponds about London and in the fouth of England, there is reared a fish called the crucian &; but as it is not mentioned by Willoughby, it is probably not a na-

VOL. III.

Pp

^{*} Willough, p. 248. † Lib. ix. cap. 18.

[§] Cyprinus Caraffius, Lin. Syft.

tive of our rivers. The colour of this species is a deep yellow; as an article of food, it is held in little estimation. It is very deep and thick, the back much arched, and the dorsal sin supported by nineteen rays; the sirst two strong and serrated.

The Roach *.

The roach, like the other fish of this genus, frequents the still or stagnated waters; it is an excellent pond fish, not only thriving well in artificial sheets of water, but replenishing them sooner than any other species, with a numerous race of inhabitants. Its offspring are not so numerous as those of the carp or tench; this extraordinary fertility must therefore be owing to a smaller number of the young perishing before they arrive at their adult state.

The roach is fmaller than the bream; the shape of the body, like that fish, is deep and laterally compressed, being four inches broad by twelve long. The back is dusky, with a shade of green; the belly pale; the ventral and anal fins are a cinnaber colour, the dorsal and tail sin having less of the red †. The scales are of a silver hue, broad and deciduous. The lateral lines are curvated, and run parallel to the swell of the belly: The tail is bifurcated.

The

^{*} Cyprinus Rutilus, Lin. Syst. Leuciscus, Rondel:

[†] Willough. p. 262.

The dace * haunts the fame places as the roach; like it, too, it is a great breeder, very lively, and during fummer, fond of gamboling on the furface of the water. Its colour is a mixture of brown and yellow; the feales are fmaller than those of the roach; the length of this fish is feldom above ten inches.

The graining is a fish fo nearly resembling the dace, that perhaps it may be considered merely as a variety: The usual fize is from six to eight inches; it is caught in the Mersey near Warrington.

The Gold-fift +.

This species was first introduced into England from China, in 1691; and that specimen having been destroyed, they were again imported about thirty years ago in a considerable quantity. From that time they have multiplied greatly in the neighbourhood of London, and from thence have been gradually dispersed over the rest of the kingdom, where they now breed as freely in the open water as the carp.

The most beautiful species of these sithes are kept in China for the amusement of people of rank. They are confined in small porcelain vessels, and placed in the courts to decorate the entrances of the houses belonging

P Cyprinus leucifcus, Lin. Syft.

[†] Cyprinus Auratus, Lin. Syft. Kin-yu, Du Halde Hift. China,

to the nobility. The beauty of their colours and lively motions, give great entertainment, especially to the ladies, whose pleasures are extremely circumscribed, from the cruel policy of that people *.

In shape, these sishes bear a great resemblance to the carp; and in their native lakes they are perhaps not greatly inferior in fize; for they are faid to equal our largest herring +. Beautiful coloured drawings of them have been executed by Edwards I; their prevailing colour is gold, of a molt amazing fplendour; there are varieties marked with fine blue, brown, and bright filver; the most beautiful are faid to be taken in a finall lake in the province of Che-Kyang |.

The Chub T.

THERE are five species, or rather varieties, of this fish enumerated by Willoughly, after the ancient naturalists : That fish known in the English rivers does not grow to a large fize, feldom exceeding five pounds. The chub is a very coarse fish, and full of bones; it frequents the deep heles of rivers, and, during fummer, commonly lies on the furface, beneath the shade of a tree or bush. It is a very timid fish, finking to the bottom on the smallest clarm, even at the pailing of a fhadow, but foon refumes its

^{*} Brit. Zoology, gen. 40.

[‡] P. 209.

[†] Vide Du Halde, p. 316. Brit. Zool. ubi fupra.

T Cyprinus Cephalus, Lin. Syft. Cephalus Fluviatilis, Rondel.

its fituation. It rifes to a common trout fly, and feeds on worms, catterpillars, grafic ppers, and other coleopterous infects, that happen to fall into the water *.

The body is longer than that of the carp; the head flattened; the back of a dusky green; the sides and belly silvery, with a shade of a gold colour, if the fish be old and in high season. The scales are remarkably large and angular, like those of the carp; the mouth is of a moderate size, round when the jaws are opened, and wholly destitute of teeth. The number and disposition of the fins resemble those of the other sish belonging to this genus; the tail is forked.

The Bleak +.

This species is frequent in the English streams, as well as those of the Continent. According to Ausonius, the taking of these sites was anciently the amusement of children, as that of the following species is at present. The sless is soft, and destitute of fat, making no very commendable morsel.

They are at fome feafons apparently feized with a diforder, which feems to occasion them great agony: They are then feen tumbling about near the furface of the water, and are incapable of fwimming far from the place;

but

^{*} Brit. Zool. Species 175.

[†] Cyprinus Alburnus, Lin. Syst. Alburnus, Rondel.

Alburnos prædam puerilibus hamis.

but in about two hours they recover and disappear. When thus affected, the sishermen on the Thames call them mad bleaks. Pennant imagines, that they are then troubled with a species of hair worm, which Aristotle observed to infest some other kinds of sish *.

It is of the fitvery scales of this species of fish, that artificial pearls are made; an art which ower its origin to the French, and is by them carried on to such an extent, that one artist in Paris used thirty hampers of fish in this manufacture, during the space of a single winter. The scales are beat down into a sine powder, then diluted with water, and introduced into a thin glass bubble, which is afterwards filled with wax.

The minnow is a beautiful fmall fish, about three inches in length, and fearcely half the fize of the bleak; it frequents most of our pure gravelly streams, and is always gregarious. It has no teeth; and the scales are so small, that they are almost invisible. The back is a dark olive colour; each fide beautifully adorned with a lateral line of bright gold. The colour of the sides and belly varies; in some it is white, and in others yellow; in some it is of a rich crimson. Taking the minnow is one of the favourite, and perhaps most innocent amusements of children. This first essay at angling is performed with a bended pin, baited with a small earth worm.

During the month of June, there appears in the Thames, near Blackwall and Greenwich, a fish evidently of this genus, called the white bait. No naturalist has yet determined to what particular fish it belongs, though all are agreed, that it is the young of some species that resorts there. Some have ascribed its origin to the shad; others to the sprat, the smelt, and the bleak. These sish,

however,

^{*} Brit. Zool. Species 176.

however, are all found in other streams, while the white bait is peculiar to the river Thames.

The white bait has a greater similarity to the bleak than any other sish; the number and disposition of the sins are the same; and the body is compressed on the sides, in the same manner. The usual length of this small species is two inches; the dorsal sin is placed before the equilibrium; so that, when suspended by it, the tail sinks down. The tail is forked, and black on the tips. These small sry are usually caught for the purpose of bait for other sish; when fried with slour, they are reckened a delicious viand by the lower order of epicures, who frequent the taverns contiguous to the places where they are taken *.

^{*} Brit. Zool. Species 176.



NATURAL HISTORY

OF

INSECTS.

CHAPTER I. OF INSECTS IN GENERAL.

Sect. I.—General idea of the number and variety of Infects: Inducements to the study of Entymology. History of this Science.

We are now to enter upon that department of Natural History which treats of Insects; by the study of which, we are conducted into a province the most extensive, and by far the most populous, of the whole empire of nature. The residence of quadrupeds, as we have seen, is confined to the land; that of sishes to the water; while birds are enabled to rise from the surface of these elements into the aerial regions. Nature, however, has assigned a still more extensive range to those animals upon whose history we are now to enter. They are seened to pervade every part of her dominions, in num-Vol. III.

bers that defy all computation: For in nothing does the immensity of her works more strikingly appear, than in the infinite number and variety of those her smaller productions.

Entymology, therefore, of all the sciences, presents the widest, field for investigation. The number of experiments and observations necessary to furnish a complete history of fo many fmall animals, is not only great; but the difficulty of making them is also immense. The number of ideas with which a skilful botanist must load his memory, before he can acquire an accurate knowledge of above two thousand plants, is no doubt great: his talk, however, bears no proportion to that of the entymologist; for, amidst so many plants, there is perhaps hardly one that does not furnish nourishment and an habitation to feveral infects; while many, fuch as the oak, afford a retreat for fome hundreds of different species *. These plants, however, are far from being the only abode of infects; vast numbers reside upon the larger animals, whom they continually fuck; many live upon and deyour others of their own order. Infinite numbers spend a part of their lives in the water; others remain there entirely: The earth fwarms with multitudes; and the air teems with others, too fmall for the human eye to observe, and too numerous for the imagination to conceive!

While the numbers of this class of the animal kingdom thus exceed all our powers of conception, its varieties are also multiplied to a degree that renders a complete discrimination of them equally impossible. The disserent species

^{*} Mem. pour servir a l'Hist. des Insects, par Reamure, Tome primage 2.

species of insects are not only numerous, but each has its own distinct history, and exhibits manners, appetites, and modes of propagation peculiarly its own. In the larger ranks of existence, two animals that nearly resemble each other in form, will be found to have a similar history; but here, insects almost entirely alike will be found very dissimilar in their habits, and in the different changes which they undergo during their short lives *. To give a full enumeration of all the animals contained in this province of nature, would prove a task extremely arduous; one which no naturalist has ever pretended to accomplish. To furnish a distinct and complete history of each, is an undertaking for which the human powers feem altogether unequal; their impersection, at least, forbids us to hope that it will ever be effected.

But although a complete history of the operations of nature in this large and populous part of her empire cannot be expected; yet, such a general picture may be given, as shall demonstrate the existence of that great vivifying principle by which she is animated, and by which she is enabled continually to pour forth into existence such immense numbers of organized beings. A history of such insects as most frequently occur, and whose manners are best known, will present to us a pleasing view of that protection which providence affords even to the smallest of its creatures;—of the means it employs for perpetuating them;—and of that great arrangement of nature, by which one set of living beings find subsistence, by devouring another, and by which life is continued through every part of the creation, without a pause.

Qq2

Natura!

³ Goldfmith's Natural History, Vol. VII. p. 239.

Natural history claims it as its prerogative to demonthrate the existence and the perfections of that Creative Power which produced, and which governs the universe: It is the hiftory of the works of God, and naturally leads every intelligent mind to their Author; for there are no proofs of his existence more level to the apprehension of all, than those which it offers to the understanding. No rational man can ever blush for having placed among his occupations a fludy which has for its object the works of the Supreme Being which leads him to the contemplation of their author. All those naturalists who have described the different parts of the animal kingdom, have willingly confessed them to be productions of infinite power and wisdom; and seem to regret, that the experiments and observations relating to their history are not more numerous; because, in proportion as they are extended, the proofs of a Creative Power are multiplied.

The manner, however, in which entymology has too frequently been studied, and the extremes into which men, according to their different capacities and taftes, have fallen, have brought that obloquy and derifion against the science, which a proper degree of discernment would have directed against the foibles of those who studied it. While the fystems of some naturalists contain only a dry repetition of shades, colours, and shapes of different infects, without entering into the more interesting and animated description of their manners, those of others, as injudiciously, ascribe to them functions, and a degree of intelligence of which they are incapable. By the former, the imagination is fatigued and difgusted with a constant repetition of the same images: By the romantic air of the latter, the mind is led into distrust with regard to the truth of the whole narrative, and to doubt of those facts

which

which are well established and certain. Hence the study of entymology has been deemed by many, an occupation the most useless and frivolous in which the human mind can be engaged: Hence too, from a fear of prostituting their talents, many have been deterred from contemplating the wonders displayed by nature, in a kingdom of animals the most numerous, diversified, and splendidly adorned, of any on the face of the globe; and thus have deprived themselves of views of the power and munisicence of the Author of Nature, the most striking and interesting that can be presented to the mind of man.

To those who can derive no pleasure from the pursuits and studies of a liberal mind, and who feel no satisfaction in any employment that is not attended with immediate profit, the relearches of the entymologist cannot feem altogether ufelefs. Had the operations of the filk worm never been examined, how could men have availed themselves of the labour of an infect that administers so profusely to our luxuries and our wants? It was not to the unobserving that it first occurred, that the produce of that animal's labour might be converted into a confiderable article of commerce, and might give rife to many arts, and afford fubliftence to thousands of manufacturers. In the same manner, wax and honey enter into the articles of commerce, and add to our enjoyments. It cannot, therefore, be denied, that they were profitably employed, who first observed the industry of the bee; who brought that animal from its native woods, introduced it into our gardens, and, by domesticating it there, have rendered it subservient to our enjoyments.

The Chinese, whose progress in many of the arts is superior to that of any other nation, avail themselves of the labours of certain insects, in procuring a rich dye,

and an elegant varnish, which is provided by a certain species of winged ant. The celebrated purple dye of the ancients was the produce of a small species of shell-fish; and we are told by Pliny, that the discovery of its virtue was occasioned by a dog, who, in eating the fish, had dyed his ears with that beautiful colour *. It feems probable that the ancients were capable of communicating to their stuffs many beautiful shades of scarlet with which we are unacquainted; and it is not unlikely that we have also some rich tints of that colour which they wanted. It is certain that our finest red colours are furnished by insects with which they were unacquainted. Cochineal, the extensive and profitable uses of which have been long known, is now univerfally allowed to be an infect, which is propagated with care, and in vaft numbers, in the kingdom of Mexico. The kermes, or grain of fearlet, which was formerly imagined to be one of the galles or excrefeences that are feen on shrubs, is now understood to be an infect, which attaches itself in that form to a species of the oak t.

The medical uses of certain insects are far from being inconfiderable; and to these purposes they have long been applied, perhaps more frequently, and with better effect, than at present ‡. The valuable purposes to which the

can-

^{*} Reamure, Tome I. p. 5.

[†] The quercus coccifera of Linnæus. The red dye collected from this tree is produced in Africa; it is not fo bright, but more permanent than cochineal. The coccus quercus forms galles upon the common oak, which are brought from the Levant, and are univerfally used in dycing over Europe. We have this plant in Britain, and also that of the coccus polonicus: The infect which inhabits these plants might in all probability thrive, if imported into this island.

[‡] Hist. Medic. des Anim. par Vanden Bosche, Liv. iv.

cantharides has been made subservient, will alone vindicate the utility of those researches which have been made concerning this part of the animal kingdom. There are other uses to which insects have been applied, and that from the most remote antiquity, which appear of a still more fingular nature. Before the time of Theophraflus and of Pliny, certain kinds of them were employed in ripening the figs throughout the islands of the Archipelago *; and it appears that the same practice still subfifts among the present inhabitants of these islands to There are two kinds of figs cultivated around the Mediterranean; the wild, and the domestic. The former produces fruit feveral times in the year; and in it are produced certain worms, which are afterwards transformed into fmall flies. It is by the affiftance of these little animals that the domestic fig is brought to maturity, which would otherwife drop from the tree in an unripe state. During the months of June and July, the peafants of these delightful climes are bufily employed collecting fuch of the wild figs as abound most with these insects, and in placing them near the cultivated fig, that they may co-operate with the climate in bringing it to maturity. Similar purpofes might probably be ferved by a judicious application of infects to fruit in more northerly climates, were we acquainted with the proper fpecies. Those prunes, pears, and apples which are first ripe, are commonly found penetrated by worms.

But there are other inducements to the study of infects, of a nature totally different from those already mentioned; inducements, founded not on any hope of advantage to be derived from these animals, but of allering

leviating

^{*} Vide Plinii Hist. Nat.

[†] Voyages de M. de Tourneforte,

leviating or preventing the numerous mischiefs they occasion. Infinite swarms of these animals annually desolate whole provinces; others attack our gardens and cultivated grounds, where they commit endies devastations upon the corn, vegetables, and fruit trees. Nor are their depredations confined to the fields; they enter the habitations of man, and by destroying the timber, gradually reduce them to ruins. They destroy his furniture and clothing; some of them spare not even his person, tormenting it long before the period which nature has destined it to become their legitimate prey. Here, then, is a wide field laid open for study; and the person who could inform mankind how to remove or alleviate these mischiefs, would describe better of his species than if he had discovered the longitude.

Thus, by a closer examination of the destructive powers of infects, we shall have melancholy proofs of their importance in the fystem of nature, and be perfuaded, that however despicable they may appear, there is no class of animals whose history more nearly concerns us, and which better deserves the attention of the naturalist.

There are four different species of the local which are remarkably destructive. Almost every year, whole provinces, the most fertile in Asia and Asrica, are laid waste by their depredation. In Tunis and Algiers, swarms of the grillus migratorius appear so numerous, that they darken the face of the sky, like a thick cloud. These pernicious animals are wasted there by the southersy winds in the month of April: In May they take their departure for the interior parts of the country, to propagate their young; these make their appearance in their larva state, during the month of June, when they come mit vast depredations. The first columns, which pervade

the country like an army, deflroy every green shrub and pile of grass; and their devastation has not ceased, when they are succeeded by other swarms, that press upon their rear, devouring the tender branches and stalks of plants, which their forerunners had left. This dreadful visitation, which the language of Scripture has justly described as a plague, does not terminate till the infects have passed into their winged state, when they sly off, leaving the whole surface of the earth naked and brown, as if scorched by sire.

Little inferior to the locust in its destructive powers is the Phalæna Graminis of Linnens, which destroys the meadows in Sweden: There the peafants are employed in cutting deep ditches in the furface to ftop the progrets of the larvæ as they pass along: If the swarm be small, this device has the defired effect; but the numbers of thefe animals are often fo great, that they fill up the trenches, and pass along over the dead bodies that are buried in them. The formica facchilifera is a native of the West Indies, where it pervades the plantations of the fugar-cane, entering the plants, and destroying them when they are tender: After long experience of its depredations, the inhabitants have never been able to invent a method of destroying this pernicious animal. In our own country, the turnip fly, the butterfly, and the goofeborry worm, have long committed depredations in the fields and gardens, which no invention has hitherto been able to prevent: Against the last of these animals, indeed, the watering the bulbes with an infufion of towacco, has been found efficacious, by killing the greater number in their larva itate.

Vol. III. Rr Another

Another object highly worthy the attention of the entymologist, is the means of preserving corn from the invasion of infects, after it is collected into granaries. Our fubfishence in almost every stage of its progress, is conflantly exposed to the intrusions of these enemies of human industry. Flour, biscuit, and almost every kind of provision, even after it is barrelled up for exportation, is liable to be devoured, or rendered useless, by the depredations of the most hideous animals. The patriotism of statesmen, and their zeal for the good of mankind, could not receive a nobler or more useful direction than in holding out rewards to fuch as might discover the most effectual means of preventing the ravages of those animals, who, by the most destructive activity, are continually converting large flores of provisions into fo many masses of corruption.

May it not be hoped, that by a careful study of the nature of infects, fome means may be discovered to prevent them from penetrating into the joists of buildings, and thereby reducing them into dust, and effecting the destruction of the most costly edifices. How often do we find wooden furniture destroyed by insects, which might have answered the purposes for which it was intended for ages, had it been preserved from the destructive attacks of these animals? How many accidents are probably occafioned at fea by those formidable worms, whose heads are armed with hard shells, and who are thereby enabled to gnaw through the thickest vessels, and make perforations under the water? The alarms they have frequently occasioned in Holland, by introducing themselves and multiplying among those wooden stakes which support their dikes, are univerfally known. The naturalist who should difcover

discover a mode of preventing such devastations, would certainly deserve well of his country and of mankind, by communicating information that tended so directly to the public good.

The tar extracted from coal by that ingenious nobleman the Earl of Dundonald, when better known, will probably be found an efficacious remedy against the corrosions of wood by infects in all cases where it can be applied. It not only penetrates so deeply into wood that it cannot be washed away; but is of so acrid a substance, as must inevitably destroy worms. The intolerable effluvia which it emits, will, however, prevent its application in the case of household furniture.

The pernicious genus of phalæna, which contains all the different species of moths, makes still nearer approaches to man in the hostilities which it commits. No person is ignorant of the destructive quality of these insects to woollen cloth, and all kinds of fur and wearing apparel. The association instinct of these animals, in providing a proper receptacle for their eggs, and food for their young, have not withdrawn that indefatigable entymologist Reaumure, from devising efficacious methods of preventing their depredations upon woollen stuffs and furs *.

Of fuch vast extent are the mischiefs occasioned by the insect tribe upon the various objects of human industry, and the necessaries of life; all these, however, would in a manner disappear, were we to resect upon their dangerous effects upon the human body, and the husbandman's attention would be withdrawn from the depredations committed on his fields, did he imagine that thou-

fands of the same noxious race were continually entering into his lungs by breathing. During the whole of the fummer months, the atmosphere teems with myriads of fmall animals, and particularly in the months of July and August: It is then that the excessive numbers that are conveyed into the flomach and lungs, probably generate those epidemic disorders for which that season of the year is to remarkable. If we pass from those dangers which are less visible, to others more apparent, with which infects threaten the human race, we shall fill find room to commiferate the inhabitants of many parts of the globe. What an uncomfortable life must the poor Laplander lead, fince, at certain feafons of the year, the number of infects is fo great, that a candle is no fooner lighted than the flame is extinguished by the multitudes that flock to it; where, after millions are deftroyed, famished millions succeed, and renew the unceasing combat. Even in Britain, which is happily free from thefe unrelenting invaders, much inconvenience is often felt from the bug, the hornet, the wasp, and the bee *.

To counteract these inveterate enemies of man, and to relieve him from the mischies they occasion, ought always to be one aim of the entymologist: and no person will deny, that whoever, by the study of insects, has sound the means of availing himself of the labours of such as are useful, or preventing the noxious from doing harm, hath rendered an essential service to mankind.

After all, however, it will readily be allowed, that in purfuing the history of infects, the number of useful observations

^{*} The first of these animals has been found to be effectually destroyed by the smoak of peat.

observations with which it presents the reader, bear but a finall proportion to those whose object is merely to gratify curiofity. But in what science is there nothing to be found but what is immediately useful? and are not objects of curiofity often nearly allied to those of utility? Is it not while we amuse ourselves with the former, that we are most frequently led to the discovery of the latter? To these circumstances, we may add another for the encouragement of those who undertake to write upon this fubject; that works the most useful are not always the most favourably received: The number of those who read for amusement, is at least equal to those who read for inftruction. A taste for the marvellous, in a greater or less degree, is universal among men: It is this which leads them to prefer romances, novels, Perfian and Arabian tales, to the incidents of real history. The generation, the metamorphofes of infects, their means of rearing their young, and of procuring food, prefent to the reader wonders, perhaps not inferior to those that are fabricated by the most licentious imagination; with this difference, that the latter are true. The entymologift therefore, has himself rather than his subject to blame, if his works are not read by the public with avidity.

SECT. II .- The Literary History of Entymology.

THE class of insects, although by far the most numerous in the animal kingdom, the most remarkable for its effects on the objects of human industry, and perhaps the most important in the economy of nature, was long before it attracted any confiderable degree of attention from men of learning. Hardly any attention was bestowed on entymology by any of the writers of antiquity. Aristotle, that father of naturalists, has allotted but a small portion of his works to the history of insects; nor does it appear that he has been always fufficiently attentive to the authorities upon which he inserts the different facts he relates, or to the order in which they are arranged. It is known, that Alexander his pupil furnished him with confiderable fums to be employed in the various refearches necessary for the compilation of his hiftory of animals; and it is not improbable, that many persons were engaged in different parts to procure him the different species, and to furnish him with such observations on their history as fell not within the reach of his own knowledge. From feveral of the facts which he has related, we are led to suspect, that these men were not altogether qualified for the task in which they were employed. The arrangement followed by the Grecian naturalist seems as defective as the authorities upon which

his observations are founded. It consists of a long and tedious enumeration of animals, whose appearance and history correspond in certain particulars; followed by another of those which differ from the former. Enumerations of this kind are seldom complete, and long before they can become so, must prove a burden too great for the memory to retain*. The histories of Pliny and Elian are formed upon the same plan, and are liable to similar desects: They consist of a number of observations ill arranged, and insufficiently authenticated.

During that long fuccession of ages, which was only diftinguished by ignorance and barbarism, entymology shared the same sate with every other science: It was condemned to oblivion. After a taste for literature had begun to revive, the history of infects again attracted the notice of the curious; unhappily, however, for the growth of science, men were then devoted to the study of the ancients with a blind admiration: It was from their writings that they imagined the moderns were to derive a complete knowledge of all the fecrets of nature; and Aristotle was principally consulted for the history of animals. Had Aldrovandus, Gefner, and Mouffet, beflowed the fame attention in studying the works of nature that they employed upon the writings of that naturalist, they would have made a much greater progress in real knowledge: But they unfortunately observed nature only to observe there what they had read from Aristotle. This excessive predilection for antiquity ough: not indeed fo much to be imputed to these authors as to the age in which they lived; a period when every thing was deemed

^{*} Vide Hist. animal. passim.

deemed contemptible, that was not found in the writings of the ancients.

The attempts of these early writers gave birth to the researches of others, who were gradually more bold and fuccessful, in proportion as their reverence for antiquity was diminished. In 1668, about twelve years after Monfist published his Theatrum Infectorum, the experiments and observations of the celebrated Rhedi made their appearance in Italy. His investigations were principally directed to the manner of the generation of infects, with a view to overthrow that abfurd and erroneous doctrine established by the ancients, of their arising fortuitously from different bodies in a state of putrefaction. Nothing can more fully demonstrate the strength of prejudice than those elaborate treatifes which Malphigi, Swammerdam, and Rhali were obliged to compole, in order to combat the notion of the fpontaneous generation of these animals; an opinion which at prefent feems as ridiculous and anphilosophical as it is untrue. Notwith tanding all their efforts to prove, that the finaller animals are produced in the fame manner with the larger, and that the organization of the Lody of a mite requires the fame apportus of limbs, and the lame delicate thructure as that of an elephant, attempts were still made to revive the ancient error by Kirker, Bonarie, and others *. And what is most mortilying to human reason, the same Rhedi, the declared enemy of prejudices, and the man of all others who knows best how to combat them, has, upon this very subject, fallen into a similar error. In order to produce those insects which are found

^{*} Vide Motivi di dubitar intorno la generatione de viventi, fecondo la opinione de moderni.

found upon the small excrescences of plants and trees, he found himself obliged to confer a vivifying power, a kind of soul upon those vegetables where they were found, and has laid aside the ordinary mode of generation, which in other instances he had laboured to establish.

Swammerdam was the contemporary of Rhedi; and, like him, he possessed the courage to examine nature, and to think for himfelf. This naturalist made many anatomical observations upon insects, which after his death were published at Leyden *, and laid the foundation of future improvements in entymology. About the fame period, Madame Marianne, a Dutch lady, contributed largely to bring the history of infects into request, by the beauty of her paintings and drawings. After having executed elegant drawings of several of the insects of Eucope, from a fingular avidity for these studies, she was prompted to cross the Atlantic, and give paintings of those in America. Having refided for feveral years in Surinam in South America, the returned to Europe with exquisite drawings of many of the splendid insects of that continent, which were afterwards engraved and published in Holland, about the end of last century +.

Goedart is another of the first authors who adorned the history of infects with the labours of the pencil. He paid great attention to the metamorphoses of the animals, and has painted many of them in the several forms which they assume, from their appearance till their death. His work was originally published in German, very badly ar-

Vol. III. Sf ranged;

In the year 1737, under the title of Biblia Naturæ.

[†] Her original drawings were purchased by Sir Hans Sioane, and are now in the British Museum.

ranged; a new edition in Latin was afterwards given by Dr. Lifler, in which many errors were corrected, and a new arrangement made out by that able naturalist, who has himself written a valuable treatise upon the intricate genus of spiders.

Some of these authors, by the extent of their labours. and others by the boldness of their genius, had convinced mankind that the ancients were far from attaining that perfect knowledge of nature which they had hitherto imagined. They were now fo far emboldened by the progrefs they had already made, that they trufed to their own judgment and observation in examining the works of nature; and many productions far more judicions, accurate, and philotophical, were offered to the public, than had ever hitherto appeared. Among the first of these may be ranked that of our celebrated countryman Mr. Ray, who had, for the greater part of his life, alliduously examined the economy of infects. His Methodus Insectorum was not published till after his death in 1710, and may be regarded as the most accurate and concife performance on the subject of entymology. In the meanwhile, there appeared a numerous lift of au. thors who treated this class of the animal kingdom; Albin described the insects of England, while Sir Hans Sloan, Petivert, Catabr, and Frisch, have detailed the history of vast numbers of exotic animals of this tribe.

In this state of the science, about the year 1754, appeared M. de Reaumar, by far the most laborious and indefatigable entymologist in Europe. Almost in every part of France this naturalist had correspondents stationed to transmit him by post descriptions and specimens of every curious and rare insect that might occur: And in order

order to examine their inflincts, their metamorphofes, and mode of generation, he inclosed vast numbers of them in a large aviary, constructed with a close net and fecured below with a pavement overlaid with green turf, and planted with shrubs and different kinds of plants. It was there that this unwearied observer of nature examined the manners and economy of those insects, which he has described in a work the most voluminous that has hitherto appeared on entymology. As a writer, he is extremely diffuse, but always entertaining and instructive. The principal defect in his works is the want of a fystematic arrangement, and the fynonima of other authors; a precaution that was alone able to have prevented confusion in fo large a performance, and amid the vast number of infects which he has described. These desects, however, were foon remedied by that excellent arrangement which the immortal author of the Systema Naturæ has introduced into entymology, as well as every other department of natural history. No philosopher hath ever yet appeared who posiciles the same powers of classifying the different productions of the various kingdoms of nature; and his fythem will probably long continue to be a flandard to all those who shall afterwards attempt to make improvements, or to enlarge the boundaries of natural history.

Since 1715, when the fystem of nature was first published by Sir Charles Linnaus, various improvements have been made on it by that author, who has enjoyed the rare felicity, not only of seeing his works gain universal approbation, but of many opportunities of bringing them nearer to perfection. In the mean time Geostey, Scopeli, Wotton, Harvey, Valifnieri, and many others of inferior

ferior note, have turned their attention to this interesting part of natural history; and while they have endeavoured to improve upon the arrangement of the Swedish naturalist without success, have nevertheless added greatly to the number of insects. Among this number De Geera Swedish nobleman, councellor to the king, may be ranked in the first station: He has published a large treatise upon this subject, in which he has not only described a great number of insects with accuracy, but has indulged in philosophical remarks upon their history, which possess very considerable merit.

Besides these, a number of entymologists have appeared, who have given most elegant engravings of insects; among which are, Roefel, Lewenbock, Baker, Barbut, Harris, and Drury. Some of these have improved this science by microscopical observation: the last has given excellent drawings of the exotic insects, while Harris has applied himself successfully in delineating those of England.

After reviewing the numerous lift of authors who have laboured in the fame field, we should widely err, were we to conclude that the science of entymology had reached perfection, or that the whole of this class of beings had been fully made known. The far greater part of insects, as we have already hinted, from their extreme minuteness, elude the observation of the naked eye; many, it may be presumed, are not perceptible even by the assistance of the best microscope; and of these small animals, whose bodies are scarcely discernible, how little can we know of their organization, their food, their manners, and their history? In the latest edition of the system of nature, near three thousand insects are enumerated by

Linnaus:

Linneus: There is, however, good reason to apprehend, that this enumeration does not contain above one half even of the larger infects that are distinctly observable by the eye. Whether in a hot or cold climate, it is probable that the number of infects in every country is greater than that of plants. In Sweden, there have been enumerated, and pretty accurately described, about one thousand seven hundred infects, while the number of plants, though no where more accurately examined than in that kingdom, do not exceed thirteen hundred. In the neighbourhood of Paris, there have been discovered by Geoffrey and Reaumur, upwards of one thousand four hundred infects, while the plants of the same district, which have been still more accurately examined than the infects, do not nearly amount to that number.

The British infects have never yet been either fully enumerated, or accurately described. Dr. Berkenboot, in his outlines of the natural history of this island, has enumerated five hundred and seventy-two different species: In all probability, a more complete investigation would discover three, or perhaps four times that number of British infects. The nearer any country approaches to the equator, the more numerous will be the infects found in it: Even in these northerly climes, however, there is still much room left for entymological investigation. If, in travelling over this ground, we cannot flatter the reader with much new matter, we can at least assure him, that he will not be led assure by theory, or deceived by unauthenticated narrations.

SECT. III.—A comparative view of the Senses and Endowments of Infects, with those of other Animals.

THE more closely we examine this class of the animal kingdom, the greater number of furprifing facts and wonderful inftincts shall we find, to indemnify us for those large portions of the marvellous, of which we shall often be obliged to divest their history. Many entymologists, it must be allowed, particularly those who wrote in the earlier stages of this science, before philosophy had thrown much light upon the operations of nature, from a defire of filling the reader with the fame admiration which they themselves felt, have much oftener had recourse to the marvellous, than can either be justified by facts or observation. Those eulogies which we find fo frequently beslowed, without measure, upon the intelligence of certain infects, afford pregnant evidence of this indifcretion. There is hardly any kind of knowledge, endowment, or even moral virtue, of which fome or other of them has not been faid to be possessed. They have been made to act and think like men; and fometimes have been celebrated for accomplishments of which few of these lords of the creation can boast; and all this on the most puerile and false foundations. For example, there is a species of the mantis, with long limbs, and of an uncommon appearance; and because this animal is frequently feen in an erect posture, having the two fore

legs crossed, it has been said to be employed in devotion; a circumstance from which it has in some parts obtained its name *.

The same infect is said to have such a regard for children, that, should one lose its way, it humanely points to the proper way with its leg, and feldom is known to give a wrong direction. Of this kind, too, is the respect for their dead, which has been afcribed to the ants, and the decency with which they bestow the honours of interment upon their deceased friends. These reports rest upon no better foundation, than that these animals, and fome species of bees, are feen carrying from the hive fuch as have perithed or been killed. How many wonders of a fimilar kind have been told of the common bee; wonders which render the greater part of the early treatifes upon them hardly more instructive than a romance. Their affociations have been deemed a perfect model of a monarchical government, conducted by one fovereign, whose commands are implicitly obeyed, and regulate all the different operations in which they are fo bufily employed. From farther investigation, however, this king of the bees has a queen, and fo prodigiously prolific, that from three or four, which are to be found in every hive, the whole progeny, which confifts of many thousands, is derived.

The different operations which infects perform, no doubt oblige the naturalist to ascribe to them a certain degree of intelligence; some degree of it we have seen possessed by the other animals: From analogy, therefore, we may reasonably suppose them not altogether destitute of fagacity. The different processes through which they

gc,

^{*} Vide Muosset, p. 118. In Provence, it is called the God-prayer.

go, are fo uniform, and that fuccession of actions which we fee them perform, are fo little varied, that nature feems to have deflined them to act a subordinate part in the great republic of animals. If their history fometimes exhibits them varying their procedure, and accommodate ing themselves to circumstances, the sphere of their addrefs feems to be confined within narrow limits; and it is feldom that they attract our admiration by any extraordinary efforts of fagacity. The bee and the ant present firiking inflances of affiduity in labour; they have, however, but one fingle method of operating, and, withdrawn from that, can turn to no other. A bee taken out of the hive is totally helplefs and inactive; far from displaying the fagacity of the dog, it feems incapable of giving the fmallest variation to its instincts. In the pursuits of the latter animal, there appears fomething like choice; in the labours of the former, the whole appears like necessity or compulsion *.

The conformation of the whole infect tribe feems also to argue its inferiority to many other parts of animated nature. In the external and internal structure of the body of one of these small animals, where every member appears completely formed, and co-operates with the rest in carrying on the vital functions, there is no doubt displayed a very wonderful organization. Insects, however, of all animals, are perhaps the most impersectly formed; nor is this affertion founded on any investigation made upon these minute creatures by means of the eye, or the knife of the anatomist; it is obviously deducible from their capacity of subsisting after being deprived of many of those members and organs, which, in the higher ranks

of

of nature, are absolutely necessary to life. Many of them, like the nobler animals, are furnished with lungs, and an heart; yet the caterpillar lives, though its heart and lungs, as is often the case, are entirely eaten away. In some, the stomach perishes altogether, and is again renewed; others, after being cut into feveral pieces, not only continue in life, but are formed into as many new distinct animals as there were segments of the old! As in mechanics, the most complicated machines are required to perform the nicest operations, so, in anatomy, the noblest animals are most variously and wonderfully made *. Of all living beings, man offers the most won. derful variety in his internal conformation; quadrupeds come next; and after them the other animals follow, in proportion to their powers and excellencies; while infects feem to fill up the last and lowest rank of animated nature; fome of them being fo imperfectly organized, that they have long remained in the fystems of naturalists confounded with the vegetable tribes.

The amazing number of infects is another argument of their imperfection. It is a rule which obtains among the offspring of nature, that the nobler animals are flowly produced, and that, in forming these, she acts with a dignified economy; while, in her meaner births, she is often lavish to profusion; and thousands of the more ignoble kinds are produced merely to fupply the necessities of the more favoured and delicately organized parts of her creatures. Of all the other productions of nature, we have feen that infects are the most numerous; that, however minute when taken individually, when taken together they are probably more bulky than all the rest Vol. III. Tt

* Idem, ubi fupra.

of her animated offspring. The plants and vegetables which cover the furface of the earth, and which at first fight appear so far to outnumber the other productions of nature, upon a closer investigation, are sound far inferior in number and variety to the insects which swarm in the wider range of the air, the earth, and the water.

From these considerations, we are induced to believe, that the degree of intelligence allotted to this class of the animal kingdom is but very fmall; and our experience, as far as it goes, coincides with this opinion. While other animals are capable of fome degree of education, thefe have one invariable mode of operating, which no art can either alter or improve. The dog is taught to carry; the bird to whitle a tune; but those infects which may be confidered as completery domesticated, an by no invention be turned from their inflinct. The filk worm completes its labours, and the foider confiruds its web, invariably in the iame manner: An existence which continues but a fingle feafon, feems too thort for the purpofes of inflruction, or of learning Hence the infects are not only or a rank inferior to the other animals, but many of them feem more nearly allied to the vegetables than to the classes above them. Many are attached to one vegetable, forme to a fingle leaf, where the period of their lives is completed in a few weeks, or perhaps a few days, and where the pleafures they enjoyed, or the purpofes for which they were produced, are in a great meature beyond the reach of our faculties to explore.

The external fenses of infects, as far as we are enabled to judge of them, correspond with the low measure of sagacity which the Author of nature has assigned them: Of some of them they seem altogether destitute, while others are enjoyed but in an impersect manner. It has

long been a received opinion, that hearing is denied to infects; or at leaft, that the exidence of this fense is very equivocal. Many of them, however, are endowed with the power of uttering founds; as the bee, the fly, the gnat, and the beetle. The Sphinx atropos squeaks when hurt, nearly as loud as a moufe; it has even the power of uttering a plaintive note, in certain circumstances, which excites commiseration. In general, the power of uttering founds agreeable to the feelings and necessities of animals, is conferred on them for the purpose of communicating these feelings to the rest of their kind. We have already feen, that fuch of the finny tribes as were vocal, were also endowed with organs for the reception of founds; the fame, probably, is the case with the infect tribe. For what purpose is the individual possessed of the power of expressing its pleasures or its pain, if all knowledge of found is denied to its tribe? Were the fense of hearing withheld from the animals of the same class, it must crave assistance in vain; for it must speak a language destined to be unintelligible to every being in nature *.

Experience daily convinces us of the truth of the foregoing positions. If a bee or wasp be attacked near the
hive, the consequence of this assault commonly is, that
the animal expresses its pain or indignation in a tone disferent from its ordinary noise; the complaint is immediately understood by the hive within, when the inhabitants hurry out to revenge the infult, in such numbers
that the offending party feldom comes off with impunity.
The same evidence of hearing is continually afforded by
the spider: Often his webs are of such an enormous

Tt 2 length,

² Vide Barbut's Genera Infectorum, p. 4.

length, that he cannot fee from the one end of them to the other; often too, in watching for his prey, he conceals himself in some adjoining crevice, where he cannot fee those animals that are ensured in his toils. The fly, however, no sooner finds itself entangled, than it makes a buzzing noise, in order to escape; this noise is instantly heard and understood by the spider, who sallies forth from his concealment, and riots in the spoil, with all the eagerness and serocity which distinguishes the most rapacious animals *.

If the fense of hearing has with difficulty been allowed to infects, naturalists have had still more in afcertaining the place or organ where that fense is fituated. The most accurate observers of nature have supposed it to be placed in the antennæ: These, from their situation in the head, from their inward structure, and their capacity of motion, have been supposed most favourable for the seat of fuch organs. Besides, it is dissicult to assign any other use for these instruments, which are common to almost the whole of this class of beings. From their extreme fensibility, they seem neither fitted for the purposes of attack nor defence: While, then, it is certain that providence hath made nothing in vain, but hath framed every limb of the fmallest animal for use, as well as ornament, it feems to follow, that the antennæ of infects are the appointed organs of hearing, fince there is apparently no other purpose in the economy of these animals which they feem fitted to ferve.

The antennes of all infects are composed of joints, varying in form, fize, and number. Among those which are confined to live mostly under water, as the gyrinus,

they

[.] Genera Infectorum, ubi fupra.

they are in general short; while such as roam at large through the air have them long and flender, as the phalenæ ichneumons, and others. They are all hollow within, and rendered flexible by the joints, which are very visible in those of the crab; which are the best examples, because the largest belonging to this class of beings. This hollowness, it is supposed, is intended to receive the found communicated to the extremities of the antennæ, by the repercussion of the air; and to convey it, by means of the joints, from one piece to another, till it arrives at the brain, in that leffened degree of tone which is fuited to the nature of the particular animal *. Such is the apparatus destined for the hearing of infects; a fense which they probably enjoy in various degrees of perfection; but in none of them does it feem fo acute as in the higher ranks of animated beings.

But, befides the fenfe of hearing, it feems highly probable that infects possels also that of smell. As many of them live on bedies in a flate of putrefaction, around which, when exposed, they are feen immediately to collest themselves, the conclusion feems obvious, that they are possessed of organs sitted to direct them to their food. Those which feed on herbs, flowers, or fruits, feem in the fame manner to require fenses adequate to their purfuits. It has therefore been supposed, and not without an appearance of probability, that the palpi, or feelers, are the organs of smell in the insect tribe. These instruments are four, fometimes fix in number; two of which are evidently destined to the purpose of handling their food, and conveying it to the mouth. The others, which are in continual motion, and confiantly applied to those objects

a Genera Insectorum, ubi supra,

objects on which they alight, feem employed, like the fnout of a hog, in fearching for food, and examining the quality of the different kinds of fustenance by which they are supported. The beetles, and other tribes whose palpi are large, are proper subjects for examining the organs of smell in this class of beings; and an accurate investigation of their manners would probably justify the foregoing conjectures with regard to the uses of the feelers, and determine how far they possess this sense.

The organs of vision among most kinds of infects are large; a circumstance which has put their fense of feeing beyond a doubt. The eyes are commonly two in number, each frequently confitting of a congeries or affemblage of lentes, covered with a crustaceous transparent fubstance, to protect them from injury. The organs which have been alletted to hearing and finell, are also protected from dust, and the fmaller particles of those fubflances to which they are applied: Their extremities are not patulous, but ciliated, like those of the mole, to prevent them from being elogged or injured by the intrusion of furrounding objects. Hence it appears, that though infects are destine; to all a subordinate station in the animal kingdom, yet nature has by no means nogl &ed them, but furnished them with organs wonderfully adapted to their humble purtuits, and to that tranfient existence which she has assigned them.

Sect. IV.—Of the External Parts and Classical Gharacters of Infects.

Institute have always been confidered as a diffinct class of the animal kingdom, though naturalists have not agreed in afcertaining its limits. The shades of nature are indeed intimately blended together; and those links by which she connects different portions of her animated offspring, are often small and imperceptible. The infect tribe comprehends those small animals which are destitute of red blood, bones, and cartilages; which are furnished with a mouth, or else a trunk, opening lengthwise; and which breathe by means of stigmata, or apertures upon the external parts of the body. They have obtained the name of infects from the frequent incisions by which their bodies are apparently divided into several parts or segments.

The body of these animals is divided by naturalists into a head, thorax, abdomen, and limbs. The head is, for the most part, distinct from the thorax, being attached to it only by a slender tendon, and is surnished with eyes, palpi, and antennæ, which we have already seen are the organs of the disserent scales. This whole tribe of animals is supposed by Linnens to be destitute of brains *.

The antenne are organs peculiar to infects; and according to their various forms and proportions, afford

fystematic

Vide Systema Natura, Tom. I. Part I. p. 533,

fystematic writers the most proper characters for the ara rangement of those animals. According to their form, they are either setaceous, filiform, moniliform, clavated, capitated, fissile, pactinated, or bearded; and in their proportions, they are either longer than the body, or shorter, or of equal length with that part.

The fielers have already been noticed, as constituting the organs both of touch and of smell; and have two, three, and sometimes four joints. Some infects are faid to have no mouth; in general, however, that organ is situated under the head, and to it the seelers are attached: In some species, the mouth is placed under the breast; in others, the restrum is of considerable length, having an upper lip, transverse jaws, teeth, and a tongue, not unsrequently rolled up in a spire. The stemmata, or small eyes, are three brilliant convex spots, situated upon the crown of the head.

The thorax is placed between the head and the abdomen, and is that part to which the fulcra or limbs are attached: immediately behind it is placed the abdomen, containing the stomach and viscera. It is divided into five fegments, each pierced on the fides with small foramina for the purpose of breathing. It was long imagined, that all the animals that are destitute of red blood lived without respiration: It has, however, been found by experiment, that this is not the case; and that, among the infect tribes, breathing is carried on, though in a different manner, from what takes place among the larger animals. Infects are all furnished with minute organs, which, in the language of naturalists, are termed fligmata; these are a number of small tubercles, ranged along each fide of the body, each having an aperture in the top, called the spiracle, by which the animal breathes.

Thefe

These stigmata are commonly situated on the sides both of the thorax and abdomen; and their number is various, being from eight to twelve. When the uses of these organs at sirst began to attract the attention of naturalists, it was imagined, that by means of them the insect only inspired, and that the air was ejected by the pores in the common manner, by perspiration.

This opinion was adopted by M. de Reaumur*; but subsequent experiments have proved, that the air is both introduced into the lungs, and emitted from them by the spiracles: If the stigmata are covered with oil, respiration totally ceases, and the animal dies; if they are covered only on one side, the vital sunctions on that part are impeded, and the side becomes paralytic. Nor is it in their winged and active state alone, that insects breathe. The crustaceous shell, by which the chrysalis is covered, is also provided with lateral stigmata, by which respiration is carried on during the period of their pupa state. In the breathing of insects, there is still another peculiarity; they thrive in air tainted by putrifying substances, and are capable of subsisting in phlogisticated air, the inspiration of which is so state to other animals.

From the limbs of infects are obtained the most permanent and striking generic characters; these are most commonly taken from the tail, the legs, or the wings, in such subjects as are surnished with them. The tail terminates the abdomen; and sometimes has two horns, and sometimes none: It is either simple; or armed with a forceps, a bristle, a claw; or a sting ‡.

Vol. III. U u The

^{*} Vide Memoir pour fervir a l' hist. des insecles.

⁺ Bonnet.

‡ Barbut's genera infectorum.

The fect of these animals are generally six; the crass and spiders have eight, and the scolopendra has a much greater number. They are divided into the thighs, which are attached immediately to the body; the tibia or shanks immediately below the second joint; and the tars, which are composed of various articulations, and are terminated by nails: The hind feet receive different appellations, according as they are formed, for executing the various movements of walking, running, leaping, or swimming *.

The wings are in some subjects two, and in others sour in number; and are so various in their colour, shape, and cendificacy, that they afford many characters for the distribution of these animals. The elytra or wing-cases are two, formed of a cruitaceous substance, and for the most part moveable: They serve as a cover to the under wings, and surnish distinguishing marks to the naturalist. Under the wings of dipterous infects are placed the halters or poisers, which are composed of a small stalk, terminating in a round knob or head; and are supposed to serve the purpose of balancing the animal, as is expressed by their French name †.

After this short account of the external parts of infects, and explanation of the technical language by which they are expressed, the reader will be able to see the propriety of the *Linnean* division of this class of animals into seven orders; an arrangement which constitutes the basis of a system of entymology the most simple and judicious that has ever yet been devised.

The first order of infects is called the colcopterous: It consists of all those animals that have membrenaceous wings,

^{*} Hence they are called, curforii, faltatorii, and natatorii, &c.

⁺ Les balanciers.

wings, covered with crustaceous elytra, and is subdivided into thirty different genera, according to the shape of the antennæ.

The fecond order is called the hemipterous: because the wings are half covered with crustaceous elytra, and are less hard and robust than those of the coleopterous infects, but more strong than those of the membren accous winged i sects that compose the subsequent orders. The upper wings are semi-coriaceous: They do not meet together in a longitudinal suture, as in the sirst order; but have part of their interior margin crossed or lapped, the one over the other along the upper part of the abdomen. The mouth and proboscis of the insects of this order, are bent inwards towards the breast.

The third order comprehends the various tribes of moths and butterflies, whose wings are covered with imbricated scales, and which on that account are called lepidopterous infects. They have four membranaceous wings; their bodies are rough, and the mouth furnished with a spiral tongue, which they can roll up, or unfold at pleasure.

The neuropterous infects conflitute the fourth order, which comprehends all those genera which have four naked and membranaceous wings, reticulated with veins. The tail of these animals is unarmed, having no sting, but is provided with appendices like pincers, by which the sexes are distinguished.

The infects of the fifth order have the tail armed with a fling, excepting the males, which have not that offen-five inftrument. They have four membranaccous wings, and are called the hymedopterous infects.

The fixth order contains the depterous infects, or fuch as have only two wings: They are faither distinguished U u 2 by

by a poifer fituated under each wing, the base of which is covered by a small scale, and the extremity terminates in a knob.

The last order comprehends in it all the apterous infects, or such as are entirely destitute of wings in either fex.

In giving the history of a class of animals so extremely numerous as that of infects, it becomes abfolutely necessary to group them in certain tribes, whose manners or external characters correspond: Nothing but this expedient can prevent endless confusion among fuch multifarious forms, or afford any prospect of finishing a task, that, at first view, seems involved in such inextricable difficulty. All naturalists who have treated of this part of the animal kingdom, have accordingly endeavoured to arrange them into orders and genera. Swammerdam and Ray feem to have founded their fyftems on the different changes which these animals undergo, and have formed them into four great divisions, agreeable to the different forms under which they appear: Valificieri has also distributed them into four orders, according to their habitation; arranging together in one group, fuch as inhabit plants; placing in another, thefe that live in the water; and in a third, fuch as conceal themselves under the earth or fand; reserving for his last division, those that inhabit the bodies of other animals *. Both those systems are defective, in having too few divifions of a class of animals so extremely numerous; the last, however, is liable to an imperfection of another kind; because many insects change their habitation, at the moment of their metamorphosis. Some are aquatic, which, after their transformation, are feen inhabiting the trees and plants; many of the fubterraneous infects in like manner rife into the air, so soon as they arrive at their winged state. Several other attempts have been made towards the arrangement of infects, none of which seem to have been so successful as that of the Swediß naturalist above described: By his system, therefore, we shall be chiefly guided in the following sketch of the history of these animals.

SECT. V.—Of the Generation and Metamorphofis of Infects.

We have already noticed the ewoneous opinion which the ancients entertained concerning the generation of infects, and mentioned the efforts made by Rhedi and others, in order to combat it: After all their refearches, however, this part of our fubject is far from being free of difficulty. Infects are remarkable for a greater variety of fexes than any other class of animals. In general, each individual is either male or female; but there is one order (that of the hymenoptera, which comprehends all the numerous tribes of bees, wasps, hornets, and ants), among which there is found a numerous race of animals that are neuters. These take no part in propagating the different species to which they belong, but seem devoted to the service of the more perfect animals. This sact is

fully

fully afcertained by the history of the common domestic bee: all those which are employed in collecting the honey, and constructing the cells, and which constitute by far the greater part of the hive, are of the class of neuters, to whom nature has denied the power of propagating their kind. It is only upon the drones, and one or two females in each hive, that she has conferred the fexual distinction, and the powers of generation.

Witherto we have uniformly beheld animals either male or female; and the neuters of the order of hymenoptera which form the first exception to the general law of nature, appear not only an anomalous but a defective race. The history of infects exhibits another instance of animals deviating from this rule, from a different cause. All the animals belonging to that genus, termed the aphus puccron, appear to be hermaphredites of the most perfect kind: A single animal of this tribe, though kept in the most careful manner from every other, will propagate its kind by itself; and if the offspring thus produced be preserved, it will also breed. An ingenious naturalish has well ascertained this fact by experiments, which he has repeated to the ninth generation *.

This mode of generation, so different from that effected by the joint co-operation of the sexes, naturalists have asserbed to a different power, something resembling that possessed by the vegetable tribes: We have already, however, seen instances of animals among whom secundation by the male, impregnated the semales for several generations. If that sact be well ascertained, it will explain the peculiarity of that tribe, without investing it with powers so different from all those possessed by the rest of the animal kingdom.

Infects

Infects are in general oviparous, producing eggs which are gradually quickened into life, by the joint influence of the heat of the fun, and of those warm substances which constitute their nidus. Bonnet mentions some inflances in the order of diptera, in which the parent infect produces living young. The genus aphlus exhibits a singular phenomenon. This animal, during summer, is viviparous, but towards winter becomes oviparous; the state of its progeny being determined by the nature of the season.

The nidus in which the eggs of infects are deposited, is generally chosen with admirable skill; being adapted equally to the fecurity, warmth, and fubfillence of the future larvæ that are to be reared in it. Some construct their nests in the earth with great labour: others depofite their eggs upon those plants, the leaves of which are to impply food for the natcent brood: while feveral of the musca eject their eggs in the body of the chryfalis of other infects; upon the juices of which the young are nourished at the expence of the defenceless animal which they devour. Inflinct is an unerring guide in directing each of these animals to a nidus fitted for the fervation of the ova. In one inftance, the common blue fly is faid to be deceived; and that is, when it dejotics its eggs upon the flower of a certain plant of a putrid fmell, mistaking it for sless in a state of putretaction. There the young are no fooner quickened into life than they die for want of proper food #.

The

^{*} The fkin, the nostrils, the anus, and vicera of quadrupeds, fome-times furnish a receptacle for intects; and there nature directs the parentanimal to deposit its eggs.

The different changes of form which many infects undergo, from their first appearance as eggs, till they arrive at their perfect and winged state, constitute an important article in their history: These have been termed their metamorphoses, or transformations; and, from the very language employed to express them, the salse notions which were long entertained, even by naturalists, are still discernable.

A fly, a spider, or an ant, insects of the most different kinds in outward appearance, do not differ more widely than the same insect does from itself, under the different forms of a worm, a chryfalis, and a butterfly. What is at prefent a worm, however, foon becomes a chryfalis, which is again as fuddenly to be changed into a winged animal. Changes apparently fo inftantaneously produced, have been compared to the metamorphofes fo renowned in ancient fable, and probably at first suggested the idea of those transformations which fable has rendered fo celebrated. When an infect in fo fnort a space, appeared under a form so different from that which it lately exhibited, men imagined that the change was real: They trufted to appearances, without giving themselves the trouble of reflecting on the improbability of the fact, They who imagined that a piece of rotten wood or putrid flesh could become the eyes, limbs, and body of an infect of fuch delicate organization, and confifting of mufcles, nerves, veins, and arteries, could have but little difficulty in admitting, that the flesh of a chrysalis might be transformed into the wings of a butterfly, or that the fixteen limbs of a filk worm might furnish six for a moth.

After true philosophy appeared, one of the first lessons she gave her votaries, was to beware of trusting too implicitly to appearances, and of admitting ideas that were neither

meither clear nor intelligible: It was then that men began to recognife, that fudden metamorphofes were none of the expedients which nature employs for the production of her offspring. This point was fuccessfully laboured by Malphigi and Swammerdam, who carefully examined those insects that appear under different forms. By diffeeling them, a short time before the period of their transformation, they observed that their first form was owing to a covering under which their different members were to acquire their proper fize and firmnels: That all the parts of a butterfly, for example, were perfectly distinct under the skin of the worm which covered them; and that under the crustaceous shell of the chryfalis, they were fill acquiring greater degrees of strength; and were fast approaching to that state in which they were destined to appear, when the animal should arrive at perfection, and be able to propagate its kind.

From the experiments of these naturalists it appeared, that all the parts of the most perfect winged infect were distinctly formed, and gradually acquired fize and strength, under the disserent forms of a worm and a chrysalis, and that its progress and growth proceeded rather by developments, than by a real change: All the marvellous ideas conveyed by the terms transformation and metamorphoses thus vanished; and a beautiful analogy was established between the growth of all organised beings, whether in the animal or vegetable kingdoms.

A filk worm, or the worm of a butterfly, which is about to enter into its chryfalis state, is observed for some time before to grow languid, and to cease from gnawing those plants of which it was formerly so voracious: After having retired into a place sit for its purpose, and un-

Vol. III. X x dergoing

dergoing a few convultive struggles, the skin which covered it, and gave it the form of a worm, bursts, and the animal within makes its appearance; at first it is fost and tender, and covered with a viscous sluid which ascends from the body, but which afterwards hardens into that crustaceous shell in which all the members are again locked up, till they acquire greater firmness and stability. This vifeous fluid, which is generally feen coloured and opaque in its crustaceous state, is at first transparent, and through it the wings, limbs, and antennæ of the butterfly, are clearly perceptible. M. de Reaumure collefted feveral hundreds of these worms before their transformation, and placed them together upon a table, where he had many opportunities of examining them as they passed from the one state to the other: It was then that he diffinally perceived all the different members of the butterfly, before the chryfalis had assumed its hard and apparently inanimate state *.

After having remained for some time in this torpid state of a nymphe, chrysalis, or pupa, the limbs of the animal acquire sufficient strength to perform their sunctions; and it employs them in breaking open the second prison. On this event all the members are set at liberty, and instantly assume that posture and arrangement which is most suited to the new functions with which they are now to be charged: By these different processes does the butters sy arrive at that state in which it possesses all the faculties which are enjoyed by the most persect of its kind.

All infects do not undergo the fame number of transformations before they arrive at the winged state.

Some

^{*} Vide Memoires pour Servir a l'hist. des insectes, tom. i. p. 354.

Some, immediately on leaving the egg, assume a form pretty nearly refembling that which they possess after their growth is completed. The whole order of aptera, which comprehends all the different kinds of spiders, come under this denomination: The viviparous infects, in like mannner, appear at first, under their most perfect form. Some kinds, without undergoing all the changes of the filk worm, or that of the common butterfly, after having grown for a certain period, deposite a covering in which their wings were inveloped, and afcend into the air. Previous to this change, however, these enjoyed the power of locomotion by means of their limbs, and in that state were distinguished by voracity and activity: Of this description are all the different species of locusts. The flies, wasps, and bees, constitute another class, which, after leaving their vermicular form, and after passing into their chrysalis state, display their limbs and wings without being capable of using them. The last and most complete example of transformation is displayed by the class of moths and butterflies. On leaving the egg, and even before it, they assume the form of worms, which they again change for that of the nymphe, aurelia, or chryfalis: and it is not till they have lived a confiderable time under this form, that they throw off a second covering, and come forth winged infects.

One of the most wonderful circumstances in the economy of infects, is the different preparations which they make, and the expedients to which they have recourse for their preservation in their aurelia state. Many dig a hole in the earth, where they remain during the whole period of their inactivity. This is the invention of all the colcopterous infects, or such as have crustaceous wings. The

gnats, on the other hand, go into water, where they remain till the period of their winged state arrives. Some eat their way into seeds and fruits, where they undergo the different changes previous to their appearance on wing. Many lodge themselves in animal bodies on the approach of their transformation; several of the aquatic tribes bury themselves among sand, encrusted with a glutinous substance; while the numerous race of phalenæ wrap themselves up in the leaves of trees, the bombyces, or larger kinds, constituting for themselves a silken web, to protect them during that trying vicishtude of their lives.

In general, all infects provide for their fecurity before their helpless state arrive, by retiring from their usual haunts into fome sheltered retreat. It is thus that the worm of the butterfly provides for its fafety, by betaking itself, while it has yet the power of motion, to the hole of a wall, or the eave of a house: There some are fuspended by a thread, which nature affifts them in providing; fome hang by the head, others by the opposite extremity, and many by the middle. The crustaceous covering with which they are then clothed, affords another instance of the attention paid by nature to the preservation of her off pring, during a period when they are not able to avoid external injury by flight, Thus protected by the munificence of providence, myriads of animals fink annually into a flate of torpor fo profound as appears to threaten the extinction of every vital power. At the return of Spring, however, all nature feems again, to quicken into life; her fervants awaken from their torpid flate, and enter upon their functions with enlarged powers.

Sect. VI.—Of the Habitation and Food of Infects, and their Uses in the Economy of Nature.

It has been afferted by Ariftotle, that every kind of quadruped and bird was inhabited by its peculiar infect *; and this affertion, which has never been contradicted, feems to admit of being much extended. We have the ready feen, in a former part of this work, that the falmon and cod were at certain feafons infested with infects; the fame is probably the case with many other species of the finny tribe, whose history is less completely known. Some particular animals afford food and refidence to different kinds of infects; and the fame is the cafe with plants. As one animal is often feen to feed upon and inhabit a great many different plants; fo the fame plant often supports a variety of infects. The oak, in this country, affords fustenance to twenty different species of these animals; and there are varieties of this tree in warmer climates, that ferve for food to a far greater number.

Plants afford the most general and copious pabulum for this tribe of the animal kingdom. Wherever any infect is found indigenous in a country, there will always be found in it plants accommodated to its wants. There are many infects attached uniformly to one plant; the filk worm always gives a decided preference to the mul-

berry,

berry, but will live and propagate, though less vigorously, upon the common lettuce.

There are other inflances of the attachment of infects to particular plants, equally conflant with that of the filk worm; and by means of these, the animal is often known from the place where it was found to reside. Hence Linnaus has frequently given them names from the plants upon which they feed; a method often fallacious; since, perhaps, the greater number reside indiscriminately upon several plants, and support themselves upon a variety of different vegetable food. Some are afferted to be capable of residing in the human intestines, and undergoing their transformations there; being ejected from the stomach in their winged state.

The most possessons herbs afford food to infects equally with those that are falubrious: The water-hemlock, which is considered as the strongest vegetable posses, is much frequented by many of those animals, and seems to constitute their favourite nourishment.

In the different stages of their lives, insects are distinguished by various degrees of voracity; many of them in their larva state are most insatiable: The different species of the buttersy and silk worms are then endowed with teeth, with which they make great havock among leaves, even though of a pretty strong consistency; their stomachs in that state being capable of dissolving these harder substances. The same animal, when a chrysalis, loses all appetite for food, and those instruments that were employed in comminuting it. The teeth are deposited with its sirst covering; and the inner coats of the stomach are voided, it is faid, along with the excrements, a short while before the first transformation *. After being liberated from their

^{*} Memoir pour servir l'hist. des insectes par M. de Reaumur.

their last state of confinement, the butterslies are far inferior in voracity and in the powers of digestion; their food being then a thin liquid substance collected from the leaves of plants, and devoured only in small quantity.

The same is the case with the different species of locufts; fome of which in their larva state are the most voracious of all animals, and defolate entire provinces. It is not till they arrive at their winged and more perfect form, that their depredations cease, and mankind are relieved from one of the heaviest calamities which fall upon the human race in the fultry climates. It is by infects in their larva state that the roots of corn are perforated and devoured in more northerly countries. If the feafon prove cold and wet, they continue long under ground in that state of voracious vermine; and the crop, in the mean time, is fo completely ate away, that in fame inflances, scarcely a tenth stalk survives their depredations. In a more genial spring, these animals continue for a shorter space in their aurelia state; and the damage done by them is proportionably less; their destructive operations below ceasing as soon as they are enabled by their wings to rife into the air, and go in quest of fustenance in another element.

Infects, even in their winged state, all take food in a greater or lesser quantity; it seems, however, to be of a more desicate nature, being often only the liquid substances that exude from vegetables. Some are said, indeed, to be so short-lived, that they do not require any sustenance, and have, as it is afferted, no mouths: Of this kind are the sphemera, the may-sly, and the gad-sly. This sact, however, would require a more accurate degree of examination, and more satisfactory proof than

has ever yet been adduced, to establish it. The smallest and most delicate insects, for ought our senses can discover, may find various particles of matter floating in the air that may serve for their support; the estimates in a state of putresation, probably support many of this class. We know at least, that several sithes are capable of living upon the earthy particles that are found sloating in the purest water; and if animals of that superior size are thus supported, there can hardly be any room to doubt that insects, many of whom are searcely perceptible, may find abundant subsistence in the air, impregnated as it is with various essential.

The economical uses of insects, and their noxious esfects upon the various objects of human industry, have alteady been confidered *; it now remains, that we point out the purposes which they serve in the general system of nature. A class of beings by far the most numerous upon the sace of the globe, and however small taken individually, yet forming collectively a mass of organised matter, superior in bulk to any other department of the animal kingdom, every person will admit, could not be originally formed for no purpose, nor continue to be propagated without answering some important uses in the economy of nature.

The first and most obvious use of this part of the animal kingdom, seems to consist in that large supply of provision which it assorts the superior ranks of animated beings. It is for this important purpose that the whole surface of the earth is annually covered with plants and herbage; and many, perhaps the greater part of the larger animals are supported by the immediate pro-

-duce

duce of the vegetable kingdom; not a few, it must be allowed, are also sustained by devouring animal food. Between these two species of nourishment there is a wide difference; and infects afford a species of nutriment which feems to partake fomewhat of the nature of both, and which supplies the wants of an infinite number of creatures whose constitutions are not wholly adapted to either. Many kinds of birds live upon hardly any other food. What a blank in the feathered race would enfue, were this copious fource of provision that up from that part of the animal kingdom! The fithes feem still more dependent on the supplies afforded by those numerous tribes of infects that either float upon the water, or are feen hovering over its furface. The whale, the largest of nature's animated offspring, as we have already had occasion to observe, is supported solely by an infect which it finds floating upon the waves. Among this class alfo, an inconceivable diminution of numbers would necessiaily enfue, were the food of infects denied to the inhabitants of the water. Farther, many of the larger infects prey upon the fmaller; all thefe, as well as the different animals they support, must be unavoidably struck out of the family of nature the moment this species of support is withheld.

By means of the food of infects, therefore, a large proportion of the fuperior ranks of animated beings are fupported, all which would of necessity perish, were this order of animals destroyed. What a dreadful chase in the works of nature would the annihilation of a class of beings occasion, which of its is constitutes to large a proportion of her living productions, and which preserves the existence of so many more! The air, the earth, and the fea, which, according to the present system, teem with Vol. III.

life, would thus be more than half depopulated; and that great vivifying principle by which nature is actuated, and by which life is multiplied and carried on in all the clements without a paufe, would then be exerted in a great measure in vain.

But befides the fupply of food which infects afford to the fuperior animals, and their proving the grand inftruments in the hand of nature by which the supports life turoughout her dominions, they have always been deemed ferviceable in the general fyflem, by preferving the falabrity of the air. Over the whole furface of the earth, thole numberless productions that enjoy either animal er vegetable life, are continually falling into decay, and making room for that fuccession of organized beings with which nature conflantly teems: Hence, it has been supposed, that the atmosphere would foon become unfit for the support of life, did not millions of infects continually confume the carrior, and other substances in a flate of putrefaction, and purge the air of the noxious effluyia emitted from them. It is probably the office of those small insects who escape our observation, to destroy those noxious particles with which that element is impregnated, and which at certain feafons render it reflilential. The operation of this class of animals upon putrid fubstances, is much more confiderable than a superficial examination might fuggest. It has been afferted by the melt judicious and differning naturalifis, that the produce of a dozen of flies will confume a dead carcafe in a thorter space than a hungry lion. If this be true, what beneficial effects may be produced by that inconceivable number of infects, which in warm countries continually fwarm in the air?

CHAPTER II.

ORDER I .- Coleopterous Infects.

This order comprehends all those infects which have their wings covered by crustaceous elytra, and is divided into three sections, distinguishable by their antenna: The first includes such as have their antenna in the form of a club, thickest at the farther extremity; the second comprehends those whose antenna are sitiorm, or of equal thickness throughout; and the third is reserved for those infects that have cetaceous antenna, tapering from the base to their extreme points. There are thirty genera of coleopterous infects enumerated by Linneus, which include under them about eight hundred different species.

Y y 2

^{*} Vide System. Naturæ, p. 541, et seq.

GENUS I .- Scarabæus. The Bectle.

FORMERLY, the name fearabass was applied by naturalists to all infects whose wings are covered by a crustaceous elytra; Linnews, however, has discriminated these animals, and confined that term to this single genus. The tribe of fearabase are characterised by the antennæ, which terminate in a club, and are divided longitudinally into different plates or lawring. The second joints of the foremost pair of legs, in most subjects, are dentated *.

The larve or caterpillars of the greater part of this genus lead a fedentary life under ground; refiding most frequently in earth, while in a state of fermentation by a large quantity of manure. Pure dung is the favourite food of many; and it is in its immediate vicinity that they chiefly delight to dwell. The cockchafer, the most common of all the beetle tribe, is produced from the ova of the female deposited under ground, where it remains during the whole of its larva flate, devouring the roots of plants. In this destructive occupation the garden beetle, and all the hairy fearabai are employed, till the period of their emancipation arrive, when they take wing, and, forfaking the roots, betake themselves to the leaves of plants +. No calculation can afcertain the mifchiefs produced by these animals, in counteracting the labours .


^{*} Syft. Nat. ubi fupra. † Barbut's Genera Infectorum, p. 14.

Plate I.

ORDER 1.







Jahours of the husbandman: The more industrious he has been, and the richer his fields are, the more is he exposed to their unwelcome visitations. Those which have obtained the name of pilularii delight in the filthiest matter; and they have obtained that appellation from those round balls of excrement which they form, in order to deposit their egge. Happily for the scanty productions of these northern climates, only a small proportion of these nau-feous and destructive insects are produced there: sewer, perhaps, are found in Britain than in the corresponding latitudes upon the continent.

Of the fearabes, properly so called, there are three diffinct families: In the first, the thorax is armed with horns; the second has horns upon the head, and is unarmed in the thorax; in the third, both the head and thorax are without horns. Some of the infect, belonging to each of these families are featellati, or furnished with that part called the escutcheon, while others belonging to each of them are without it. Our enumeration shall comprehend a species belonging to each of these families.

The Bull-comber *.

I'ms species is not very frequent in Britain; and is of a very singular conformation. The body is broad and short; the clytra have longitudinal streaks that disappear gradually

[&]quot; Scarabæus Typhæus, Lin. Syft.

gradually as they proceed along its fides; the head has jects forward, and the antennæ are very apparent. The colour of this infect is black, except the inferior parts of the body, which are partially covered with a few vibrille of a brown colour. What particularly diffinguished the typhreus is the peculiar form of the thorax; the two lateral points of which project beyond the head, having a fmall protuber mee upon the fide, whill the middle point is thorter and formwhat raifed. These long projecting horns feem given to the infect as offensive wearons, although it is incapable of using them: It is not always that the members of this tribe of beings are applied to the purposes that feem most obvious and natural to the observer.

The refemblance which these thoracic horns bear to the long spears carried by the soldiers of the Macadomian phalanx, has occasioned the name of phalangill, which the French naturalists apply to this insect. The larve of the typheus is most commonly found in cow dung, where the grown animal is also frequently to be seen, both for the purpose of collecting food and depositing its eggs.

The Dung-beetle *.

This search one is remarkable for choosing the most file thy residence, heaps of dung and excrementitions matter. There

^{*} Scarabæus pilularis, Lin. Syft.

There are feveral other species nearly refembling it, in the impurity of its habits, as well as its external form, viz. the blue fearabeus, with the elytra and buck smooth, and the head armed with a rhomboidal helmet prominent at the top †: The fearabeus of an azure colour on the head and thorax, black legs, and pale coloured elytra ‡: The fearabeus of an oval figure, and black colour; the body smooth, and the elytra striated: And the scarabeus, whose head and thorax are black and without hairs, the legs pale ||.

These and many other pilular beetles are enumerated in the sauna swecica of Linnous; their size and form are not very different, while they are characterised by the same manners: They prowl continually amid every kind of silth, some of them having the same nauseous smell with the impure substances which they devour.

The dung-beetle, which is here felected for description, is smooth and of a black colour above, some mixed with green; on the lower side there are a few straggling vibrisse. The head resembles a hood, raised in the middle, and projecting at the edges: The jaws extend beyond the head. The thorax has a groove in the middle; its circumference is marginated, and its figure is round and smooth. The elytra are ornamented with a great number of longitudinal streaks: underneath the colours are various and brilliant, consisting of different shades of blue and green. There is observable on the anterior thighs, a spot formed by some red hairs, which, however, in some subjects is wanting. The tars appear weak

[†] Systeme Naturale du Regne Animal, Ord. i. gen. 1. spec. 14. ‡ Idem ibidem, spec. 15. || Rai, Insecta, p. 166.

weak and flender in proportion to the thighs, and this is uniformly the case with all the feet *.

The Golden beetle +.

THE last species of the beetle is deemed the most beautiful of the English insects. The whole body is of a glossy green, tinged with yellow; underneath, these colours partake of a shade of red, which some naturalists have compared to leather, others to finely polished copper ‡. In the brilliancy of its hues it rivals the emerald, and has sometimes been distinguished by that name.

The fearabæus in its vermicular flate remains on the furface of the ground, or covered to a little depth with the mould; there it preys on the roots of tender plants. After it is transformed into a winged infect it chiefly delights in the flowers of the rose and piony: It is consequently an inhabitant of the gardens, where its transfent beauty is reckoned a small compensation for the mischief of which it is guilty §. In the impure abode of the dung-beetle, that animal may be less pleasing to the spectator; but in return it is productive of less harm, and is less exposed to injury in a dwelling, of which it remains the unenvied possessor.

^{*} Barbut. Genera Infector, p. 13.

[†] Scarabæus auralus, Lin. Syft.

Vide Regne Animale, Tom. ii. p. 8. § Idem ubi fupra.

GENUS II .- Lucanus.

The animals of this tribe are the largest British infects; They are distinguished by the antennæ, which terminate in a knob, flattened on one side, and divided into a number of pectinated laminx. The jaws are extended beyond the head, incurvated, and beset with teeth *.

The lucani feed upon the liquor that exudes from oaks, which they fuck with their tongues. The females deposite their eggs in the trunks of decayed trees, such as the oak and the ash. The larvæ or grubs lodge under the bark or in the hollow of old trees, which they gnaw and reduce into a sine powder, and there transform themselves into chrysalides. They are found all over the ideal of Great Britain, but most frequently in Kent and Saffert. With the noxious effects of these animals on growing timber we are not acquainted; but notwithstanding their enormous size, it is probable that they are far less dearestive than those that prey upon the roots of corn.

* Syft. Nat. p. 559.

† Barbut Gen Infect. p. 17.

Vol. III.

The Stag-beetle *.

This beetle is readily diftinguished by its superior magnitude, which entitles it to the first rank among the infect tribes: It is however characterised by another peculiarity no less singular, and that is the large moveable maxillar, resembling in form the horns of a stag. These instruments are broad and slat, projecting from the head nearly one third of the animal's length. They have in the middler towards the inner part, a small branch, and at their extremity are forked. Their similarity in shape to the horns of the animal above mentioned, has struck every naturalist, and has, with equal propriety, procured for the insect the appellation of the slying stagt.


An infect of fuch an uncommon fize as the flag-beetle, and of fuch fingular conformation, has attracted the notice of all the entymologists. The female is diffinguishby the fize of her horns, which are not above half the fize of those of the male; both, however, are armed in the anterior side with small teeth, throughout the whole of their length; and both are sometimes as red as coral, which gives these animals a very beautiful appearance.

The head that supports these romantic horns, is broad, short, and irregular; and the thorax, which intervenes between

^{*} Lucanus Cervus, Lin. Syft.

[†] Cervus volans. Vide Mouffet, p. 148. Aldrovand, insect. p. 1513. Emperati, p. 169, and Charleton, p. 46.

Plate II.





between it and the body, is narrower than either, and marginated around. The elytra of this beetle are very plain, being unadorned with either streaks or lines. The colour of the whole animal is uniformly of a deep brown.

The residence of these animals is commonly the oak; in some parts of the country they are but rarely to be met with; and though the largest of all the coseopterous in this part of the world, they are much smaller than those of the same species, in countries where woods are more extensive, and the climate is warmer. In these they acquire an amazing strength and vigour; and the maxillæ, whose ordinary office is to tear the bark of trees, are occasionally converted into offensive weapons, which are carefully avoided by such as have experienced the severity of their bite.

The Parallelipiped Beetle *,

The body of this species is black; the horns smaller than those of the preceding, which in other respects it nearly resembles, and the body is oblong, and of that shape expressed by the Linnæan name, which we have borrowed †. It frequents the meadow grounds, and its

Lucanus parallelipipedus, Lin. Syst Sestabæus platyceros, Rzi.

habits are therefore probably different from those of the stag; but very little relating to its economy is known.

Linneus enumerates five other species of the lucano viz. the capreolus, the tridentatus, the interruptus, the carinatus, and the caraboides: Of these the hidentatus is distinguished by three teeth, with which the thorax is armed on each side. The maxillæ, like those of the last species, are in the form of a crescent, prominent, and indented on the inner side. It is a Swedish insect, not much inferior in size to those already described, and bears a strong resemblance to them in many particulars of its economy.

GENUS III .- Dermestes.

WE are now come to a tribe of beetles, much inferior in fize to those already mentioned; but of superior beauty. Many of the infects belonging to this genus exhibit a variety of the richest colouring that flows even from nature's pencil. They are near neighbours of man, and often troublesome companions: One species of the lardagius is destructive to meat, and is very difficult to prevent from entering into the repositories of the cook. It is a fill more unwelcome intruder into the cabinets of the curious; being very destructive to birds, infects, and other subjects of natural history, in a state of preservation. Arfenic is the most certain preventative against its depredations there. Many species of this family, as well as their larvæ, inhabit dried ikins, the bark of trees, rotten wood, feeds, flowers, and the carcafes of dead animals.

The antennæ of the infects of this tribe principally exhibit their generic characters. They terminate in a perfoliated club of an oval form, and divided into different plates or leaves, which feem to be united together by a small stalk. The thorax is of a convex form, and slightly margined: the head is bent inward, and, as it were concealed in the thorax *.

^{*} Systema Nat. p. 561.

The Domestic Dermestes *.

The form of this infect is oblong, and almost cylindrical: The elytra are striated; the thorax thick and gibbous. There are many varieties in the species, differing considerably both in size and colour; some being found of a dark brown, and others of a much lighter hue. When touched, it is struck with such an apprehension of danger, that it instantly draws back its head under the thorax, and its feet under the abdomen, remaining metionless in that position till the danger is over †.

Of all infects this is the most destructive to wooden furniture, where it is found in vast numbers, their larva state perforating it into those small round holes that appear on the external surface, while the inner parts are reduced to powder. There are few kinds of wood so hard as to be proof against the attacks of these animals, after they have remained long enough out to be drained of their natural juices. It does not seem to be ascertained how the eggs of this animal came at first to be deposited in timber: it would seem, that the winged animal which produces them, has the power of perforating cells for their reception; but it is evident, that the young in their larva state have a much greater degree of voracity, than the perfect animals, which are seldom found devour-

ing

Defmestes domesticus, Lin. Syft.

ing furniture, in comparison with those worms which no invention has been able to destroy. The acrid varnish, which is now extracted in such abundance from coal, promises to be the most effectual remedy against their energoachments*.

Dermeftes Lardarius +.

This infect is of a black colour, and easily diftinguished by a large transverse stripe covering the anterior part of the elytra, which is of a cinercous hue. The stripe receives this pale colour from a number of small gray hairs which grow upon it; it is irregular at the edges, and intersected through the middle by a small transverse streak of black spots, three in number, on each side of the elytra; the middle spot somewhat lower than the rest, gives the black streak a serpentine or undulating form ‡.

This animal feeds upon cadaverous carcases, and meat verging to a state of putresaction: It makes its way into the museum, and is seen among preparations, in numbers more than sufficient to gratify the curiosity of the possessor. In sact, it is among the greatest enemies of his labours: and in the larva state, makes the greatest havock among the collections of insects §. It then assumes

^{*} Vide Chap. i. fect. r.

[†] Scarabé Dissequeur poir, Regne Anim.

F Barbut gen. insectorum. § Regne Animale, Ord. 1. gen. 2.

fumes the form of a hairy oblong worm, which is divided and into fegments alternately of a dark and light colons. In that form it is often found deeply penetrating into old bacon, which it renders useless.

The infects of this genus are extremely numerous; Limeus has enumerated thirty species, and is probably far from having completed the catalogue. They are of sizes and colours too various even for the pencil to describe: In their manners, however, there is a strong refemblance; most of them inhabiting dung or putrid sless. The beautiful tints displayed by some, overcome our aversion to their impure habits: The violet dermestes exhibits the most brilliant shades of that colour. The thorax is covered with greenish hairs; the legs black, forming upon the whole a pleasing object, if we could forget that its residence, both in the larva and perfect state, it in the bodies of dead animals.

* Syst. Nat. Ord. r. gen. iii. spec. 13.

GENUS IV .- Ptinus.

This genus is characterised by its siliform antennæ, the extreme articulations of which are longer than those nearer the animal. The thorax is without a margin, rounded, and assords a receptacle for the head, into which the insect frequently draws it *. There are probably many varieties of this tribe which, from their extreme minuteness, escape observation: Linnæus has enumerated six different species, whose habits nearly resemble those animals last described. They attack household surniture, cloths, surs, and particularly dried animals in a state of preservation. Some of them when caught, have the artistice to counterfeit death. They draw in the head and limbs, and remain till the danger is over in a state of inaction, from which nothing but the application of heat can rouse them.

In order to deposite their ova they retire among hay, dried leaves, and other substances of a similar nature. from these retreats their larvæ issue, and penetrate rotten wood, and decayed household furniture +.

The prinus prefinicornis is the first species enumerated in the system of nature; it is produced from a small worm that ledges in the rotten parts of the bank of trees, where it makes a deep hole. The worm is there trans-

[?] Idem, gen. 4.

[†] Barbut's Gen, Infect. p. 23.

formed into a winged infect, which has obtained its name from the form of the antennæ, being pectinated on one hae. The elytra and thorax are of a deep elay-coloured brown; the antennæ and legs are of a paler shade of the same colour.

Genus V .- Hifter.

THERE are fix species enumerated under this genus, all varying in size, but uniformly of a dark colour: in some the elytra are striated, in others spotted; while some are smooth and uniform in their colour. The larvæ, as well as the winged insects of this genus, reside most frequently in the dung of horses and cows, and often upon sand. The general characters in which they all resemble each other, are drawn from the shape of the antennæ; the last articulation of which is larger than the others, and terminates in a solid knob; while the first articulation is compressed and incurvated. The mouth is forcipated; the head drawn back within the body; and the fore-legs are dentated *.

The hister unicolor is one of the most beautiful insects belonging to this tribe: The body is black, polished, and brilliant. The figure of the animal is almost square; the thorax large, and highly polished, having a flight margin that bounds its circumference. The anterior part is formed with a slope, into which the head is withdrawn, so completely, that it is only perceptible by the projection of the maxillæ. From this situation of the head, the insects of this genus often seem altogether deprived of that part. The elytra end abruptly, as if cut away by the middle, before they reach the whole length of the abdomen: They are smooth, having only a few strue on the outer side, that are scarcely perceptible †.

^{*} Syftema Nat. ord. 1. gen. 5. † Barbut Genera Infestorum.

GENUS VI .- Gerinus.

There are only two species belonging to this family of insects; the Natator and the Americanus; both easily diftinguished by their form, as well as their manners. The antenne are clavated, shiff, and shorter than the head: The hinder legs are also short, slat, and very broad. These animals are said to have four eyes*, two on the upper and two on the under side of the head; they appear in sact, on both sides of the head, but their number is only two:

The gerinus natator has an English name, being known in this country by that of the water-flea. It is that small animal which is seen describing circles on the surface of the water, by running on it with great swiftness. When an attempt is made to take it, it plunges below and eludes the grasp. The larvee are said to be found together with those of the elytiscus, an animal considerably larger.

The colour of the water-flea is a resplendent black, with a shade of brown: The elytra are adorned with beautiful strice, consisting of a number of points so minute, as hardly to be observed without the assistance of a microscope. On the hinder part of the margins of the elytra are seen, by the same means, small protube-

rances

rances, born upon pedicles, and so deciduous, that the smallest friction sweeps them away *. The feet of this infect are of a yellowish brown, and the hinder pair so short, that in certain positions the animal seems to have but two.

* Barbut Genera Infectorum,

GENUS VII. - Byrrhus.

This tribe when in its larva flate, refembles the worms of the dermefles; like them also they are extremely voracious, and are but too well known to those who are employed in collecting the subjects of natural history. Their antennæ, when they arrive at the winged state, are clavated, terminating in a capitulum or knob of a solid substance and oval shape.

The byrrhus verbasei of Linneus is of this genus, whose elytra are so thickly set with small scales, that the black colour of them is no where to be seen. These scales were of a white colour in the subject we examined, and so arranged, that they formed three undulating stripes across the elytra; between these stripes there were others of a similar form, but of a reddish brown. The infect is sometimes to be found stripped of this scaly covering, which so dissignres it, that it can hardly be recognised for the same animal.

GENUS VIII .- Silpha.

The infects belonging to this class are very voracious of carrion, upon which they deposite their ova in vast numbers. Thirty-sive different species of this fertile race have already been discovered; and it is probable that many have hitherto escaped all notice from naturalists. The antenne grow gradually thick towards their extremeties; the elytra are marginated; the head is prominent; and the thorax slattened, and surcounded with a border.

The different species of filphæ vary but little from each other, either in form or colour: In Britain, many of them are feen early in the fpring, under the loofe bark of trees. The filpha vespillo is one of the most remarkable of this genus, from which it differs confiderably in the form of its antennæ. They have at their extremity a reddish knob, formed by four small plates firung through the middle, one upon the other; the last being thicker, forms a fmail sharp pointed knob. The head. thorax, and body are black, charged with a few hairs of a yellowish hue. The thorax is of a rounded form, marked by feveral protuberances, and has the circumference terminated by a broad, flat, margin. The elytra are short, and as it were, cut across, leaving about one third of the body uncovered. Their colour is black, and variegated with two yellow transverse stripes.

Genus IX .- Cassida.

In this tribe the antennæ are nearly filiform, thickening a little towards their extremities: The elytra have a broad margin, and the head is entirely concealed under the thorax, which is flat, and forms a kind of shield to it*; a peculiarity from whence this genus takes its name.

The casside that are sound in Britain have, in their larva state, two prongs projecting from the extremity of the abdomen, with which they form a kind of umbrella of their own excrements, to shelter them from the sun and rain. When the umbrella becomes unsit for that purpose, it exchanges it for another, which is sabilicated of the same materials. These insects, which in their larva state are distinguished by manners as disgusting as those of the Hottentots, are afterwards transformed into the most elegant of the beetle tribe. Before they undergo their last change, they cast the skin several times; their food all the while being different kinds of thissless, and verticillated plants.

There are thirty-one species of the casside already deferibed; of these the green oval shaped cassida ‡ is the most remarkable for its beauty. Above, the clytra are wholly of a green colour, variegated with small strice, and projecting from the body like the shell of a tortoise; underneath, the body is entirely black, and supported by pale coloured seet: It inhabits the gardens and fields.

^{*} Syft. Nat. p. 574. † Reaumur, Tome III. Mem. vii.

[‡] Rai Inf. p. 107.

GENUS X.—Coccinella.

THE coccinellæ are generally didinguished by subclavated antenna; the palpi are also club-formed, the last articulation being shaped somewhat like a heart. The body is hemispherical, the thorax and elytra margined *. They are the most brilliant insects of this order; some searlet, others yellow, pale green; and these splendid colours are finely variegated with spots.

When the females of this tribe have been impregnated, they depolite their eggs upon the leaves of trees; and from them there are produced larvæ that are great devourers of plant lice. Their chryfalid flate continues for about a fortnight, when they are feen attached to a leaf by the hinder part; their bodies bent and fwollen. Their wings, after burfling the covering, foon become hard, and change from their pale colour to the red or fearlet, according to the particular species. Their infects are hardly ever capable of hight; their motions appear tather like jumping than flying.

The lady-bird of the *longlish* is a beautiful species belonging to the tribe of coccinellæ; the elytra are red, bordering upon yellow, and adorned with two black spots, one on the middle of each. The colour of the under parts is black. This species is found on the leaves

Vol. III. 3 B

^{*} Systema Nat: p. 579.

of trees; and in its larva state is a devourer of plant lice *. The different species of these animals are known principally by their colour, and the number of spots with which their wings are adorned. Linneus has enumerated no less than forty-nine kinds.

^{*} Vide Rai Inf. p. 86.

GENUS XI .- Chryfomela.

From the variety of different colours with which the infects of this tribe are adorned, they have been called chrysomele. There are no lefs than an hundred and twenty-two different species described by naturalists, diffinguishable by their size, colour, and manners. They all agree, however, in having the antenne moniliform, and increasing in thickness towards the ends: Neither the thorax nor elytra are margined *.

The chrysomelæ are to be found almost every where; some frequent the open fields; others the woods, and many the gardens. Their larva prey upon the leaves of trees, rejecting the fibrous parts; some of them dettroy the grass while in that state; and there is in Sweden a particular species, said to be guilty of very extensive depredations on the pasture grounds. This insect, when it has reached its winged state, is of a sine glossy green, mixed with a shade of blue; the whole differentiated with a number of small spots, which form irregular strice to

Chrysomela fanguinolenta, has the elytra of a deep black, interspersed with small spots, which gives them the appearance of shagreen: Their margins, and the wings which they cover, are marked all along with a broad band of light red, which resembles streams of blood; the belly, thorax, and head, are of a blue celour.

3 B 2

Chryfomela

f Şyft. Nat. p. 586.

Chrysomela cruciata †. This singular species has the elytra yellow, and marked with two black lines, resembling a cross; the head of a dark blue, the thorax brown, narrow, and cylindrical. This species is frequent at Hamburg, where it devours the asparagus, and is supposed by the gardeners to have been transported from Russia along with that plant ‡.

† Frisch, p. 27. Regne Animale, p. 26.

GENUS XII .- Hifpa.

ONLY two species of these insects are sound in Europe; two others are enumerated by Linneus, which are not indigenous in this quarter of the globe. They have the antennæ formed like spindles, growing gradually thicker from each extremity to the middle : their infertion is between the eyes; the thorax and elytra of this tribe form their most distinguishing character; both being covered with protuberances like spines *.

The hispa atra is of a deep black; the upper part of the body entirely covered with firong spines, somewhat like the shell of a chesnut; the elytra are also set thick with finall vibrifcæ, which makes this very fmall infect prove a hedgehog in miniature j.

* Syft. Nat. p. 603. † Barbut, p. 437


GENUS XIII .- Bruchis,

This family of infects is superior in fize to the last, and more diversified: It contains seven species, having club-formed antennee; four palpi placed at the extremity of a short rostrum *.

The bruchus of the pca blossom frequents that flower; is covered with ash-coloured down, forming cloudy spots upon the thorax and elytra; the latter are so short, that they leave a great part of the abdomen exposed, and thickly covered with longitudinal streaks.

^{*} Systema Nat. p. 603.





GENUS XIV .- Curculio:

THESE infects are in some parts called weefils: Near an hundred different kinds are enumerated in the System of Nature, a fifth part of which are sound in Scotland. The antenne are subclavated, and situated in a projected rostrum, of a horny substance.

As fome of the birds who are supported by the fruits of man's industry repay their benefactors with a song, so the infects of this genus make some compensation for the ravages they commit, by the enchanting richness and variety of their colours. Upon them nature seems to have profusely lavished the most refulgent and vivid tints which dazzle the eye. Often, however, the curculiones peculiar to our country are so diminutive in size, that we are only admitted to survey their beauties by means of the microscope.

In their larva state, the animals of this genus are neither so inessensive nor so beautiful, as to entitle them to the savour of man. They penetrate into the grains of corn, while yet of small size, and in proportion as they grow, they increase the boundaries of their habitation, by eating out the heart of the grain. As the number of these insects is great, the mischief they occasion in granaties and corn losts is often very considerable. After lying for some time, the grain which they frequent is entirely consumed to the huses; and however sound it may externally

ternally appear, it is no fooner expected to the wind, than it blows before it like chaff.

It is with the hulks of grain, after they have ate the fubfiance, that there animals are transformed into chryfalids; and when ready to come forth winged, they perforate their manifon, to make room for their cfcapes. Beans, peafe, oats, and other kinds of grain, afford food and a retreat to this tribe; many of which also perforate plants, and dwell in the interior parts; some of them mine into the leaves of trees, devouring the parenchyma that lies between the outward pellicules.

The green curculio. This species is of a bluiss green, thining with a fine resplandent shade of gold, like the rock of a pigeon: The head, thorax, addomen, and feet, are all of this beautiful colour; the antennæ are black, having the last articulation longer than the rea.

^{*} Pelivert: Gaza; p. 77.

GENUS XV .- Attelabus.

This tribe very much refembles the last, but is inferior in fize and beauty: The animals which compose it have the head inclined, and gradually tapering towards the thorax; the antennæ increase in thickness as they approach the extremity*.

Attelabus curculionoides has obtained its name from its resembling the curculio; the antennæ are short, fixed on the point of the head, which is of a triangular shape, having the sharp angle joined to the thorax; the head, part of the thorax, and belly, are black; the rest of the insect is of a fine red.

3 C

Vor. III.

* Syftema Nat. p. 619.

GENUS XVI .- Cerambyx. The Capricorn.

The tribe of capricorn beetles owe their original to larvæ refembling foft flender worms, whose heads are scaly, and who are provided with fix hard legs. These larvæ, which produce a race of the most beautiful insects, are in general white, and are found in the inner part of trees, which they perforate for a double purpose, that of obtaining food and a retreat, while transformed into chrysalids. It is from these cavities that the winged capricorn is seen issuing, as soon as it has completed its last change; and in this act it is easily caught. Many of these insects emit a strong smell, which is selt at a considerable distance; and when laid hold of, they utter a cry, occasioned, as is supposed, by the friction of the thorax and abdomen.

Cerambyx viridi-cærulescens*. This species is green, with a mixture of azure blue; the antennæ are as long as the body; each side of the thorax is furnished with a sharp protuberance; the wings are black, and the feet of a resplendent blue. It frequents the leaves of the willow, and, when approached, emits an agreeable slavour, like that of the rose †.

Fauna Swecica & Lister, p. 384. † Regne Animale, p. 35.

GENUS XVII .- Leptura. The Wood Beetle.

The antennæ of these insects are setaceous; the clytra diminish in breadth towards their extremity; and the thorax is round and slender. The larvæ of the wood beetles are found, like those of the last genus, in the perforated trunks of trees, and, like them, they devour the pulverised wood when digging their retreat.

Leptura arcuata. This is one of the most beautiful of infects; when viewed with a microscope, it appears like velvet inlaid with precious stones; and if exposed to the rays of the sun, it shines with infinite splendour. The ground colour of this animal is a velvety black; the elytra are variegated with transverse bars of a bright slame colour, formed by streaks of down, of a refulgent yellow *.

3 C 2

Barbut, p. 57.

GENUS XVIII .- Necydalis. The Carrion Eater.

Or this genus there are eleven species, which have obtained their English name from their supposed predifection for carrion. In their winged state, they generally frequent the woods; their residence and habits, while larvæ, are wholly unknown. The generic characters of these insects are, the setaceous form of the antennæ, and the shape of the elytra, which are either shorter or narrower than the abdomen †.

The first infect of this family enumerated by the Suchest naturalist is the larger carrion eater, which is distinguished by its large and prominent eyes: The antenne are raised upright as far as the first articulation, after which they are bent, and turned aside. In some individuals, the thorax is black, in others yellow; the elytic are generally black, lighter towards the middle, which contains a lemon-coloured spot.

[†] Systema Nat. p. 641.

GENUS XIX .- Lampyris. The Fire Fly.

THESE infects are characterifed by filiform antennæ, flexible elytra, a round flattened thorax, flurrounding and concealing the head; the females are, in most species, without wings †. It is the female of one species of these infects that is in this country termed the glow-worm, from that phosphorescent light which it emits during night. Two or three of these animals inclosed in a glass vase, will give a light sufficient to enable a person to read in the darkest night.

This fingular phenomenon is observed most frequently in the month of June, when the animal is in motion. The female can withdraw or display this light at pleafure, by contracting or unfolding her body; her purpose in shewing it is said to be in order to attract the male. When crushed with the hand, this luminous substance of the glow-worm adheres to it, and continues to shine till it is dried up.

The noctiluca, or glow-worm, in its infect form, poffesselytra, and wing: under them, longer than the body. The head and antenne are black; the former entirely concealed by the broad plate of the thorax. The four last rings of the abdomen, which emit the light, are in the male not so bright as those of the female, and are nearly destitute of that luminating quality which renders her so remarkable †.

GENUS XX .- Cantharis.

THE infects of this family are numerous, and diftinguished into twenty-seven different kinds. From the similarity of the name, some have supposed them the same with the cantharides imported from Spain, whose virtues are so well known in medicine; that infect, however, belongs to a different genus. The cantharis frequents flowers; its larva resembles that of the cerambyx already described.

All the infects of this tribe have fetaceous antennæ, the elytra flexible, and the thorax margined, and fhorter than the head. Like the preceding genus, they have papillæ upon the fides of the abdomen.

Cantharides ænes-viridis †. This infect is of a kind of brass green; on each side of the thorax is a small nipple, of a sasson-colour, with three tips. The thorax and elytra are of a deep green, the latter terminating in a red spot. Beneath, the insect is of the colour of brass, and red under the wings ‡.

† Rai Inf. p. 101.

† Regne Animale, p. 69.

GENUS XXI .- Elater. The Skipper:

The infects of this genus are diftinguished by setaceous antennæ, but more particularly by an elastic spine which springs from the under side of the thorax, near the extremity. By means of this spring, these animals, when turned upon the back, are capable of jumping into the air, and recovering their position.

In the state of larvæ, the insects of this genus inhabit the trunks of decayed trees, and are there metamorphosed into the winged form. Their residence is then changed, and they are seen in various haunts, slowers, thickets, or open fields.

The chesnut-coloured elater †. This insect is found both in the corn and pasture fields. The antennæ are branchy, and the tips of the elytra black; the rest of a pale slessh-colour; the thorax is covered with a sine ash-coloured down ‡.

† Lister Loqu. p. 387. ‡ Regne Animale, p. 64.

....

GENUS XXII .- Cicindela. The Sparkler.

The porrected jaws afford an eafy mark to discriminate the infects of this genus; they are both armed with teeth; the eyes of these animals form another striking peculiarity, being singularly prominent; the thorax is round, and margined †.

Cicindela campestris. This insect is one of the most beautiful which this country assords. The whole upper parts of the body are green, tinged with blue; underneath is a mixture of yellow, red, and copper. The elytra are delicately marked, each with small white spots. The insect, and its larvæ, are found in fandy dry situations; the latter resemble soft whitish worms, and immerge themselves into a perpendicular hole in the sand, with the head at the bottom, ready to catch the insects that sall into it. These animals are all rapacious, devouring whatever they can overcome ‡.

GENUS XXIII .- Buprestis. The Cow-burner.

These animals, both in their external figure and manners, nearly refemble the skippers already described; their colours, however, are far more resplendent; and, when viewed with a microscope, their essugence dazzles the eye. From this circumstance, they have obtained in France, where they are more numerous than in this country, the name of Richards.

The characters of this genus are, fetaceous antennæ; the head drawn back within the thorax; the mouth armed with jaws, and furnished with palpi.

Buprestis guttata. This elegant species is found on fandy ground, below timber that has begun to rot. Its head enters within the thorax, and the elytra are surrowed, and adorned each with sour or sive spots of a bright yellow; the upper side of the abdomen is blue *. The larvæ of this and the other species belonging to this tribe are unknown.

Vol. III.

3 D

Regne Animale, p. 58.

GENUS XXIV .- Dytiscus. The Diver.

This genus comprehends twenty-three species of aquatic infects, mostly of a very large size; they are distinguished by the antenne, which are either setaceous, or terminated by a persoliated knob; the hind legs are armed with claws, and furnished with vibrish, that enable the infect to swim *.

The divers have obtained their name from the fudden escape they make from danger, by plunging into water. By day, they frequent the flagnated pools, from whence they iffue in the evening, to make excursions on the wing. The males have furrowed elytra, while those of the females are plain. The latter, when impregnated, deposits her eggs in the water, wrapped up in a filky cod, of a texture fufficiently strong to protect them from predatory infects, especially those of their own kind, which are mutually rapacious. Their larvæ are worms confifting of eleven fegments, distinguishable by the largeness of their heads, and by four filiform antennæ; they deyour other water infects, and even each other. When arrived at their winged state, their bodies are protected by a hard fealy coat of mail: If caught unguardedly, they are in that flate capable of giving a fevere bite, as well as of wounding the hand with a sharp spine.

Dyticus

^{*} Systema Nat. p. 664.

Dyticus piceus. This is one of the largest of the tribe; all over of shining black; the antennæ are of the shape of clubs, and perfoliated. Their colour, and that of the sour palpi, is brown; the head of this insect is slattened, and surnished with a pair of formidable jaws.

GENUS XXV .- Carabus. The Ground Beetle.

In this family, which contains no less than forty-three species, are found some of the largest of the British infects. They are characterised by setaceous antennæ, and by the shape of the thorax, which resembles a heart; the point cut off and margined. The elytra are likewise surrounded with a margin *.


In their winged state, the heads of these animals are prominent; their mouths, like those of the preceding genus, are armed with jaws, and sour palps. Their eggs are deposited under ground, or in decayed trees, where the larvæ reside till they are metamorphosed. It is during their caterpillar state, that they are prejudicial to gardens, and hence have been ironically called gardeners by the French †. It is not, however, vegetable productions alone that they devour: They are the greatest tyrants to other insects, and destroy indiscriminately, as many as their strength enables them to overcome.

Carabus violaceus ‡. This species is of an oblong thape; the colour a dark violet: The edges of the thorax and elytra, are violet with a shade of purple. The latter have neither dots nor streaks, but are marked with strong longitudinal wrinkles. It is found among rotten wood.

^{*} Systema Nat. p. 668.

[†] Regne Animale, p. 44,

[‡] Rai Inf. p. 96. n. z.





GENUS XXVI .- Tenebrio. The Darkling.

These infects are commonly of a dark fombre colour: Some have wings under their elytra, while others are apterous; and from this circumftance, they have been divided by naturalists into two fections, containing together thirty-three different species. The larvæ of some species relide among rotten sticks, or rubbish; those of others take up their abode among flour, and different kinds of food. After they have become perfect infects, they enter houses and devour every fort of provision. Their principal refort is to damp cellars, where putrid air and darkness attract them. It is from their precipitately avoiding light, as well as their gloomy appearance, that they derive their generic name.

Tenebrio Mortifager*. This fable animal is found among rubbifh and dung in gardens; when it enters houses, as is frequently the case, the credulous believe it the certain foreignner of death. It is entirely of a deep black, without the least lustre: The antennæ are long; the thorax margined; and the elytra wrinkled. This, as well as some other species, is remarkable for emitting a fetid smell.

Tenebrio primus Voyage d'Alande. † Charleton exercit, p. 48.

GENUS XXVII .- Meloe. The Bloffom Eater.

In the first edition of the System of Nature, there were only one species of this genus taken notice of; in the later improvements of that celebrated work, there are sixteen kinds of these animals enumerated; while, perhaps, a still greater number remains to engage the refearches of suture historians. This singular tribe, contains in it the genuine cantharis, or blistering say, and is distinguished by having moniliform antennæ, with the last articulation of an oval shape. In all these insects the head is insected and gibbous; the thorax round; and the elytra slexible *.

The infects of this genus are either winged or apterous; in the former, the elytra are fhort, extending about half the length of the abdomen: In the latter, these parts reach the whole length of the body, and conceal under them two slender wings: In all of them the antennæ resemble a necklace, composed of a number of small rings; those of the middle larger than those near either extremity.

Meloe profearabæus †. This infect is foft to the touch, and of a deep black, with a flight shade of purple towards the under part of the body. The feet and the antennæ

are

^{*} Systema Nat. p. 679.

are of a violet hue. The elytra are coriaccous, and feem as if cut through obliquely from the inner to the exterior edges, being thicker at their future than at the fides. Some naturalists have termed this species the unctuous cantharis, from that fat, oily, and fragrant substance, which exsudes from the body *. This oleagineus matter enters into the composition of salve for plague-fores; and is deemed an excellent antidote against the poison of the scorpion's shing.

Meloe vesicatorius. This is the species so frequently used in pharmacy; an innect to which man is perhaps more deeply indebted, than to any individual belonging to this class of beings. The blistering cantharis is about nine lines in length; the colour a resulgent green, mixed with azure. In the southern parts of Europe it multiplies exceedingly; some of the provinces of Spain onnually receive a large sum for those they export to the rest of Europe. They are there seen slying in vast swarms, and alighting upon trees and shrubs, whose leaves they devour. They are faid to prefer the ath leaf to that of any tree in the forest; but whatever leaves they devour, they are uniformly accompanied with a heavy nauseous smell, like that of mice, and thence their haunts are discovered by those who go in quest of them.

In their humid and living flate, the odour exhaled from these infects is so corrolive and irritating, that the gathering them is attended with danger. In that occupation, the labourers, who imprudently collect them in the heat of the day, and with their hands uncovered, are frequently seized with a violent heat of urine, and voiding of blood. The same accidents beial those who unwarily sleep under the trees they frequent.

The female cantharidis feems to feel the access of amorous desire in a more violent degree than the male: It is she that courts the male; and in the great act of fecundation, it is she that occupies that place, to which in most animals nature directs the other sex. After impregnation, she deposits her eggs in the ground, where they remain till they have undergone the various changes that are to bring them forth winged cantharides.

When collected and dried, these insects become so light, that sifty of them hardly weigh a dram: it is in that state they are grinded down into the well known powder, which constitutes the basis of the common blistering plaister. Of the other purposes to which they have been applied, ignorance is perhaps better than information; and we freely resign to the annalists of dissipation, the task of recording those vain attempts in which they have been employed by the enervated debaucher to restore he virility. The enterprises of love, tike the fatigues of war, require certain intervals of rest and tranquillity, without which neither the lover nor the soldier can take the field without hazarding his reputation.

Militat omnis amans, et habet sua castra cupido, Ovip.

GENUS XXVIII .- Mordella. The Nibbler.

This genus has the antennæ filiform, and ferrated; the head bent under the neck; the palpi clubbed, and obliquely truncated. The elytra are curvated, or floping down towards the tail; and a broad lamina is feated at the base of the abdomen before the thighs*. There are only fix species of the mordella known to naturalists; of these the mordella aculeata is quite black and smooth. The abdomen terminates in a sharp point, and extends a considerable leagth beyond the tips of the elytra.

The yellow mordella is of a fmall fize, and of a dark shade of that colour. The elytra bend downwards at the extremity, where they assume a ferruginous colour: The point of the abdomen, which extends beyond them, is of an obscure black.

Vol. III. 3 E

* Systema Natura, p. 682.

GENUS XXIX .- Staphylinus.

The antennæ of this fingular tribe are moniliform: The elytra, by which they are readily diffinguished, feem as if cut through by the middle, and do not extend half the length of the abdomen. Under these cases the wings are folded up, and concealed. The extremity of the abdomen is unarmed, but provided with two oblong vesicles which the infest can shoot out or retract, at pleasure.

The staphylini are sometimes seen upon slowers; but their principal resort is to the dung of cows. All of them when touched turn up the tail, as if with an intention to sting. They are, however, unprovided with that instrument, but bite severely with their jaws when laid hold of. In their larva state, these insects so much resemble the perfect animal, that they cannot easily be distinguished.*

Staphylinus maxillofus, is the largest British infect of this genus. The jaws, from which it obtains its Linneau name, are sharp, hard, and rather longer than the head. The whole infect, both above and below, is of a deep black: The elytra do not cover more than one third of the abdomen, which appears beyond them, rough and hairy. The legs are long; each of the tarsi is provided with a tust of hair resembling a brush. This and the other species can sly by means of large wings, which unfold from under the elytra †.

^{*} Barbut, p. 95.

[†] Rai Inf. p. 109. n. T.

GENUS XXX .- Forficula. The Earwig.

This genus contains fome species of infects that are universally known. Their common characters are setaceous antennæ; elytra shorter than the abdomen; wings solded up and covered by the elytra; and a forceps, by which the extremity is armed *.

Forficula auricularia. This infect is feen every where; it even enters our immost apartments. The body is of a yellow colour, inclining to brown; dark above, and lighter underneath. The forceps with which the earwig is provided, procured the generic name forficula to this tribe. The formidable name of earwig has arisen from a notion, that these intects entered into the ears of people while asleep, and from thence penetrated into the brain, where they occasioned much pain; often madness and death: The French appellation, which sign see the Ear-piercer, urges this accusation in still plainer terms.

The smallest acquaintance with the structure of the human ear, will easily prove the fallacy of these suspicions, which have in all probability often cost this animal its life. It, however, destroys slowers; and wherever fruit has been wounded by stronger insects, the earwig generally attends for a second feast.

All these animals after secundation lay eggs, from which caterpillars are produced, that differ but little 3 E 2 from

Syftema Nat . p. 686.

from the perfect infect. It is amazing in how little room the long wings of these animals are packed before they burst from their chrysalid state; and even after they are protruded, the animal, by means of the joints and mutcles with which they are furnished, folds them up under the elytra, that you would imagine it altogether deprived of these instruments *.

The life of the carwig, after it enters upon its winged state, lasts but a few days; each fex is then constantly in pursuit of the other, and that short existence feems chiefly employed in the business of procreation: After having provided for posterity, the parent animal dries up, and dies to all appearance consumptive †.

^{*} Swammerdam, p. 114.

[†] Goldsmith, p. 358,

CHAPTER III.

SECTION I.

ORDER II .- Hemiptera.

WE are now come to the order of hemipterous infects, which comprehends the different kinds of locusts and grashoppers, and, in general, all those animals whose antennæ are setaceous, and heads inflected. Their elytra and wings are extended, smooth, and semicoriaceous; being of a substance somewhat like vellum. The thorax is horizontally slattened, of an orbicular form, and marginated round the extremities: The seet are formed, in most species*, for running; and the abdomen is terminated by two small appendices resembling horns. They are termed beniptera, because only half of their wings appear when in a state of rest: They do not then meet together in a longitudinal suture like those of the last order, but have some part of their interior margins crossed, or laid one over the other, above the abdomen.

^{*} Pedes curforii, Lin. Syft.

GENUS I .- Blatta.

This order of infects has been fubdivided into eleven different genera; the first of these is the blatta, or cockronch, which comprehends that well known species of animals which frequent kitchens and bakehouses. Their appearance is ugly and deformed; they seldom, however, present themselves by day; and though in our immediate vicinity, and of considerable size, they are but rarely seen. If surprised in their hauats while it is light, they speedily escape by running; their wings being unsit for slight, except in the males of some species who make aukward attempts at slying. The night is their season of activity, when they issue from the crevices near the chimney in quest of crumbs of bread or dough.

In this island the cockroach is probably always a domestic animal: In warmer countries, however, there are different kinds which haunt the fields in great numbers. The hakkerlac of the American isles, that voracious animal, which so greedily devours the provisions of the inhabitants, is of this genus. That infect, like our cockroach, shuns the light of day; during which it lies concealed in some hole, till darkness approach, to favour those depredations for which it has long been so infamous in the New World.

The larvæ of all the different species of blattæ differs but little from the perfect insect, excepting in the total want of the wings and elytra. In that state, meal is its

common

common food; and of that necessary of life it is then extremely voracious. When meal is wanting, as must happen in the fields, it corrodes the roots of plants. After having attained its full complement of limbs, and acquired its most perfect form, the cockroach displays a mouth armed with jaws, and furnished with palpi. Its antenne are generally as long as the body, and the abdomen as broad as the thorax. The tarsi of the fore feet have sive joints; those of the hinder only four; while the horns which terminate the abdomen are furrowed, or wrinkled transversely. There are ten species of this genus, named, for the most part, after the different countries which they inhabit.

^{*} Barbut, Order II. Gen. S.

GENUS II .- Mantis, or Soothfayer.

THE infects of this genus possess a form the most roa mantic and extraordinary that is perhaps prefented by any animated being; and fo powerfully have their fingular attitudes operated on the minds of the credulous and ignorant, that superstition has invested them with certain powers that are altogether unexemplified by any part of the history of animated nature. By the fingular manner in which the foothfayer stretches out its fore legs, it has acquired the reputation of a diviner, who could unfold all the fecrets in the bosom of futurity; and because the infect often fits upon its four hind legs, having the two fore ones raifed up and folded together, the believing multitude have supposed it to be then holding intercourse with the Supreme Power, in the exercise of devotion *; a circumstance from which it has obtained from the peafants of Languedoc the name of Pregadiou, or the God-prayer. It is in that province, where these animals abound, that the country people have also ascribed to the foothfayer another very commendable quality; that of obligingly shewing the way to strangers: This it is supposed to do, by that peculiar habit which it has of ftretching its fore legs fometimes to the right, and fometimes to its left fide. These superstitions of the vulgar have been as favourable to the security of these animals, as they are disgraceful to human reason; for they have procured them a character almost

[·] Vide Goedart.

almost facred; and the injuring them is reckoned a crime of no small enormity +.

The infects of this genus are diffinguished by an unfleady nodding motion of the head, which gives it the appearance of being flightly attached to the thorax. The mouth is furnished with jaws and pulpi; and the antenne are setaceous. The four wings are membranous, and wrapped close round the body. The anterior pair of feet are compressed laterally, and servated on one side, their extremities terminating in a single nail. The four back feet are smooth and gressory, being formed rather for advancing slowly than for performing quick movements. The thorax is throughout of an equal size, narrow, and in some species extending to a disproportionate length ‡.

The most remarkable of these animals is the mantis googylodes of China. The thorax is uncommonly long and narrow; the head small and slat, with two short siliform ansennæ; behind these, two large polished eyes are placed; the rostrum has the shape of an awl, but often split towards the extremity into two points. The elytra, which cover two thirds of the body of the insect, are reticulated, and crossed the one over the other; the wings which they cover are veined and diaphanous. The four hind legs have the appearance of being winged, on account of those large membranous lobes which emerge from their joints. The anterior pair are armed with spines at their first articulation, and towards their extremities they are ferrated on one side of the same street in the same street are street in the same street are street and towards their extremities they are ferrated on one side of the same street in the same street are street and same street are same street.

Vol. III.

SF

[†] Reamur, Tom. IV. ‡ Systema Nat. Ord. II.

T Vide Barbut's Gen. Infect. Ord. II. Gen. 2.

GENUS IV .- Gryllus.


This defiructive genus comprehends all the different kinds of locults, and is divided by linnaus into five tribes, containing fixty-one different kinds of animals. Their general characters are, a head inflected, armed with jaws, and furnished with palpi; the antenna in some species are setaceous; in others filiform: the wings are wrapped round the sides of the body; the under one folded, and concealed under the elytra. All the seet are armed with two nails; the hind pair formed for leaping.

The tribe of achetæ, which comprehends in it all the different species of crickets, is distinguished by the bristles which spring from the extremity of the abdomen, and by the three articulations of which their tars are composed. This samily have obtained their English name from the continual tiresome noise which they produce. The domestic achetæ usually take up their abode in overs, and in the holes of walls around kitchen chimneys, where they are attracted by the heat, and are very noisy companions to the country people, who, from some prejudice in their savour, seldom wish to destroy them.

The field cricket remains, during the day, pent up in some subterraneous habitation, from whence it iffues forth about sunset; when, in countries where these animals abound, the whole fields ring with their noise.

The mole cricket is by far the most hideous and extraordinary infect belonging to this tribe. It has obtained

Plate V..





its name * from the fingular structure of its fore legs, which are extremely broad and flat, and terminate in fix large serrated claws, somewhat resembling the fore feet of a mole. The whole animal is of a brown, dusky colour, very large, and active. It frequently takes up its residence in hot-beds, to the great disturbance of the gardener; for it digs under ground, like the animal after which it is called, committing dreadful havoc among the tender roots of the plants that are artificially raised there. It is frequent in France, where the gardeners know it by the name of courtilliere.

The tettigonia, or tribe of grashoppers, is the next section into which the grylli are divided; and of it, the gryllus viridishmus is the most remarkable. Its colour is a pale green, the antennæ setaceous, and longer than the body. The elytra are clouded, and the wings reticulated; both extend beyond the body about one third of its length. The semale carries, at the extremity of her abdomen, a kind of servated spine, composed of two laminæ, and in shape broad, and turned up like the blade of a cutlass. These implements are employed by the semale in digging in the ground, or in wood, holes for the reception of her ova; and this being a function in which the male has no share, he is unprovided with the instruments by which it is performed †.

The female grafhopper possesses an amazing fecundity; she regularly deposits from four to seven hundred eggs at a time ‡. The wonderful precautions which she takes for providing them security, and food for the young as

3 F 2 foon

^{*} Gryllo-talpa, Linnæus.

[†] Barbut's Gen Insect. page 115.

Memoir. pour servir a L'Hist. des Insect. Tome IV.

foon as they are disclosed, merit our particular notice. With that lancet, which we have already described, she excavates a number of holes in the dried branch of a tree; into each of these holes eight or ten of her eggs are dropped; there they are furrounded with that kind of food which is most suitable for them in their larvæ state. The disposition of the eggs is in rows, and placed in the middle of the trees; the soft substance of which is the sirst food of the insect after it leaves the ovum. The insect that proceeds from each of these eggs, after it has grown for some time, and before reaching a fize incompatible with escaping by the narrow mouth of the hole, takes a final departure from the place of its birth.

The larvæ having thus left their egg flate, and acquired the use of their limbs, the two anterior of which are formed for digging the ground, soon apply them to that purpose, and excavate for themselves a subterraneous retreat among the roots of plants, which they gnaw, and support themselves upon the juices that exsude from them*. In this state they remain till they are ready to undergo another transformation, which introduces them into the open air in the form of winged insects.

A fhort time after, the grathopper appears in its last stage of perfection; it spreads over the meadows, which it fills with its chirruping strains, which are the calls of the male inviting the semale to love. Some naturalists are of opinion that the notes of the grashopper are produced by rubbing the two hind legs of the animal against each other. M. de Reamur and Linnaus, who have minutely examined these insects, derive their vocal powers from a very different source. On examining the male,

his

his body has been found provided with a fmall hole below the infertion of each wing, delicately confiructed with organs of found within, and covered over externally with a fine transparent membrane. It is by means of these organs, which, in the completeness and delicacy of their structure, may vie with those of the human voice, that some species of the grashoppers produce their melody. The cicada of the ancients, so samous for beguiling the labours of the husbandman by his melodious notes, is an animal very different from our grashopper: The former either walked or slew; and it was from the summit of a tree that it poured forth those delicate notes, so much celebrated by the Grecian poet *.

The note of the grashopper is feldom heard, without being returned by another male of the same species; and the two little animals, after many mutual insults of this kind, are seen to meet and sight desperately. The semale is generally the reward of victory; for, after the combat, the male seizes her with his teeth behind the nock, and thus keeps her for several hours, till the business of secundation is performed: They are at that time so strongly united, that they can searcely be separated without tearing them as and errors.

After focundation by the male, and towards the close of fummer, the female is seen diffended with the feeds of a future family; and the prepares for depositing her numerous ova in the manner we have already mentioned. The eggs are white, of an oval shape, and horny consistency. In size, they are nearly equal to a grain of anise; and while in the body of the semale, they are inveloped within

^{*} Vide the Odes of Anacreon.

[†] Goldfmith's Nat. Hift. Vol. VII. p. 334.

within a covering, branched all over with veins and arteries. In this form they remain deposited under the furface of the earth, or inclosed in wood, apparently unaffected by the rigour of winter, till the genial heat of fpring begin to hatch and vivify them. Then, the fun beginning with its warmth to animate all nature, the infect eggs feel its benign influence; and generally about the beginning of May, each egg produces a larva about the fize of a flea, at first of a white colour, but afterwards gradually turning brown.

After having taken these measures for perpetuating her kind, the parent animal does not long survive; as the winter approaches, she dries up, seems to feel the effects of age, and dies from a total decay. Some affert that she is killed by the cold; others, that she is eaten by worms; but certain it is, that neither male nor semale are seen to survive the winter.

When examined internally, the grashopper discovers a very singular and complicated structure of viscera; besides the gullet, there is observed a small stomach; and behind that a very large one; still lower down, there is yet a third: So that it is not without some soundation, that all the animals of this tribe have been supposed to chew the cud, as they so much resemble ruminating animals in their internal conformation. Aristotle informs us, that they were greedily sought after as a delicate morfel by the Greeks; and that the season when they were deemed most delicious was a short time before they left their chrysalis state*. The metamorphoses from that state is performed with great difficulty and agitation; many perish in this severe effort of nature, and those who survive are for some time in a languid and debilitated state †.

[·] Historia Animalium.

[†] Reamur, Tom. IV. p. 188.

SECTION II.

THE locusts occupy the next section of this genus; & face of animals, unhappily for mankind, by far too well known by their destructive effects on the productions of the earth in many of the warmer regions of Afia and Africa. They are distinguished by a simple tail, and filiform antennæ*. The former of these members is destitute of those setaceous bristles which characterise the achetæ, and of the tube which is peculiar to the tettigoniæ already described. The locusts constitute a multifarious group of the most voracious and prolific insects which are visible to the naked eye. Some of them are of a large fize, others fmall; fome are green, others red or yellow. Among us they receive various denominations, according to their fize and colour: When of one form, they are called locuits; those of another, grashoppers; and a third fort are termed crickets +.

Many of the locuits far exceed the grashopper tribe in bulk, in rapidity of slight, and the powers of injuring mankind, by swarming upon the productions of the earth. The latter animals are unable to sly any considerable length; and the quantity of vegetable food which a few of them sporting among the grass can destroy, is scarcely perceptible; but when a swarm of locusts, several miles in length, and two or three yards deep, settle upon a seld, the consequences are truly alarming. The annals

of

of many countries in the East have recorded their devaflations, which at different periods have threatened the
extinction of the human race. Happily for the inhabitants of Europe, the coldness of their climate, and the humidity of their foil, are unfavourable to their production.
In some of the southern kingdoms, their visitations,
though far less frequent than formerly, are still an object
of terror: In general, however, their incursions are satal
to these animals; they visit us and perish.

The facred feriptures, which were written in a country where the fwarms of locults made a diffinguished feature in the picture of nature, have given us many firiking reprefentations of the numbers and rapacity of their infects: They compare, according to the manner of the oriental languages, an army of which the numbers are beyond computation, to a fwarm of locults: They defectibe them as rifing out of the earth, where they were produced; as purfuing a fettled march, for the purpose of destroying its fruits; and as co-operating with the divine indignation in punishing the fins of men.

When the locusts take the field, they have, as it is faid, a leader at their head, whose slight they observe, and pay a strict attention to all his motions. They appear at a distance as a black cloud, which, as it approaches, gathers upon the horizon, and almost hides the light of day. It often happens that the hostandman sees this imminent calamity pass away without doing him any nuschiles, and the whole swarm to proceed onward, to settle upon the labours of some less fortunate people: But wretched is the district upon which they alight. They ravage the meadow and the pasture ground, strip the trees of their leaves, and the gardens of every we yetable.

The vilitation of a few minutes destroys the hopes of a year; and a famine but too frequently ensues.

In their native tropical climates, the locules are faid not to be so dreadful as in the more southern parts of Europe, and the opposite coasts of the Mediterranean: There, though the plain and the forest be stripped of their verdure, the power of vegetation is so great, that an interval of three or sour days repairs the calamity. But our verdure is the livery of a season, and we must wait till the ensuing spring repair the damage; besides, in their long slight to this part of the world, they are samished by the tediousness of their journey, and are therefore more voracious wherever they happen to settle *.

But it is not by what they devour, that the localls do fo much damage, as by what they deftroy; their very bite is thought to contaminate the plant, and prevents its vegetation: to use the expression of the husbandmen, they burn whatever they touch, and leave the marks of their devolutions for two or three years. The bite of many animals operates like a poison to vegetables; and it this be the case, it does not appear that the depredations of their infects can, in any country, be soon repaired.

Such are the fatal effects of the voracity of the locusts while alive; and when dead they sometimes prove still more nomous, by infecting the air with a stench that is insupportable. Or of us tells us, that in the year of the world 3800, there was an incredible number of these infects which intested Africa; and after having eaten up every thing that was green, they slew off, and were Vol. III.

Goldsmith's Nat. Hist. Vol. 7th p. 345.

drowned in the African fea, where they caused such a stench that the putrifying bodies of many thousand men could not have produced essure for pestilential.

The eastern borders of the Russian empire, are subjected sometimes to the awful visitations of this insect: That which happened in 1690, which extended from Russia over a great part of Poland and Lithuania, was singularly destructive. In some places the locusts were seen lying dead, heaped upon each other sour feet deep; in others they covered the urface like a black cloth; the trees were seen bending beneath their weight, and the damage which the inhabitants sustained was beyond computation.

In Barbary their numbers are also formidable, and their visits more frequent: During the year 1714, a traveller from Britain, remarkable for the accuracy of his observations, witnessed the havoc they committed in that ill sated country. Towards the end of March they began to appear with a southerly wind: During the succeeding months, their numbers continued to increase so prodigiously, that during the heat of the day they rose in swarms so large as to darken the sun. In the middle of May they began to retire, for the purpose of depositing their eggs in the drier plains of the interior country.

About the middle of fummer the young, already ripe for devastation, made another incursion, in several bodies of a vast extent; although then in the form of worms they crawled forward, climbed the trees, walls and houses, devouring every plant in their way: It was in vain that the inhabitants dug trenches through their fields

and

[.] Dr. Shaw in his travels through Africa,

and filled them with water; in vain they collected large rows of heath, stubble, and other combustible matter to fet them on fire on the approach of the locusts. The trenches were soon filled, and the fires extinguished by the immense swarms that succeeded each other.

The locusts which are thus active in their larva state, remain only about a month in that form; after having completed their growth, they cast that skin which gave them their vermicular shape; and, in order to prepare themselves for this transformation, they attach their hind legs to some twig, where, after some laborious efforts, and several undulating motions, they at last burst the skin: at first, the head only appears, but soon after the rest of the body is disengaged, the whole operation continuing only for seven or eight minutes. After casting their covering they remain for a little in a languishing state, till the air hardens their wings, and the heat of the sun again invigorates them, when they resume their former voracious habits, with an increase both of strength and agility.*.

In fome parts of the world the inhabitants convert what so generally is considered as a plague, to an advantage, by making the locusts an article of food. It is for this purpose, that in many parts of the east they are caught in small nets, which are constructed for entangling them. When a sufficient quantity is thus procured, they are roassed over the sire in an earthen pan till the wings and legs drop off: when thus propared they are reckoned tolerable sood, and are said to taste like cray-sish t.

In Tonquin nets are used in dragging a species of them out of the rivers, into which they frequently fall from their unwieldiness

[&]quot; Dr. Shaw's travels.

⁺ Dampier's voyages.

unwieldings in the months of January and February. This locust, which is reckaned a great delicacy both by the rich and poor, is caten fresh after being broiled on the coals: It is of the fine of the top of the finger, and as long as the first joint; inhabits the low grounds, where it breeds, till the winter feafon, when it fpreads over the country in vall fwarms. In all countries where the locust is an article of food, it is regularly brought to market, like shell-fish or small birds in Europe. They must have conflituted a common food among the Jews, fince Melia, their lawgiver, has condescended to specify the different kinds which they were permitted to eat *.

From participating of this luxury of the Yews, however, the inhobitants of Britain may deem themselves hapby in being excluded. Since the year 1748, none of the destructive species have visited this island: At that peried, the great brown locust was feen in feveral parts of Encland, and many dreadful confequences were apprehended from its appearance. The body of this infect is about three inches long; the antennæ are extended about one third of this length from the head, which is brown. The shield which covers the back is greenish; the upper fide of the body is brown, fpotted with black, and the under fide is purple.

The great West Indian locust, of all the animals of this genus, is the most formidable, when individually considered. The body is about the thickness of a goofe quill, divided into ten annuli, and is fix inches in length: It has two small eyes standing out of the head, like those of

crabs

^{*} Vide Levit. chap. xi. v. 22. Even these thou mayest eat; the locust after his kind; the bald locust after his kind; the beetle after his kind; and the grashopper after his kind.

crabs; and two filiform antennes. The whole body is shaded with small excresences, which are not much larger than the points of pins: Its shape is roundish, diminishing in circumference towards the tail, which is bifurcated; between these forks, the animal is armed with a dangerous sting, which invariably wounds any person who touches it. The poisonous matter with which this instrument is loaded, occasions a shivering and palpitation all over the body, which the inhabitants cure by rubbing the wound with palm oil †.

Goldfmith's Nat. Hift. Vol. VII. p. 348.

SECTION III.

GENUS IV .- The Fulgora, or Lanthorn-fly-

The infects of this genus are rare in Britain, only two species being caught in the island. It is the foreign animals of this tribe which have the singular quality of emitting stashes of light when they sty. This phenomenon is thought to be occasioned by the waving of the elytra, whose thinness renders the spots upon them transparent: The effect is probably heightened by some resplendent quality peculiar to the tribe, and by the golden yellow of the under wings bordered with black. However this may be, it is certain that in China there is a certain species of the sulgora, which blazes with an amazing lustre, that marks the path of the animal wherever it directs its course, and which in the minds of the timid and superstitious, creates images of danger of the most tremenduous kind.

Linnaus enumerates nine different species, one only of which he has termed European from its residence in that quarter of the globe. They are all characterised by a long projection from the forepart of the head, of a light and empty substance. The antennæ are seated below the eyes, having two articulations, the exterior of which is largest, and of a globular form; the rostrum is bended inward

inward under the body; the feet are greffory, or formed for walking *. In these animals the elytra commonly decline along the sides, and when at rest cover the whole of the abdomen; for they gradually increase in breadth from the base to the extremities.

The fulgora candelaria of Linneus, is of a fresh green colour; on the ground of the elytra figured and striated with lines of a pale yellow; the wings are of a deep and brantiful yellow, with a band of glossy black, bordering their extremities; the head and thorax are generally of a ruddy brown, and the animal emits slashes of lightning as it slies in the dark.

^{* 3}yftema Natura, Ord. ii. Gen. 4.

GENUS V .- The Cicada, or Flea locust.

THE infects of this tribe, obtain fometimes the name of grashoppers, or froghoppers, as well as the tettigoniae already mentioned. They are far inferior in fize as well as in their destructive powers, being produced from small larvæ that are found on plants resembling the fix-legged worm. Some of these larvæ have the singular property of voiding from the anus and the cuticular powers, small bubbles which unite together, and form a species of soam in which the young animal remains enveloped.

This foam, which is not unlike a spittle, is in some parts called the cuckoo-spittle, and is probably intended by nature as a shelter to the larvæ, against the search of other animals to which it would fall a prey: Perhaps, too, this moisture may be designed for serening the infect, in its tender age, from the beams of the sun, and the rough elements by which its body might then be injured *.

Whatever purposes this froth may be intended to serve, it is undoubtedly necessary to the subdiffence of the animal †, for it is no sooner wiped away, than a fresh quantity is seen exsuding from all parts. During its residence in this spume, the animal acquires sour tubercles on its back, in which the wings are inclosed; as soon as

thefe

^{*} Barbut's Genera Insect. p. 122.

[.] De Geer artes de Stockholm, 1741. p. 221.

The

these burst, from a reptile it becomes a winged insect; thus rendered complete in all its parts, it slies to meet its mate, and propagate its kind *.

Other larvæ are unprovided by nature with a frothy covering; their bodies are not fo easily injured; and they are enabled to escape from their pursuers by the nimbleness of their running and leaping. The chrysalids and larvæ that produce them differ but little from each other, either in their form or habits: Both run, leap, and climb upon plants; the former distinguished by small knobs upon the back, the rudiments of future wings.

When arrived at their most perfect form, the males of many kinds possess the faculty of singing, by means of an organ situated under the abdomen, furnished with valves and muscles. It is by the help of this sonorous instrument that the male amorously tolicits his mate; and from this lively and animated song, the country people presage a sine summer and plentiful harvest;

The cicadæ have for their generic marks, the rostrum inslected; the antennæ setaceous; the sour wings membranous, and the seet formed, for the most part, for leaping ‡. Their heads are generally of a triangular shape; their bodies are oblong; and the wings sastigiated, like a roof. The semales possess, at the extremity of the abdomen, two large laminæ, between which is inclosed a kind of blade, serrated on the edges. This apparatus appears plainly intended for digging holes for the reception of the young, and for sinking them in those plants upon which the larvæ are to feed, after their exclusion from the eggs.

Vol. III. 3 H

^{*} Goldsmith's Nat. Hist. Vol. VII. p. 258.

[†] Barbut ubi supra. ‡ Syst. Nat. Ord. II. Gen. 5.

The genus of cicadæ is divided by Linnæus into five tribes or families, viz. into those whose thorax is compressed, membranous, and larger than the body; those which have their thorax armed on each side with a spine or horn; the third, the manniferæ, whose feet are not formed for leaping; the sourth consists of those whose hind feet are made for that purpose; the last family comprehends those whose wings are wrapped round the body *.

The cicada fanguinolenta of Linneus is the most beautiful of all our freghoppers: It is of a shining black, both above and below; the elytra have each three large spots of a fine crimson.

⁹ Systema Nat. ubi supra.

GENUS VI .- The Notonecta, or Boat-fly.

The flies of this genus have the roftrum bent in towards the thorax, the antennæ very flort, and the four wings croffed over each other *. The fix feet are all formed for fwimming; and the abdomen is terminated by four horny appendices. This tribe comprehends in it only three species; the grey, the striated, and the small notonesta.

The first is of a pale colour, mixed with black, and is very common upon stagnated water †; the head is round, and for the greatest part occupied by two large brown eyes; the antennæ are very small, of a yellow colour, and inserted in the under part of the head; the elytra are of a rusty cloudy colour, large, and crossed one over the other.

The fecond species also frequents the water, is much smaller than the former, and emits a disagreeable odour. The elytra are pale, striated with a number of undulating transverse lines ‡. Its shape is oblong; the forehead and feet of a golden yellow; the thorax brown.

The small notonecta is of a fize scarcely perceptible; and, according to the Swedish naturalist, has neither wings nor elytra. The whole insect does not appear larger than a small grain of sand, and is of a brown colour, transversely striated §.

3 H 2

The

^{*} Regne Animale, p. 100.

⁺ Vide Petivert, Gaz. t. 72.

[‡] Fauna Swecica, Ord. II. Gen. 3.

[§] Regne Animale, p. 102.

The notonectæ have obtained their name from the finagular manner in which they fwim on the back, prefenting the belly uppermoft. This fituation feems admirably fitted for the creature's manner of feeding, which is faid to be on the under fide of plants that grow on the furface of the water; the animal, by having its mouth turned upwards, is capable of taking its food with greater convenience: Nor are its motions in the least impeded by this aukward posture; on the contrary, it is very nimble, diving down inflantaneously, when it perceives danger, and rifing again to the furface, the two hind legs all the, while ferving for paddles *.

[&]quot; Barbut, page 129;

GENUS VII .- The Nepa, or Water Scorpion.

THE characteristics of these animals are drawn from their inflected rostrum, the chiliform * antennæ, and the four wings folded crosswife, the anterior part being coriaceous. In the System of Nature there are fix species enumerated, three of which are found in Britain. All thefe dwell in the water, both in the state of larva and chrysalids. The eggs of the water scorpion are also deposited in that element; they are of an oblong shape, and have at one extremity two or more vibrifcæ, the only part of them which is feen, the rest being funk and concealed in the stalk of a bulrush, or other water plant. These plants may be removed, and placed in water near the naturalit, who may thus have an opportunity of feeing them hatched and coming forth animated larvæ immediately under his eye. The water scorpions, when thus excluded from the eggs, remain for fome time in the state of worms, in the place of their nativity. When they arrive at their full fize, and are metamorphofed into complete intects, they are fometimes an inch in length, and nearly half as broad. The antennæ appear in the usual place of the fore legs +; they are armed with a forceps, well adapted to the rapacious purpofes to which they are applied. The nepæ are of all animals the most tyrannical;

^{*} i. e. Formed like a hook, or the claws of a crab.

[†] Vide Geoffroy, as quoted by Barbut, p. 131.

tyrannical; they deftroy, like wolves among fleep, twenty times as many as their hunger requires. If one of them is placed in a bason of water with thirty or forty worms of the libellula, each as large as itself, it will deftroy them all in a very short space, getting on their backs, and piercing them through the body with its rostrum *.

These animals, though they live upon the water by day, are capable of taking long slights from one pool to another in quest of food: This they are probably often obliged to attempt, from the sierceness of their manners, by which the infects in their vicinity must be soon destroyed. Though so formidable to other creatures, they are nevertheless haunted by a little louse, which probably repays the injuries which the water scerpion so frequently commits upon others.

The cincreous feorpion is the most common insect of this tribe: The head is small, and as it were sunk in a slope of the thorax; the elytra are broad, cross each other, and cover almost the whole abdomen: In the females, the extremity is furnished with two appendices, equal to three sourths of its length.

^{*} Goldfmith's Nat, Hift. Vol. VII. p. 360.

GENUS VIII .- Cimex. The Bug.

We are now to examine a tribe of infects equally remarkable for their hideous appearance and their voracious habits. Of this naufeous race Linneus reckons up no lefs than an hundred and twenty-one different kinds, all diffinguished by these general characters: an insected rostrum; antennæ longer than the thorax; wings solded together crosswise; the forepart of the upper ones coriaceous; the back is shat, the thorax margined; and the feet are formed for running *.

For the fake of a clearer arrangement, this numerous genus is distributed into eleven different sections or families; and among these the cimex lectuarius holds the sirst station. This troublesome inmate has attracted the notice of almost every naturalist; and it were well for the rest of mankind, that the knowledge of it had been consined to them. By day, it lurks, like a robber, in the most secret parts of the bed and apartment; takes advantage of every chink and cranny, to make a secure lodgement, and contrives its habitation with so much art, that scarce any industry can discover its retreat. So cautionsly does it avoid the light, that is candles, or a strong sire, be kept burning,

^{*} Syftema Nat. Ord. II. Gen. 8.

[†] Meuffet, p. 269. Bonani, Micro. p. 65. Rai Inf. p. 7. Aldrov. Inf. p. 534.

burning, it will feldom venture from its hiding place. As foon, however, as darkness promises security, it issues from every crevice, drops from the roof, or crawls from behind the arras; and unhappy is the patient to whom these creatures direct their course. There are some perfons whom they do not bite; but such as are selected for their repast are foon covered with wounds, which swell to a confiderable bulk, and are often painful. In vain the fufferer endeavours to relieve himself, or avenge his wrongs, by killing fome or driving others away: famished multitudes renew the attack, and occupy their place; while, in the meantime, the bed becomes insupportable by their naufeous fmell; nothing remains for him, but to make a timely retreat, otherwise he will be disfigured and fwolen with their wounds, and fpend in reftlefs agitation those hours in which the weary are detlined to rest, and even the wretched to forget their forrows.

The retreats of the house bug, in its wild thate in the sields, are unknown: About a century ago, it was scarcely seen in Britain; and happily for the inhabitants, it multiplies less in this country than almost any part of the continent. Throughout France, Italy, and Spain, the beds in most of the inns swarm with bugs; and every piece of surniture seems to afford them a retreat. They acquire a greater size in proportion as the climate is warmer; they are more active, and bite with a more cruel appetite. There the weary traveller, who is subject to be bitten, remains the whole night like a centural upon duty; and, instead of inviting the approaches of sleep, watches the attacks of innumerable invaders, who are ready to gorge themselves with his blood *.

3

The

The house bug differs from all the rest of the genus, in having neither wings nor elytra. It has two small brown eyes, and two antennæ, having each three articulations: Underneath these lies the crooked trunk, its instrument of torture, which, when the animal is in motion, lies inslected upon the breast. The parts of generation are obvious in both sexes: they are often seen coupling together in the act of procreation; two days after which, the semale deposits her eggs, to the number of an hundred and fifty.

Cleanliness is the best antidote against these animals, as their hostile attacks appear the proper punishment of its neglect. Many secret compositions are made use of to destroy them; but that object seems rather the effect of assiduity than of a cure. Mixtures have been made for this purpose, of soap, verdigrise, and Scotch souls, which are said to be effectual. After having taken down the bed three or sour different times, and washing it with a solution of corrosive sublimate, they have been banished. The smoke of peat, where that suel is to be had, is the most efficacious of all applications, and uniformly destroys these troublesome animals wherever it is applied.

The field bugs have all wings, and inhabit plants as various as their shape and colour. In their larva state they are very active, and differ but little from the perfect animal, except in their wanting wings: In this state, however, as well as that of a chrysalid, the animal is incapable of propagating its kind. After the last transformation, the wings unfold; and the impregnated semale lays her eggs commonly ranged one beside another, upon the leaf of a plant.

Vol. III. 3 I Thefe

These eggs, when viewed with a microscope, present singular varieties of configuration: Some are crowned with a row of small hairs; others wear a circular sillet; and the greater number have a cappice, which the larva pushes off when it bursts open the egg. Released by nature from their prisons, the young overspread those plants upon which they feed, extracting, by means of the rost-rum, those juices which are proper for their nourishment. Even in this state, the larvæ begin to renounce their gramenivorous and peaceable habits; some of them are voracious in an eminent degree, and spare no small animal that comes in their way *.

This feems indeed but a prelude to that ferocity which these creatures display in their perfect state: Then they destroy caterpillars, slies, and many of the coleopterous insects, which seem protected from their assaults by the hardness of their wings: They are then mere cannibals, who glut themselves with the blood of every animal, not excepting even those of their own genus.

The cimex hiosciami of Linnaus, or hen-bane bug, from its residence on the plant that bears that name, is described by Ray under the appellation of the smaller wood bug. Its body is of an oblong and narrow shape: Above, the insect is red, variegated with black spots.

Another species is also found on the hen-bane, with brown wings, decorated with white spots; it is described by Linneus in the Fauna Swecica; and is twice the fize

^{*} Barbut's Gen. Infect, Ord. II. Gen. 8. † Rai Infect. p. 55. † And his Voyage to Oelande, p. 155.

fize of the former, but refembles it nearly in shape, and the distribution of its colours. Of the various races of bugs, some are found in the meadows; some frequent slowers; and a great many reside among bushes and shrubs, which seem almost to defy the pencil to discriminate them, or imitate their various hues.

S I a

GENUS IX .- Aphis. The Plant Loufes

Few infects are more common than thefe, while few prefent greater fingularities to the observation of the naturalist. Plant lice are feen on the leaves of a great variety of plants, and often in fociety, and in confiderable numbers. At certain feafons, they are viviparous, and at others they produce inanimated eggs, which remain in that flate till the action of the fun call forth their vital powers. The foctus, when the parent is viviparous, thews figns of life before it is fairly excluded from the body. Sometimes the same mother gives birth to near twenty in a day, without appearing lefs in bulk than before *. If one of these pregnant semales are taken and pressed between the finger and the thumb, a still greater number of young will be forced from her body, one following another like a flying of beads, and growing fmalder and fmaller, in proportion as their period of natural delivery was more remote.

Reaumur is of opinion that fecundation is performed among the females of this tribe, without copulating with the male; and the experiments of fucceeding naturalists have confirmed an affertion that feems to be contradicted by the analogy of all animated nature. Take a plant louse, the inflant it issues from its mother; shut it up carefully from all access to congenerous animals; and if

it

^{*} Reaumur, Tom. III. p. 291.

A Memoir, pour servir a L'Hist des Insectes, Tom. III. mem. g.

it prove a female, it will produce young: In like manner, the young produced from this virgin loufe, kept feparately, will produce an offspring. Some naturalish have repeated this experiment to the third and fourth reproduction. Bonnet observed nine successive generations of this nature, all in the space of three months. The most probable account of this singular fact seems to be this; that as in all animals secundation is accomplished by the copulation of male and semale; so that happens with the plant lice; but with this difference, that the secundation produced by that act is transmitted for several generations, before the prolific virtue becomes entirely exhausted.

The young, immediately after being protruded from the mother, are always of a paler colour than the parent: They have then the entire use of their limbs, and go in quest of food. All the different kinds cast their skin; and it is after these developments that such of them as have wings obtain these parts.

It is not the males only that are winged, as we have feen in feme other genera; many of the plant lice of both fexes are deprived of these instruments in their most persect state; for many of the semales without them are seen producing young. As the larvæ of these insects possess all the activity peculiar to their genus in its last stage of growth, so they are equally distinguished by voracity. They are surnished with a small trunk, which pierces the leaves, and enables the animal to extract the juices proper for its nourithment. Many plants grow deformed by the number of punctures thus made upon their leaves, and decay from the want of their usual sapsome thrive even when covered with these insects, which on being

being broken open, disclose the numberless families that inhabit them +.

These galles or excrescences formed by the plant louse, and which often so much disfigure the shrubs upon which they grow, are in *China*, *Persia*, and the *Levant*, applied successfully in dying. Some species formed in these more northerly climes might probably be turned to the same useful purpose, were their virtues understood.

The plant lice are feen to void from the anus a clear liquid fubstance, of a faceharine nature, which is greedily fought after by other infects; the same kind of sugary liquid oozes from the extremities of two small horny appendices that terminate the abdomen. It is in pursuit of this sluid that the ants are seen to frequent those plants upon which the plant lice settle in greatest plenty. From this circumstance, the former animals have been supposed hostile to the latter; while other naturalists have deemed it an indication of the friendship of the two genera for each other: but it is neither from motives of love nor of hatred that the ants repair in such numbers to these haunts; they seek to gratify their avidity, by devouring their excrements.

The most effectual method of preventing the depredations of plant lice upon flowers and the leaves of plants, is suggested by those insects which live by preying upon them. The plant louse-lion, or aphidivorous sly, either by instinct or foresight, deposits her eggs in the midst of these animals; and as soon as the larvæ are produced, they devour hundreds around them, without the necessity of any other movement than turning to the right side or the left. This formidable insect is furnished with two tubercular

[†] Reaumur ubi supra;

unbercular horns, with which it pierces the plant loufe, and fucks its juice. A number of its eggs placed upon the leaves frequented by the lice, would foon produce a number fufficient to suppress or destroy them *.

The plant louse is a very minute insect. Its generic characters have been thus described by naturalists: The rostrum is bent inwards; the antennæ are longer than the thorax; there are four erect wings belonging to many genera, while others altogether want these parts; from the extremity of the abdomen there project two horny appendices; the feet have one articulation, and are formed for walking †.

M. de Reaumur supposes that almost every plant has its peculiar louse, the colour varying according to that of the leaves upon which they reside. Thirty-three species are reckoned up by the Swedish naturalist.

GENUS X .- Chermes.

NEARLY allied to the last tribe of infects, but of superior size, is the chermes, a race termed gall infects by M. de Reaumur*. Their characters are drawn from the situation of the rostrum, which is placed in the breast; and from the shape of the antennæ, which are longer than the thorax. The wings are four in number, folding close along the sides of the abdomen; the feet are formed for leaping, their tars having two articulations.

These animals are sound inhabiting a great variety of different trees and plants, upon which they produce very singular excrescences: The Linnean names assixed to each species are for the most part derived from the particular tree upon which they feed; that of the sig tree is the largest, and therefore more easily examined than any of the rest of the tribe.

The whole body of this infect is brown above, and green beneath; the antennæ are large, hairy, and of the fame hue with the back. The wings, which are twice the length of the abdomen, are placed fo as to form a kind of roof, as if to protect the animal from rain +. Other species, of inferior size, frequent the clm, the ash, the cherry, and the sir; that which inhabits the last of these trees is provided with a snarp pointed instrument,

by

by which it makes punctures in the extremity of the branches, in order to deposit its young. By this means the fir-tree chermes produces that enormous scaly protuberance, which is often seen at the summit of the branches, and which is formed by the extravalation of the juices through the wounds thus made †.

The larva chermes has fix feet: In figure, it refembles the perfect infect; its shape is oblong, and its motion flow. In the chrysalid state, the form is somewhat changed, by two small protuberances upon the thorax, the rudiments of future wings. When the chrysalids are about to undergo their last metamorphoses, they retreat to the under side of a leaf, to which they remain attached without motion.

On the approach of their change, the membrane above the head and thorax is feen to split and open: The perfect insect then comes forth with its wings, leaving the spoils of its chrysalis still adhering to the leaf, and rent on the anterior part. The empty sloughs of these insects are often found in great plenty beneath the leaves of the fig-tree.

The tubercles raifed upon the branches of trees by the punctures of the chermes, not only become the refidence of the animal, but also of its eggs and larva, which are contained in those cells with which they abound. The box-tree chermes produces no excrescences upon that plant: Its punctures make the leaves bend in towards each other at their extremity, where their union forms, at the summit of the branch, a hollow knob, in which the larvæ of that infect find shelter.

Vol. III.

3 K

Both

Both in their larva and chryfalid flate, many of the chermes eject from the anus a white faccharine fubflance refembling manna: within the hollow balls formed by the box leaves, there are fmall foft grains of this fubflance deposited; and in that flate it is frequently seen issuing from the anus of the insect that dwells there it.

† Barbut's Gen. Infect. p. 152.

GENUS II .- Coccus. The Cochineal.

THE infects of this tribe, present the most fingular form which we have, perhaps, yet surveyed in this department of animated nature. They were long imported into Europe as an article of commerce, before their claim to rank among animals was admitted.

The males are commonly of a smaller size than the fermales; the former have two erect wings, while the latter are apterous; the rostrum is situated in the breast; the antennæ are setaceous, and the poisers or halters are wanting; the posterior part of the abdomen is bristly, or velveted *.

Several species of these insects are found upon European plants; that, however, upon the pimpinella roots, called the grain of scarlet of Poland, was long known in this part of the world, as a dye, before the Mexican cochineal was introduced. A particular history of this insect, which is now no longer an article of commerce, since the discovery of Mexico, may be found in Breynius, as quoted by Reaumur †.

The drug at present most in estimation, for the fine colour it gives our clothes, is the cochineal of Mexico, an insest domesticated and reared with great care by the Indians. It grows not upon the root, but upon the 3 K 2 leaves

leaves of a plant known by the various names of oputatia, nopal, racket, and cardassia. The method of cultivating them upon this tree, practised by the Mexicans, is by collecting ten or twelve together into one nidus, made of moss or cotton; these insects being put into these mests, are fixed to the branches of opuntia, which is plantéd around their houses.

Cochineals have not long remained in this state when they produce swarms of young, which disperse and feed among the juicy leaves, and there produce a new generation. The insects having thus multiplied, are gathered three different seasons in the year: the first is performed by taking away the nests that were placed originally upon the tree; the second, by picking the cochineal from the leaves with pincers; and the third, at the approach of winter, by cutting off the leaves, which are yet loaded with insects: This last contains animals in various stages of their existence, and of different sizes; it is therefore reckoned of inferior quality on that account, and also, because, in scraping the leaf, some part of the epiderm necessarily comes off, and mixes with the cochineal. The Spaniards call it Granilla.

Having thus collected the infects, the next part of the process is killing them and preparing them for fale; and according as this is performed, the cochineal is supposed to be of different quality, and obtains various names, according as its colour is supposed to be more or less affected. That which is prepared by the gentle heat of overs, is of an ash-grey, or mottled colour, and named Saspeada. If the infects are deprived of life by plunging them with baskets into hot water, it then goes by the name of Renegrida, and is not covered over with that white powder, common to the other kinds. Lastly, is

bears

bears the appellation of Negra, if destroyed upon the hot plates that have been used for the roasting of maize.

But whatever be the method of preparation, that cochineal is always found fuperior, both in quality and quantity, which has been produced upon the cultivated trees. The cultivation of the opuntia, therefore, and the raifing cochineal, is a trade almost universal throughout Mexico: Eight hundred and eighty thousand pound weight of this article, is faid to be annually imported into Europe, where it fells at a high price *. Various are the uses to which our luxuries, and the arts subservient to them, have applied this drug. It is used in dyeing, and produces along with other fubftances, a variety of shades of fearlet: It furnishes painters with many lively shades of scarlet, crimson, and red: When ground into powder, it enters copiously into that carminative mixture which imitates the bloom of youth on the human cheek, and which is fo frequently had recourse to by the fair, who would repair the injuries of time +: had it really this effect, the Mexicans, we must allow, would command a freafure of greater value than all the mines of the new world.

Some attempts have been made by the French to find out the cochineal in St. Domingo, and to rival the Spaniards in their lucrative commerce in that article. The specimens, however, that have been produced in Europe, by one of the correspondents of the academy of sciences to in that island, have not, hitherto, answered expectation, but produced a colour so weak as gives reason to doubt whether the insect be of the same species. It is proba-

ble,

^{*} Reaumur, Tom. IV. p. 106.

⁺ Barbut, page 156. + M. du Hamel, Doctor of Medicine.

ble, however, that the art of transporting the cochineal, like the filk worms, into whatever country the opuntia may grow, will foon be discovered, and the monopoly of that drug be no longer engrossed by the Spaniards *.

^{*} Reaumur Tom. IV. p. 109,

GENUS XII .- 7 brips.

In this race of infects the rostrum is so obscure, from its minuteness, as to be scarce perceptible: The antennæ are stillsform and as long as the thorax; the body is of equal thickness through the whole length, and slender; the four wings are extended along the back, narrow in proportion to their length, and crossed at some distance from the base; the abdomen is bent upwards, and the seet commonly have two articulations. Only sive species are enumerated under this genus by Linnæus*.

The animals of this family are so minute, that they are not objects of discussion, unless viewed with a microscope; to the naked eye they appear rather like atoms than living animals. They reside commonly upon slowers, and under the bark of trees, and it is there also that their larvae are found, which only differ from them by their wanting wings and clytra. The thrips jurisperina, is one of the largest o'i this diminutive race; the thorax and abdomen are black, the clytra white. In autumn it is found in numbers among slowers, and on the juniper Linnaus, for a long time, was uncertain how to arrange it, from not discovering more than one pair of wings †. M. de Geer sirst observed that it had four exceedingly narrow ones, with which it slies but little, but runs quickly ‡.

²

^{*} Systema Naturæ. Ord. II. Gen. 12.

Fauna Swec. 4 Actes de Stockholm, p. 3. 1744.


CHAPTER IV.

SECTION I.

ORDER III .- Lepidopterous Infects.

This order contains all the different kinds of the moth and butterfly; infects characterifed by four wings, which are imbricated with scales; and by a mouth furnished with a spinal tongue, which can be extended or rolled up at pleasure. The bodies of all this class are covered with down, and their wings decorated with such a variety of brilliant colours, that they may be regarded as the most beautiful with which nature has enlivened and adorned the fields.





GENUS I .- Papilio, the Butterfly.

THE antennæ of this genus grow thicker, and are generally terminated with a knob or capitulum. When fitting, the wings are not folded down, but erected for that their extremities nearly touch each other above the body. They are distinguished from the moths, by sying in the day time.

Of this genus there are 273 different species, already known and described: To prevent confusion, they have been arranged by Linnaus into five distinct classes, viz. the equites, the heliconii, the danai, the nymphales and the plebeii*. Each of these classes is characterized by some particular part; so that every species when examined, discovers immediately to what class it eught to be referred.

There is no class of infects which hath been more accurately examined, or whose history hath been so fully detailed, as that of the buttersty and moth. Two large volumes have been dedicated solely to this tribe by the indefatigable Reaumur. The metamorphosis of infects is better illustrated by their history, because in them these changes are more frequent, and more perceptible.

Some of these animals frequently cost their skin, besides undergoing those more considerable transformations which introduce them into a new sphere of action; be-Vol. III.

^{*} Systema Natura, Ord. III. Gen. 2.

fore these develope, the outer skin is seen to wither and lofe the vivacity of its colours, owing to a new coat which already covers the animal beneath, and intercepts the juices which formerly circulated through it: After fome efforts, this dried covering is rent towards the back part of the head, where a fresh skin appears; and through this aperture, the worm makes his escape leaving his fpoils behind *.

After undergoing feveral changes of this kind, the animal prepares to undergo another fill more confiderable, which is to introduce it into the flate of a chryfalid, deprived of almost all motion, and incapable of taking food. This change is effected nearly in the fame manner as the foregoing; but in some it is very long in being accomplished. Several species of the butterfly worm, construct in a very ingenious manner a coque or nut of filk, into which they enter before their transformation, and in which they contime for nine months, without food, before their metamorpholes be accomplified f. During this long period, they are apparently inanimate, and take no food.

Various substances enter into the composition of the habitations configurated by these animals before their metamorpholes; fome are of filk; in others that is combined with other matter; several kinds construct no habitation, but are protected by a crustaceous shell, formed by a glutinous fabiliance, exfuding from their bodies: Some are fulpended vertically, while others hang horizontally by a thread which furrounds the middle of the body ‡.

The external form of the chryfalids varies according to the foccies of butterfly that inhabits them; in all, however, there are apertures opposite to the thorax, by which respiration 3

^{*} Reaumur, Tome I. mem. 3.

⁺ Tome II. mem. I.

[!] Idem Tome I. mem. II & 12.

respiration is carried on during the whole period of their inactive state. After the appointed time, when the animal was acquired sufficient vigour, the shell is broken, which at once constituted the grave of the caterpillar, and the cradle of the buttersty: the down already grown upon the infect has completely separated it on all sides from the shell, which, by the action of the head, is broken opposite to that part, and assords free egress to the prisoner it so long confined *:

The wings of the butterfly, on its first appearance, are closely folded; but by the help of a liquor condantly circulating through them, they are soon expanded, and sufficiently hardened, by the estion of the air, to endure the efforts of flying. It is then that the insect enters upon a more enlarged sphere of astion, with increased powers: He ranges from flower to flower, darting his rostrum into their nectaria for the delicious flores they contain. Then too in the full possession of every faculty granted to his race, he prepares to multiply and perpetuate it.

This last and most considerable metamorphosis, is attended with a greater change in the economy of the animal, than any of the preceding; not only the skin, but the treth, jaws, and cranium, are left behind. The largeartery which passes along the body, may be considered as a succession of different hearts employed in circulating the blood, which is at that important sera, observed to slow in a different direction from what it aid before, like the sectus of a quadruped after birth to Formerly it circulated from the extremity to the head; it now pursues a course directly opposite.

3 & 2

The

^{*} Idem, mem. 14.

[†] Vide Malphigi and Reaumur. Tome I. mem. 19.

The quantity of food taken by these animals in their last state, is comparatively small to what they formerly devoured. For a short time after their appearance on the wing, their excrements are voided in a greater quantity, and red like blood; this is, perhaps, the remains of that food, which they contained before their last change. Its appearance on the surface of the earth, has at different times been regarded as portentous of some heavy calamity, being supposed to be blood that had dropt from the clouds *.

Some of these animals are gregarious, and live in society during every stage of their existence; others live in that state during one period of their lives only. The duration of their life is various according to the weather; its warmth accelerates every step of their progress, and cold retards all their developments: A worm produced in a certain period of the summer lives only for three months, while the same species, if hatched a little later in the season, lives another year; hence, Reaumur, has devised a method of prolonging the lives of these animals greatly beyond their natural course. †.

The butterflies of every species are extremely prolific; a single semale at one birth produces several hundred eggs: and the most wonderful particular in the history of these insects; is the precautions by which they provide for the security of the young; some tear off the down from their own bodies to supply them with a covering ‡.

Various

^{*} Vie de M. Pierese par M. Gaffendi.

[†] Tom. II. mem. I.

[‡] Reaumur, Tome II. mem. 2.

Various infects prey upon the butterfly, or haften the approach of its diffelution. A certain species of fly makes its way into its body, while yet a worm, and there deposits its eggs, and, although this worm continues to live, and be metamorphofed into a chryfalid, no butterfly is produced from it, those internal parts that were effential to its prefervation, being confumed by the larvæ of the fly From the great fecundity and variety of the infects of this genus, they probably would foon cover the furface of the earth, did not nature provide a bar to their increase, by multiplying their enemies: hence they are destined to become the food of a great number of animals of various kinds, some of which swallow them entirely, others macerate their bodies; while many accomplish their destruction by gradually sucking their juices. A fingle pair of sparrows, in order to supply themselves and their young, may destroy, as is supposed, three thousand, three hundred and fixty butterflies in one week t.

The vast number of animals belonging to this genus, and the variety and richness of their colouring, afford ample scope to the painter for the exercise of his art. It was chiefly of these insects that Madame Marianne exhibited those exquisite drawings that constitute the peculiar merit of her work. It is principally in America, the East Indies, and China, that those species are found, which constitute the ornament of our collections. From the policy of the latter country, the ladies are circumferibed in their annuluments; there they domesti-

3 L 3 cate

[†] Vide a General Treatife on Husbandry, &c. by Mr Bradely, Profesför of Botany in Cambridge, 1724.

ate various species of this tribe of infects, and make their economy and manners the principal fludy of their lives; large glass apartments, furnished with proper food, being fitted up for the accommodation of these animals.

GENUS II .- Sphinx.

This tribe contains all those infects which hold an intermediate station, between the butterslies and moths. Their antenne are of a prismatic form, thicker in the middle than at the extremity; their wings sold down over the back when at rest; when on the wing their motions are slow and heavy. The moths, or phalene are nocurnal infects. The buttersty makes its appearance in the open day; while the excursions of the sphux are chiefly in the twilight of morning and evening *.

The infects belonging to this genus, though far lefs numerous in their varieties, prefent not uniformly the fame characters to the observer; some have angulated wings; in others, the wings are entire; some are known by a barb, or tust of hair growing from the anus; while many are distinguished by other differences in their external appearance, both in the larvæ and winged state. These last, Linnæus has termed adscitæ, as if brought into the samily by a fort of adoption, and thus distinguished from the legitimate sphinxes.

The name of fphinx has been given to the infects of this genus from the fingular attitude in which they are often feen in their caterpillar state; the hinder part of the body is then feen attached to the branch of a tree,

^{*} Systema Naturæ, Ord. III. gen 2.

456 SPHINX.

while the rest of the body like the celebrated sphing stands in an erect posture: The cod which they form, previous to their metamorphosis, and into which the animal enters, is constructed below ground of earth and grains of corn, interwoven with threads of their own manufacture: When invested with the powers of a winged insect, their voracity is diminished; they sly heavily from slower to slower, introducing the tongue into the nectaria for the liquor contained there.

The fphinx atropos is the most remarkable animal of this genus; the upper wings are of a blackish brown, waved irregularly above and below, with bands of a lighter hue; the under wings and abdomen, are of a sine yellow, variegated with transverse bands of black *. The worm from which this intest proceeds is large, smooth, and green, terminating in a kind of horn at the extremity †.

The most remarkable part of this animal is the reprefentation of a death's bead, which is seen upon the upper part of the thorax: This mournful picture is formed by a large irregular gray patch, marked with two black dots near the middle ‡. In the province of Brittany the people were afflicted during a certain season with an epidemic disorder, which often proved mortal; the inhabitants were greatly alarmed by the unusual numbers of these insects, which the common people imagined came to forecode their destruction, by this portentous representation upon their backs. The Royal Academy was confusted, whether the sphinx, by its uncommon numbers, might not be the cause of this calamity, so great was the alarm which their appearance created §.

^{*} Fauna Swecica n. 809. † Rai Insect. p. 362. n. 62.

[‡] Reaumur, Tom. I. p. 293. § Idem, ubi fupra.

The sphinx atropos, when hurt, has the power of uttering a dismal and melancholy cry, like a mouse; owing, as is supposed, to the violent rubbing of some of its scaly members. There are forty-seven different kinds of these animals.

VOL. III.

3 M

GENUS III .- Phalene. The Moths.

THE moths are the most numerous genus hitherto known, either in the animal or vegetable world: Linnaus has enumerated about four hundred and fixty species; and probably is far short of the real number. They are more frequent in warm and dry climates than in ours; and they are more plentiful in South than North Britain. In the former country, feventy different kinds have been collected; while in the latter, fearcely half that number have yet been noticed. The characters by which this numerous genus is deflinguished are the following: The antennæ fetaceous, diminishing in thickness from the base to the extremity; their wings, when at rest, are in general folded down; and they fly during the night. The race of moths have been turther fubdivided into eight different families. The first, called the attaci, have the wings hangirg downwards, and spread open; and the antennæ are cither pectinated or fetaceous: The fecond are named bombices, and have their wings covering the body horizontally, with pectinated and fetaceous antenna: The third family comprehends the nocture, whose wings are incumbent, and the antennæ fetaceous: Those of the fourth tribe are termed geometræ, and have their wings, when at rest, extended horizontally: The tortices are diffinguished by obscure wings, the exterior margins forming a curve: The pyralides have the inner margins of the wings laid one over the other: In the tince, the wings

wings are wrapped up, and folded round the body, fo as to give the infect a cylindrical form: And, lastly, the family of alucitæ have their wings split, or divided into different branches, almost to their base *.

The greater part of these phalance, or moths, when at liberty in the fields, only sly during the night, or towards the evening: Those which are domesticated in boxes made for that purpose, give indications by their sluttering within when the natural period of their activity approaches. During the day they remain quiet, and apparently reconciled to their consinement: they slutter throughout the whole extent of their prisons, and testify their impatience at their want of freedom.

All the durinal butterflies are provided with a rostrum for gathering, and for the reception of their food; a great part of the phalænæ are entirely destitute of that organ, while in others it is so small as scarcely to be discernible with the naked eye. This singular fact hath been fully investigated by the indefatigable Reaumur, who, after examining many moths with a strong magnifier, has not been able to discover the smallest vestige of a mouth †. A considerable number, therefore, of these animals, must pass the whole of their winged state without food: These cannot be destructive to cloth or surs, except while they remain in the form of worms.

The caterpillars from which the various fpecies of moths are produced, exhibit nearly the same variety of appearance as the winged infects which spring from them. Some are large, while others are extremely minute; many are provided with ten, others twelve and sourcen fect; the largest and most common have sixteen. Some

3 M 2 of

[&]quot; Vide Syft, Nat. Ord. III. Gen. 3. † Tom. I. Mem. vii. p. 286.

of the smaller caterpillars are smooth, and others covered with hairs, which produce an itchiness and an instammation, when they touch the human skin *.

All the caterpillars of phalænæ, after having feveral times cast their skin, spin for themselves materials of a habitation, in which they are to be transformed into chryfalids. Of all the inventions of infects to protect themselves during their state of imbecility, that practifed by the filk worm is most universally known; and if animals acquire a confequence or reputation from their connection with man, and the conveniencies with which they accommodate him, this infect may challenge, perhaps, a larger thare of it than any other animal whatever. Our luxury has brought filk into fuch general request, that it now may be deemed a necessary of life: The poor, in some countries at least, would find it almost impossible to procure the necessary articles of clothing, were woulden stuffs worn by all those who at present are supplied with filk.

The produce of the common filk worm has been found most proper for the purpose of manufacturing into cloth: That glutinous substance with which the filk is always covered when it first comes from the worm, and which gives it that adhesive quality so proper for constructing their edifices, sooner dries than in that of any other infect. The cods constructed by some other species are so firmly glued together, that no operation can separate the threads. The produce of many is by far too sine for any purpose in our manufactures, while that of another class is too coarse.

Several

Several very laudable attempts have been made, but hitherto without any confiderable degree of fuccess, to rear the filk worm in Britain. The public have lately been informed by a manufacturer in Paisley, of his having prepared a web entirely of the filk produced by worms of his own rearing *: And in the transactions of the fociety for the encouragement of arts, manufactures, and commerce, a number of very useful experiments are recorded with regard to the food and management of these insects †.

Probably the want of a fufficient number of mulberry trees has hitherto rendered ineffectual the efforts of our countrymen to introduce and rear any confiderable quantity of filk worms. From the attempts that have already been made, it appears that the white mulberry is preferable to the black in feeding, and that the latter is to be preferred to the lettuce. Twelve cocoons, the produce of worms fed upon the white mulberry, weighed feven penny weights two grains; while an equal number of those that had been sed upon the black mulberry weighed only six penny weight three grains: Six penny weights were obtained from the same number of worms fed upon common lettuce \$\mathbf{I}\$.

Endeavours to produce raw filk in Britain feems the more worthy of encouragement, as that country appears to possess some advantages of which Italy and many of the filk countries are deprived. In Italy, the chrysalids soon come to life; and it is there necessary to destroy them, lest, by eating their way out, they should injure the filk. In order to effect this, they are collected and placed

^{*} Vide Glafgow Newspapers, November 1791.

[†] The Rev. M. Swayne's Letters to Mr. More, Vol. VII.

Transact. ubi supra.

placed in heated ovens, where the filk, without fingular caution, is apt to be damaged. In our climate, where every progression of the infect tribe is slower, there is sufficient time to wind off the filk without killing the chrysalis.

But befide the injury that may be done to the filk in Italy, from the length of time which it is necessary to keep the chrysalis in these ovens, they are there obliged to suffer the moth to eat its way out of the largest cones, in order to have eggs from the most vigorous and healthy. Hence, they lose all the filk of these cones; whereas, in this country, the filk may be gathered while the moths are preserved. Thus, we seem to possess two striking advantages, which may probably compensate for the want of others which our climate has denied us.

Even the climate, however, of England is in some respects superior to those where silk is raised. In the south of France, the frosts are often so intense as to kill the mulberry leaves after they are out. At that season of the year, this is seldom the case in South Britain; which is also more free from lightning, and those sultry heats that have always been deemed prejudicial to the silk worm. From these considerations, the time may probably arrive, when our countrymen, by farther knowledge and experience upon this subject, may be enabled to avail themselves of these advantages, and become entitled to a rank as distinguished among the raisers, as that which they have long held among the manufacturers of silk.

The filk worm, however, is perhaps far from being the only infect of whose labours man might avail himself. There are many species very common, and immensely fertile, that might be beneficially employed in procuring

filk, did we know how to avail ourselves of their labours. M. de Reaumur has mentioned several whose productions ought to be subjected to experiment by the manufacturer; and in some suture period, some of these may be turned to account *.

Of many of the caterpillars belonging to this class of insects, the filk, it must be allowed, is altogether unsit for our purposes: their coques are not only coarse, but so seatily provided with filk, that the animal is obliged to join dry leaves, bits of wood, and other materials, in order to give stability to its edifice: Farther, many of them spin under ground, and their work consists only of joining and connecting together, by means of their threads, different particles of earth, of which their house is composed. These caterpillars, when kept by the naturalist, who waits for their phalænæ, must be supplied with earth in the boxes in which they are lodged; otherwise they will perish, from not being able to construct an edifice sit for their reception †.

The phalenee, in their chryfalid state, disser from the butterslies, in remaining for a much longer period before their metamorphoses into perfect infects is completed. Their form, too, is then different, being oblong, and not angular, like the chryfalid of the buttersly. Some remain in their coques for several years successively; especially if a cold damp situation has retarded their progress. So great is the essect of heat in precipitating their developements, that a moth in a warm exposure may be produced from its chryfalid, even in the depth of winter.

After the phalænæ issue from their last covering, some of them are destitute of wings: These are the semales of certain

Torne I. Mem. vi. † Barbut's Gen, Infect. p. 189.

certain kinds, who, instead of wings, have only short protuberances, altogether unsit for the purpose of slying. They have the appearance of large creeping animals of a different order, and can only be recognised for moths, by the shape of their antennæ, which are similar to those of the males, and by those scales with which the body of these animals is covered.

The phalænæ, though less brilliant in colour than the butterflies, are some of them far superior in size. The pavonia major of *Linneus* is much larger than any species of the butterfly yet known: In bulk, it seems rather to resemble a bird than an insect; and in the richness of its clothing sew of the seathered race can vie with it.

The three pavoniæ * have their name from those rich and brilliant colours which are supposed to have some resemblance to the gaudy dress of the peacock. Their wings appear covered with a glossy fur; and when spread, display tour eyes, decorated with vivid colours. Their caterpillar is also very beautiful; it is covered with hair, and it is the property of such to produce those scaly phakenæ, which are most distinguished by the brilliancy of their colours.

The moths in general do not fly by day; yet in the fummer evenings it is the light which attracts them into the dwellings of man: Then they are feen entering the rooms, and fluttering around the candles, where they often meet with a painful death. This fondness for light has suggested to the curious a method of catching these animals, by carrying a lanthorn into a bower, around which they all slock, and numbers are led into captivity.

The

^{*} Syft. Nat. Ord. III. Gen. iii. fp. 7.

The phalana of the filk worm, bombyx moris, frequents the leaves of the mulberry, upon which it is univerfally known to feed. It is one of those infects that has deservedly attracted the notice of almost every naturalid; and to a particular description of it we direct the reader in the note below *.

The most beautiful of the English infects are the following phalænæ, viz the large emerald, or geometra papilionaria of Linneus; The thorax, wings and abdomen, are of a lovely pea-green; the wings are decorated with small indented bars, and marginated with a fringe of golden yellow.

The peach blossom: The head and thorax of this insect are dark brown; the superior wings are of the same colour; but each is ornamented with five spots, of a pase pink, or rose colour, and appear very like small roses painted on the wings: The abdomen and inferior wings are of a fine pale brown colour: The caterpilar seeds on raspberry leaves; changes into a chrysalis in June, and in July comes forth a moth ‡.

The phalæna which in its larva flate is destructive to woollen cloth, is of a narrow oblong shape, having the wings raised towards the extremities: The head of this infect is white; the superior wings black towards the base, and white as they approach the tips: The inferior

Vol. III. 3 N wings

[&]quot;It is named in the Fauna Sweeiea, phakena pedinicornis, bombyx dicta, Aldrovandus describes it, Ink. 2. p. 280 & seq. Johnston, Inf. Mousset, p. 181 M. de Merian, Europ. Lister in Goedart, p. 82. Charleton onom. p. 40. Albin, Engl. Insects, p. 60. Reaumur, Tom. II.

[†] Systema Nat. Ord. III. Gen iii-

[‡] Vide Harris's expectation of English Insects, p. 53, where many exceltent drawings of the English insects may be seen.

wings are of an ash-colour; the body and the legs black, and the antennæ setaceous*.

Another moth, diffinguished by its descructive effects among woollen articles of apparel is thus characterised in the Fauna Swecica: The autennee are setaceous, the tongue spiral, and the thorax marked on each side with a white spot. This animal is of small size; the wings associoured; the head, legs, and the whole under side of the body are white f.

These moths construct the abode in which they reside, of the grains of wool which they gnaw off the cloth: Their food is of the same substance; and what greatly increases the extent of their devastations is, that every step they advance upon cloth, feeling themselves incommoded by the wool in their way, they gnaw a smooth passage for themselves, like a man with a scythe in his hand, cutting down the grass of the meadow as he goes along ‡.

These habits of the moth render them among the most destructive of insects. The most costly articles of fur are those which are not worn every day; and for that very reason they are most exposed to their attacks. The methods for preventing their devastations may be reduced to the two following: either we must destroy the insects, or render our clothes disagreeable food for them.

The moths may be destroyed by oil, or the sumes of tobacco. And as wool is never subject to their depredations, so long as it remains upon the sheep, the most likely

likely

[&]quot; Rai Insect. p. 204. n. 98.

[†] Vide Reaumur, Tom. III. & Regne Animale, Tom, II. p. 160. n. 83:

Reaumur, Tom. III. p. 5.


likely method of rendering it disagreeable, is by restoring that greafy substance which is taken from it by washing, and other processes which it undergoes when manufactured to

[🛉] Reaumur, Tom. III. Mem. iii.

CHAPTER V.

ORDER IV .- Neuropterous Infects.

I us order contains fix genera of infects, all of which are diffinguished by four naked wings, of a reticulated structure, and by appendices springing from the extremity of the abdo non, which is unarmed with a sting: These appendices are characteristic of the male sex in this order of animals.





GENUS I .- Libellula. The Dragon-Fly.

Or this tribe of large infects there are twenty-one different species remarkable by the formidable apparatus with which their mouths are armed. Their jaws are always more than two in number, and fitted for gratifying that voracity by which the libellulæ are diffinguished. The antennæ are shorter than the thorax; the wings are always extended, and, in fome, measure feveral inches from tip to tip. The male is diffinguished by a kind of hooked forceps, by which the extremity of the abdomen is terminated +.

The dragon flies are univerfally known, from their large fize, and the frequency with which they every where appear. The organs of generation are differently fitnated in the male and female of this tribe: In the former, they are placed upon the under fide of that part of the abdomen which lies between the inferior wings; while, in the latter, they are found at the tip or end of the abdomen. The feafon of love in all infects, from the brevity of their lives, is necessarily short, but it is bufily employed. It is only the large infects, fuch as the libellulæ, that are frequently feen colebrating the rites of Venus; and among them the operation is performed with 2 frequency and dispatch that feems perfectly to corre-

fpond

^{*} Linnaus Syft. Nat. Ord, IV. Gen. i.

ipond with the transient nature of their existence, and the numerous race they transmit to posterity.

The addresses of the libellula to his semale seem carried on in a rough and intrepid, but essications manner. He hovers about on the wing, till the object of his almours make her appearance; he then watches an opportunity of seizing her by the head with those pincers with which his tail is armed. It is thus that the ravisher travels through the air, till the semale, yielding to superior strength, perhaps to inclination, forms her body into a circle, that terminates at the genitals of the male, and thus accomplishes the great purpose of nature. It is while these kind of rapes are perpetrating, that the libellular are seen coupled in the air, exhibiting the form of a ring *.

The female, when pregonnt, retires to the fide of a ditch or pond, when, by the affiltance of a flick or reed, the creeps, or lowers herfelf down, by moving backwards, till the tip of the tail is immerged about half an inch in the water; the is then feen feized with a kind of trembling or flaking of the bedy, during which efforts the deposits a fingle egg in the water; afterwards the immerges her tail a second and a third time, when the same operations are performed. The tail is withdrawn from the water, by contracting its annuli; and by the pressure of these against the extremity of the tail, from whence it is nitimately separated, by shaking that part in the water.

The eggs thus protruded by the libellula are of a white colour, and oblong form, refembling these produced by the vomitoria or common blow-fly. The caterpillar which

^{*} Barbut's Genera Infectorum, p. 205.

which iffues from these eggs, at first feeds on small aquatic infects, gradually acquiring more voracity and boldness in the pursuit, in proportion as its ilrength increases, at length it becomes fo infatiable that it is faid to devour even those of its own kind +.

The form and colour of these worms are extremely difgusting: They are supposed to have gills like sishes; and beneath the head is placed an inftrument excellently adapted for fe-zing and holding their prey. It is furnished with a forceps at the end, and can be advanced or drawn back with all the agility of the human hand.

The caterpillar remains in the fame state for nearly twelve months before it has attained to its full fize: When the period of transformation has arrived, the worm repairs to the margin of its pend in quest of a convenient place of abode during the feafon of its inaction. It there attaches itself to a plant or piece of dry wood; and the fkin, which has gradually become parched and brittle, at last splits opposite to the upper part of the thorax. Through this aperture the winged infect quickly pushes its way; and being thus extricated from confinement, begins to expand its wings, to flutter, and finally to launch into the air, with that gracefulness and ease peculiar to this majestic tribe I.

No particular time feems appointed for the metamorphofis of the libellula into its winged flate; the different species are continually emerging from the water from April to August: For as the times of copulation are various and frequent during the whole fummer, fo the larvæ or caterpillars are found of different fizes, according to their age. The finaller kinds, however, generally

make

make their appearance before the largest; because, from breeding in shallow water, they sooner feel the influence of the sun on the approach of spring.

The manners of these insects must no doubt be greatly altered by a change, which not only confers upon them a new form, but introduces them into a different element. The complete insect, in its winged state, however, still continues to pursue the same food, and remains insectivorous. The lepidopterous insects, the buttersies, and phalænæ are destined for the support of the larger libelluse; they are a part of those numerous tribes that are appointed to confine those prolitic genera within due bounds †.

The libellula grandis is the largest of this genus found in Britain, and is perhaps not inferior in bulk to any infect which this country produces: The fore part of the head is yellow; the eyes brown, and so very large, that they meet upon the top of the head. The thorax is dun coloured, with two oblique bands on each side, of a lemon colour. The abdomen, which is very long, is likewise of a deep busse colour, often spotted with white upon the top and bettom of each segment; the small appendices which terminate the abdomen, are in this species very long; the wings have more or less of a yellow complexion, and are distinguished by a brown spot on the exterior edges ‡.

The libellula forupata § is another large infect of this genus, which expands four inches and an half. The nofe is yellow, having a black line on the prominent part; the thorax is black, with feveral broad yellow firipes,

twe

Harris Exposit, p. 51. | Barbut, p. 206.

Syft. Nat. Ord. IV. Gen. i. No. 11.

two of which appear on the front, and two others between the ligaments of each pair of wings; the abdomen is also black, having two streaks resembling a crescent on each segment; the wings are transparent, and, but for a slight tinge of amber, almost white *.

The libellula whose body is of a thining green, and the wings of a gilded brown, is smaller than either of the foregoing, and is thus characterised by Mr. Ray †: The body is green and azure; the wings adorned towards the middle each with a large spot, of a deep blue, inclining to black. The feet of this insect are black, and the wings have no spot on their exterior margin. From the brilliancy and richness of its colours, it has been called the king's fisher; it frequents little rivulets of water, over-shaded with bushes ‡.

Vol. III.

30

* Harris's Exposit. p. 78. † Infect No. 9.

GENUS II .- Ephemera. . The Day Fly.

THE mouth of this genus has neither teeth nor palpi; the infects which compose it have two, sometimes three stemmata, situated above the eyes, and larger than is usual in other intects. The wings, when the animal is at rest, stand erect, the upper much larger than the inferior pair; the tail is surnished with an indefinite number of setaceous appendices †.

The infects of this genus have been denominated ephemeræ, or day flies, from the shortness of their lives; but the life of a day is in proportion to that of several animals of this class, what the lives of the antedeluvian patriarchs were to ours. The period allotted by nature for the existence of certain ephemeræ is only a few hours; of some, even less than an hour. To some infects, indeed, that bear the generic characters of ephemeræ, she has been more indulgent; for these are destined to see more than one revolution of the sun ‡.

All the day flies are produced from hexapodal worms, which are afterwards transformed into chryfalids, of a form nearly refembling their own; like the parent worm, they walk on the same number of legs. Both the worm and the chryfalis are aquatic, and are provided with gills like



like those of fishes, and probably destined to serve the same purpose.

They are feen arranged along each fide of the abdomen, one on every fegment, from the first to the fixth or feventh. When the infect is at rest, these branchiæ are seen moving with great rapidity; and from their position, Reaumur has distributed the chaysalids of ephemeræ into three distinct classes.*

Some of these chrysalids walk or swim up and down in the water, attach themselves to plants, or conceal themselves below the stones: Others have the first pair of feet formed for digging into the ground, and besides, are provided with two instruments on the forehead, still better adapted to that purpose: with these, each animal digs itself a hole on the river bank, where it constantly resides. The brinks of the Marne and the Seine in France are seen thickly perforated by these animals: The holes thus excavated are generally nearly on a level with the surface of the water, and have commonly two apertures; one by which the worm enters, and another by which it returns.

These animals, while they remain in these subterraneous retreats, offer but sew materials to the naturalist who would record their history: Though many of them remain there for two years, and feed upon the earth in which they are lodged, yet they are effectually concealed during all that period from his observation. Probably the most interesting incidents of their lives are those which happen in our view, and are confined to the narrow space of a few hours. Their story becomes then eventful; for in that short period they are transformed into slies, the second of the same secon

^{*} Tome VI. mem. II.

males impregnated, and the eggs deposited; and these functions are no sooner over, than the active beings who performed them sinish their operations for ever *.

On the banks of the Siene and the Marne, in the vicinity of Paris, the ephemeræ exhibit a fingular spectacle for some days about the middle of August. For several hours after sunset, they rise in such vast multitudes, that they appear like the slakes pressing upon each other in a heavy sall of snow. By and by, the whole surface of the earth is covered with the swarms that have salen upon it, after having sinished their short existence.

The chryfalids that remain under the water begin their transformation in the evening, and complete it by eight or nine o'clock; an operation painful and difficult to other infects, is with them performed with great celerity and apparent cafe. No fooner has the chryfalis reached the top of the water, than its prifon burfls open, and the winged ephemera foats into the air. Millions after millions are thus conflantly taking wing, till the air becomes darkened with their numbers: They are in that element but an inftant, when abundant showers of them fall back to the ground.

The females, after their fall, are bufy in performing the last function of their lives, which is, depositing their eggs. Such as have dropped upon the ground leave them there; while these that have tumbled into the river produce two separate clusters of ova, each containing no less than three hundred and fifty. All this is the work only of a moment; for other insects are not sooner delivered of a single egg, than the ephemera of seven hundred.

Some

Some naturalists have pretended, that these eggs were all impregnated, like those of fithes, by a kind of sperm ejected from the body of the males. This opinion, however, is combated by Reaumur *, as being attended with many difficulties. That naturalist, by the help of a lighted torch, has examined their animals narrowly in their winged flate; and although that method of furveying infects, whose numbers occasion so much consusion, be uncertain, he afferts, that he observed them engaged in fhort copulations; an affertion the more probable, fince, if the ephemerm copulate, that act behoved to be more instantaneous than in any other race of beings.

Another species of the day fly is seen to cast a skin. even after it has arrived at its winged flate: Thus, however flender their wings might at first appear, they must, as well as the whole body, have been enveloped in a coat which the animal then drops +.

The ephemera vulgata, or common day fly, is the largest of these insects with which naturalits are yet acquainted; it carries at the extremity of the abdomen three brown threads, nearly of equal length with the body, which is all over brown. The wings are ornamented with brown veins, which form a net-work.

In Carniole, a province of Germany, this species is so numerous, that the peafants think they make a bad harvest of them, if they do not unload upon their land many carts filled with these insects. They make excellent manure; but the number of animals of fo small a fize necesfary to fupply the quantity, must exceed the power of imagination to conceive.

^{*} Vide Biblia Natura. † Reaumur, Tome VI. Pref.

GENUS III .- Phryganca.

THESE infects, as well as the last, are of a small size, and are at first sight easily distinguished by the antennes, which are longer than the thorax, or even the whole body. The mouth is without teeth, but surnished with sour palpi; the sour wings are incumbent; the under ones, when unfolded, are very broad in proportion to the size of the animal that wears them *.

There are twenty-four species ranged under this tribe, which Linneus has divided into two samilies; the first char Sterited by having the sets which spring from the extremity of the abdomen truncated like the beards of an ear of corn; and the second, by having the abdomen simple, without appendices.

The phryganeæ, before they become inhabitants of the air, have, like the ephemeræ, been aquatic animals; their apartments in the water are fingularly conftructed; their form is that of a tube, the inward texture of which is filk, while it is outwardly composed of fand, straws, or chips of wood. When the animal within finds its chryfalid state approach, it stops up the ends of this tube with threads of a loose texture, through which the water may penetrate, while it prevents the appreach of voracious insects.

The

^{*} Systema Naturæ, Ord. IV. gen 3.

The precautions of these aquatic larvæ, in securing themselves in these retreats, are also necessary to protect them from sishes, who are very greedy of them. Some of the phryganeæ of stagnated water, such as lakes and ponds, cut the water lentil in regular squares, from which they construct their edisces. The common trout is one of their greatest enemies; and many other sishes covet them so eagerly, that, when stripped of their crust, they make excellent bait.

The phryganea bicauda of Linneus frequents equally the banks of rivers and of standing pools; the wings are a third longer than the body, narrow at the top, and broad below, and reticulated with brown veins; the colour of this insect is a dark brown, with a single longitudinal streak passing across the head and thorax; the antennæ are long, siliform, and of the same colour with the legs and body *; the appendices of the tail are of equal length with the antennæ.

The phryganea faltatria is one of the smallest species of this genus; the wings are decorated each with a green and white spot; the antennæ are longer than the body; and when examined with a microscope, appear velveted; the wings are of a bright colour; the small veins by which they are reticulated being white at the base, and blackening towards their extremity ‡.

^{*} Barbut's Gen. Infect. p. 216. † Fauna Swecica, phrygan. fpec. 11. † Regne Animale, Tome H. p. 117.

GENUS IV .- Hemerobins.

This tribe of infects have obtained the name of hemerobii from the thortness of their lives, although they continue in their winged flate longer than a day. This genus is diffinguished by having the mouth armed with two teeth and four palpi; the wings hang down, without being folded; the antenne are setaceous, advanced before the head, and longer than the thorax, which is convex.

Nature feems to have fpread plant lice upon almost every plant, for the mantainance of thousands of her offspring. In their larva state, the hemerobii are great devourers of these animals; and from that circumstance they have obtained the name of lions of plant lice. Even after their transformation, the hemerobii retain their carnivorous habits: Not satisfied with making war upon the plant lice, who tamely suffer themselves to be devoured, they do not spare those of their own kind.

The moment these insects enter upon life, they find themselves, by the precaution of their ancestors, situated among thousands of small animals, which, without any effort, they destroy. In less than a minute after the plant louse has been seized, the whole viscera and justices are sucked from its body. Some species make a covering for their body of the careases which they have thus empticed, and carry them about as trophies upon their backs.

The

The eggs of the hemerobius offer a curious spectacle to the observer, and for a long time were deemed parafitical plants. Each of them is supported upon a long pedicle, resembling the stalk of a plant, of which the egg appeared to be the slower. In this form they are seen in clusters upon leaves, where, as soon as they are hatched, they find subsistence among the surrounding plant lice.

In about fifteen days, the larva attains to its full growth, when it weaves to itself a small white silken cod of a close texture: there the hemerobius lodges about three weeks, till its metamorphosis into a winged animal is completed: If, however, the cod has not been spun till Autumn, it remains in it till Spring, when it is transformed. The slight of these insects is slow and heavy; some of them have an excrementitious smell.

The hemerobius perla of *Linnæus* is one of the most beautiful animals of this tribe: The eyes are golden, and the whole body of a yellowish green *; the wings are very large, diaphanous, and of nearly the same colour with the body; it inhabits garden plants, and when touched, has the smell of ordure †.

Vol. III.

3 P

Pai Insect. p. 274. Vide Mouffet, p. 62. & Grew, mus. p. 156.

GENUS V .- Myrmeleon. The Lion Ant.

THE mouth of this voracious race is armed with jaws, two teeth, and four pulpi. In the male fex, the tail is familhed with a kind of forceps, for ned by two straight threads; the antennæ are in the shape of a club, and of equal length with the thorax

The lion aut, concerning which fo much has been faid by naturalitis, is of five kinds, all distinguished by an uncommon voracity, and cunning in fecuring their prey; their habits of rapacity feem born with them; and during their whole lives they are fapported by the game which they catch. At first fight, these animals feem but ill fitted for their manner of life: The most unwieldy infect can early escape from them; for they cannot advance upon it a fingle step; all its progress in walking being backwards; by stratagem, however, this lion masters infects far superior to it in strength. In the loose fand it digs a hollow refembling a funnel, where it takes its ftation at the bottom, every part being concealed except the forceps with which the head is armed. This inftrument, which it can open or shut at pleasure, is happily constructed for feizing and penetrating the hardest infect; and unhappy is the animal, who, in purfuing its journey, flumbles into this cavern. In vain it endeavours to fcramble up by the edges, which are continually giving way under its feet; it tumbles to the bottom, where it is pierced by the forceps of the lion ant lurking below.

If the infect be fmall, and the grains of fand, notwithfranding the declivity of the funnel, do not give way under its feet, the myrmeleon has nother invention by which he renders himfelf mafter of his prey: With his head, which is flattened, he throws up repeated showers of fand from the bottom of the funnel, which falling upon the fides, force down the infect till it comes within reach. The fatal instruments with which this animal feizes its prey, are each a fort of mouth or trunk, by which it sucks out the whole intrails before it is drawn out of the body *.

When the lion ant has attained its full fize, it confirmeds for itself an edifice, the external parts of which are particles of fand or earth combined together by filken threads: the interior cavity is lined with pure filk, white and gloffy like fat.n. Within this ball the myrmeleon is changed into a chryfalis, of a curved or semicircular shape, displaying all the parts of the perfect infect that is foon to iffue from it.

After the chryfalts burfts, the winged infect which makes its escape is of a gray colour, with a long slender body, resembling the libestula. In this country, the myrmeleon is very scarce; a few, however, are found to breed among the loose earth at the bottom of walls which have a south exposure. In that dry, pulverized, and sandy earth, their eggs are protected from rain, till they are hatched by the sun. Vallishieri and Poupart first gave the history of the lion ant; that of the former is in the form of a cialogue between Malphigi and Phiny, in which the modern informs the ancient naturalist of the singular manageners and metanacipitosis of these animals.

3 P 2

Externally,

^{*} Reaumur, Tome VI. mem. z.

Externally, the lion ant presents no seatures that are remarkable. The head is broad, of a brown colour, spotted with yellow: On each side are placed two large eyes, and beneath them the antennæ, thickening towards the point, and shorter than the thorax. The neck is longer than that of most insects, and can be stretched or shortened at pleasure. The thorax is of a dirty white, and upon the annuli which compose it are seen by the microscope, the organs of respiration †.

Reaumur, Tome VI. mem x. p. 339e

Maseria W. E.

GENUS VI .- Panorpa. The Scorpion.

FORMERLY there was only one species of these animals known, till Linneus discovered four different kinds, which he has enumerated in the latest editions of his system of nature. They are all distinguished by a cylindrical proboscis, of an horny transparent substance; by two palpi, and three stemmata above the head. The tail of the male sex is surnished with a weapon resembling the dart of a scorpion; a circumstance from which almost all naturalists have given it that name †.

The formidable inftrument with which the tail of the male fcorpion is armed, is not formed for any hostile purpose; it is used by him in laying hold of his female during their amorous embraces.

There are varieties of the common fcorpion; that most frequent is of a dark brown, and yellow upon the sides. The tail is formed by the three last fegments of the abdomen, and is of a maroon colour; the wings, which are as long as the body, are diaphanous, and reticulated, with sibres and streaks of a brown colour. There are some

fome varieties of this fingular animal: Inflead of the threaks on the wing, we find fome with a fingle transverse firipe of black irregularly placed across the wing; others are found with their wings entirely white, excepting the tips, which are black.

GENUS VII.-Raphidia*.

This tribe is characterised by two teeth, a flat head, of a horny substance, four palpi, and three stemmata; the antenne are siliform, and as long as the thorax; the anterior part of which is lengthened out like the neck of a bird. The tail of the semale is terminated by an appendice resembling a slexible crooked bristle †.

There are only three species of this genus yet discovered; the first of which is termed the ophiopsis, from the sigure of its head and long thorax. Its form is the most singular, perhaps, of any animal in this class of beings. The shape of the head is broad before, and narrow where it joins the thorax, resembling a heart; it is slat, smooth, and black like horn: the thorax is of the same colour, narrow, and cylindrical: the antennae are setaceous, and composed of many articulations: the abdoment is brown, marked with white transverse lines, the wings are radiated and diaphanous; the exterior edges in some subjects have each a black spot: the anus terminates in a curvated setaceous bristle, about half the length of the abdoment.

^{*} Barbut's Gen. Infect. p. 227.

[†] Syst. Nat. p. 916.

A Regne Animale, p. 112.

CHAPTER VI.

ORDER V .- Hymenopterous Infects.

This order contains all the bees, hornets, wasps, and ichneumons, insects which are diffinguished from those which we have already reviewed, by possessing a sting, and four membranous wings. Some of them the industry of man has turned to advantage, by reducing them into a domestic state, while the greater part still remain the free fervants of nature, and may be regarded among the most inveterate enemies which men have to dread among this race of animals.





GENIII.











GENUS I .- Cynips. The Gall Fly.

THE mouth in this genus is armed with jaws, but have no probofcis; the fting is fpiral, and most frequency concealed within the body.

Those innumerable and various excrescences which are seen upon the leaves, branches, and roots of trees, are all the productions of different kinds of insects. Of these, some so nearly resemble the natural productions of the plants, that they are sometimes taken for their fruit or their flowers: A particular kind of ivy produces galls that are actually eaten as fruit by the peasants in some parts of France; and in Constantinople, there are some of these productions brought to market for sale *.

Some of these excrescences have within a single large cavity, in which several insects live and associate together: Others have a number of smaller cavities, with communications between them; a third class contain a variety of separate cells, sometimes a hundred, each occupied by a single insect, which has no communication with the rest. These productions are also of various sizes, forms, and consistency; some are spongy, while others are hard like a nut. Of the latter kind, those are best known which are imported from the Levant for the purpose of dyeing cloths.

Vol. III. 3 Q All

^{*} Reaumur, Tom. III. Mem. xii.

All these apparently monstrous productions are occasioned by the puncture of insects when depositing their eggs, or by their bite when collecting food; in both cases, the animals subsist upon it; and the more they eat, the more vigorously does the protuberance continue to grow, till at last it forms a fort of impenetrable fortress, to protect its inhabitants till they have gone through their different metamorphosis, and at last taken wing. But the best concerted schemes of insects, and of man, are unequal to secure either from every accident that may occur. Impenetrable as the habitation of the gall fly may appear, its walls are often perforated by other insects, who deposite eggs there, that are soon to become rapacious worms, and to lay the dwelling waste.

The ancient opinion concerning the animals found in these receptacles was, that they were spontaneously produced from the rotten wood of the plant. Afterwards, it was believed, that the roots of plants had the power of fucking up, along with fap from the earth, the eggs of infects, and that these were animated as soon as they flopped circulation through the fibres of the tree. Even the intrepid Rhedi, who combated the prejudices of his age with fuccefsful boldness, had recourse to a kind of vegitative foul in plants, by which he accounted for the production of these animals. Malphigi at last explained their true origin, from eggs deposited there by those of this kind. The fame naturalist gives an ingenious account of the formation of these excrescences themselves by means of a liquor deposited by the sly, mixing with the fap of the tree, and caufing a fermentation at the part. The fimple extravalation of juice from the wounded

plant,

plant, is however, fufficient to account for the largest galls that have yet been observed *.

The cynips quereus folii †, is of a burnished shining brown colour; the antennæ are black; the legs and feet of a chesnut brown; and the wings white, without marginal spots. Those smooth round galls that are seen under the oak leaves, are the birth places of these insects: commonly a single one is found in each gall. Instead of the natural inhabitant of this gall, a larger insect of a brown colour is sometimes seen to proceed; this is the ichneumon, who is neither the builder nor the legitimate owner of the dwelling, but a parasite produced from an egg deposited there by his progenitors.

Of all the trees with which we are acquainted, the oak affords food and an habitation to the greatest number of insects. There are above sifty different species that in this country inhabit that plant, and probably in warmer climes the numbers of its tenants is much greater. The Norway ink, so celebrated for its colour and permanency, is the produce of a gall in that country, similar to that seen upon our oaks in the month of June. The sly which produces it is perhaps the same ‡.

The gall fly of the rose, cynips rose, is found on the gall of that shrub, and is distinguished by the black protuberances of the antennæ; the abdomen below is ferrugineous, the seet yellow, and the wings without spots §.

^{*} Reaumur, Tom. III.

[†] Syft. Nat. spec. 5.

^{. ‡} Barbut, p. 234.

[§] Fauna Swecica, No. 938.

GENUS II .- Tenthredo. The Saw Fly.

THE family of faw flies have two jaws, but no trunk or probofcis. Their fling, which is almost hid by the abdomen, is of a fingular structure, being composed of two plates, and dentated like a faw. The abdomen is closely united to the thorax, without the intervention of a small stalk, as in the former and succeeding genera.

From the shape of the antennæ, the tenthredines have been arranged into six tribes, consisting of about thirtynine different species. The most remarkable sast in their history seems to be the application of her saw by the semale, in cutting out a space in the twigs or buds of trees to contain her eggs: The whole process is performed with great ingenuity; and from the tameness of these animals, may easily be observed.

The eggs thus deposited have the peculiar property of increasing their fize every day, till the young worms are ready to burst from them. The worm of the tenthredo has often been taken for the filk worm; an animal which it nearly resembles, except in the feet, which are more numerous: Like the filk worm, too, the young tenthredo constructs a cod in which it may undergo its last transformation; and its edifice consists of two coats of filk, the external coarse and strong, the interior of a saner substance *.

The

The tenthredo femorata is, as the name imports, burdened with two large thighs; the whole infect is black, except the antennæ and tarfi of the feet, which are yellow †.

The tenthredo luctuosa alni. This quiet, melancholy, and beautiful fly is found among alder plantations, and is often satally entangled in the clammy juice that issues from their leaves. The head and trunk are of a fine yellowish brown; the antennæ a more dusky shade of the same colour, and the eyes blue: The body is coal black; the segments of a brownish yellow; the wings are of a pale yellowish brown, with little yellow risings upon their sibres, and a margin of yellow ‡.

There are no less than fifty-five species of the tenthredo enumerated in the Linnan system; but the Swedish naturalist seems himself to suspect, that some kinds are twice mentioned. Colour, though an obvious, is no certain character of insects. In some, it differs with the seafon, in others, with the sex; and in all it glows according to the creature's health and vigour.

† Syft. Nat. spec. 1. ‡ Barbut's Gen. Insect. p. 2382

GENUS III .- Siren. The Tailed Wafp.

THE animals of this genus are not fo numerous nor diverfified as the laft, but are greatly superior in fize; the mouth is armed with two firong jaws; the palpi short, and articulated; the antennæ are filiform, and confift of near thirty articulations; the ling is long, sliff, ferrated, and projecting beyond the abdomen; the wings are plain, and of the shape of a lance; the abdomen is slender, terminating in a point towards the sting +.

Of the feven different kinds of firices, the gigas of Linnaus is the largest; the male, indeed, as is frequently the case, is one third less in fize than his female, and has neither spine nor sting at his extremity: That instrument, however, is formidable in the other fex; it confifts of three laminæ; two at the fides, which ferve for fheaths, and one in the middle, fomewhat ferrated, which is the real sting.

The wings of this species are large, yellow, and veined; the thighs short, and black; and the legs, as well as the tarfi, are yellow; the antennæ are of the same colour, and are inferted into a black head, which has apparently 3

parently four eyes. What appears to be the fecond pair of eyes is only two large yellow fpots behind the other. The beliy is cylindrical, and confifts of nine different fegments, fome of a black, and others of a yellow colour*.

Barbut's Genera Insectorum, p. 242.

GENUS IV .- Ichneumon.

This genus contains a numerous lift of infects, whose manners appear singular, even amidst the wonders which this class of beings is continually displaying. It contains upwards of seventy different species, all characterised by the antennæ, which have more than thirty articulations; by the mouth, armed with jaws, but without a tongue; and by the abdomen, which is joined to the body by a pedicle or stalk, and which terminates in a projecting sting, inclosed in a cylindrical sheath †.

We have already had occasion to observe one striking peculiarity in the manners of some infects of this genus, who make their way into the body of the caterpillars of different kinds, and there deposite their eggs. It is from the signal services which they perform, in destroying caterpillars, plant lice, and other insects, that they have derived their name: These animals are devoured by them, as the genuine ichneumon, or mangouste, destroys the crocodiles.

As the species and varieties of the ichneumons are prodigiously numerous, so their manners are extremely diversified: In the general outlines of their character, they all agree, being remarkable for their depredations among the insect tribes. In some, the semale is surnish-

ed with a wimble, attached to her abdomen; and with this instrument, though delicate, she is capable of piercing through lime and plaster. The larvæ of wasps and mason-bees are the devoted prey of this species, which no fooner espies one of their nests, than with its wimble it perforates the mortar of which it is confiructed. This operation, which is performed with fingular dexterity, is no fooner finished, than it deposits its eggs in the infide, to the number of one, two, and fometimes more. Some are contented with gluing their ova to the skin of a caterpillar, while others penetrate through it, and depolit the egg in its body +. The ova hatched within the caterpillar, after being quickened into life, preys upon the intestines of that animal, without, however, destroying its vitals. Upon the life of the one animal that of the other feems to depend; and the ichneumon spares the caterpillar upon which it feeds, till it is about to enter into its chryfalis state. In the mean time, the caterpillar in which it is enclosed is apparently healthy, and prepares to undergo the same transformation; a function which it is feldom able to accomplish, because the interior parts effential to the butterfly, though not to the worm, are destroyed t. Often these caterpillars, which have been the cradle of thirty or forty ichneumons, are feen fixed to the bark of a tree, as if they were fitting upon their eggs; and it is discovered that the larvae which were within their bodies have foun their threads, with which, as with cords, the unhappy caterpillar has been fastened down, and perished miserably.

The eggs of the ichneumons which have been fimply agglutinated to the outside of the larvæ, remain not long Vol. III.

3 R in

Regumur, Tome II. Mem. xi. | Idem, Tom. II. Pref.

in that position; for the young animals produced from them easily make their way into the body of the caterpillar on which they were placed.

Various are the fizes of this carnivorous race: Some are fo finall, that the plant loufe, the curculio, or the fpider's egg, is made the cradle for their young; and all the infects upon which they fix undergo the fame fatal deftiny, fromer or later perifhing by their depredations. Those carcases of plant lice which are seen motionless on the rose-tree leaf, have been each the habitation of a small larva, which, after having devoured the intrails of the plant loufe, has performed its metamorphosis under the empty skin, from which it has afterwards fallied forth a winged ichneumon.

But though many of these insects are so extremely minute, there are others, which, from their size and intrepidity, are formidable to the spider. Many of these challenge that animal to open combat, and having run him through with their stings, tear him to pieces; thus avenging the whole race of slies of the injuries they suffer from that most dreaded enemy of their kind.

Ichneumen aphidum. This infect is almost wholly black; the abdomen, towards the base, and the sect, are yellow; the antennæ are black. This species, when about to deposit its eggs, bends the abdomen till the anus nearly approaches the thorax, and thus penetrates into the anus of the plant louse, in which it lodges its progeny †.

The parafitic ichneumon ‡. The larva of this infect adheres to the body of a caterpillar, upon which it feeds.

The

Regne Animale, Ord. V. Gen. iii. spec. 3.

[†] Vide Actes d'Upfal, 1736, p. 29, N. II.

The head and the antennæ are black, the latter longer than the body; the feet are yellow *.

The black ichneumon +. The antennæ, head, thorax, and abdomen of this species are all black: it is farther distinguished by the three siliform appendices of the tail, which are longer than the body, and that in the middle of a red colour. It frequents the meadows and gardens ‡.

^{*} Derham's Physico Theol. 1. 8, c. 6.

[†] Vespa Ichneumon tota nigra, Rai Ins. p. 256.

M. de Geer, Vol. I. p. 703.

GENUS V .- Sphen. The Savage.

THE mouth of this tribe is armed with jaws, but has no tongue; the antennæ have ten articulations, and the wings are extended, and laid horizontally upon the back; the sting lies concealed within the body.

This genus contains infects, perhaps the most fierce and rapacious of this class of beings. The favage, or ground bee, does not feed upon honey, or accommodate its young with that kind of provision. They attack infects much beyond their own fize, and that whether they are defenceless or armed; for they are provided with strong jaws, and a sting poisoned with a liquor fatal to every animal they engage. The favage feizes boldly on the infect he attacks, and gives it a stroke of amazing force. After this first encounter, he falls down, as if he had himself received the mortal blow; but it is only to rest from his fatigue, and to observe the effects of his prowefs. By and by, the wounded animal dies; and while yet palpitating with life, the favage devours those parts which he finds most palatable, leaving the greater part entire.

It is thus that the fphex riots in the blood of hundreds of intects; and his family is no fconer increased, than the carnage becomes proportioned to the number of young he has to support. After the female has deposited her eggs in the bottom of a cell dug in the ground, or in the mud

mud wall of a cottage, the whole apartment is crammed with multitudes of living and dead infects, destined to be the food of the future progeny. Thus their houses, like the renowned caves of the giants, are strewed with dead. This operation is no sooner over, than the parent infects stop up the hole at the entrance of the cell, to prevent the escape of such of the wretched captives as may yet be alive. The young, when they leave their eggs, find themselves amply supplied with provision. They devour, one after another, all the carcases with which they are provided; and when their last sty is eaten, they have no longer occasion for food, but are changed into chrysalids, which afterward become savages of one species or another, according to that of the parent pair *.

Sphex femi aurata viridis †. The head, thorax, and abdomen of this beautiful species are of a finely gilded blue; the antennæ are yellowish; the eyes of a gilded brown, and legs of a dark blue.

The spirifex, or turner savage, is of a chesnut brown, with a shade of blue; the eyes are large and black; the antennæ brown; the body is ferruginous; the thorax and abdomen connected together by a yellow thread; the wings are of a dusky brown, and the sting yellow 1.

^{*} Reaumur. Tom. VI. mem. 9. † Syst. Nat. Ord. V. Gen. v. sp. 336 & Barbut's Gen. Insect. p. 253.

GENUS VI .- Chrysis. The Golden Fly.

This tribe, like fome of the preceding, has no probefcis, but a mouth furnished with jaws: The antenuæ are filiform and bended, consisting of several articulations, of which the first is the longest; the abdomen is arched underneath with a scale on each side; the anus is dentated, and armed with a projecting slive; the body is covered above by the wings, through which it shines as if gilded *.

Among this family the ignita, or flaming chryfis, flands distinguished by its resplendent colours. The fore part of the head is green and gold, and the hind part of a beautiful azure; the thorax is likewise azure, with a mixture of green: The abdomen is green and yellow before, and behind of a copper red, resembling that metal when highly polished †.

^{*} Syst. Nat. Ord. V. Gen. 6. † Barbut, p. 256.

GENUS VII.-Vespa. The Wasp.

There are no less than twenty-eight species of the wasp; many of which are universally known. The common characters by which the tribe is distinguished, are, the jaws with which the mouth is armed; the four wings, the first pair of which are folded, and the sharp pointed sting which is usually concealed within the body; the abdomen is joined to the thorax by a short stalk, and the whole body is smooth *.

The crabro or hornet is not found in Scotland, though common in South Britain. The antennæ, head, and legs, are of a brown or chefrut colour: the abdomen is of a fine orange brown; but on the extension of the annuli, it discovers on each side a line of black: the wings are of the colour of amber; and the animal, for size, merits the first rank among the vespæ. It builds its nest in the body of decayed trees †.

Vefpa vulgaris, the common wasp. The thorax of this species is black, surrounded on the anterior part by a yellow line; the abdomen is of a golden yellow, having triangular spots down the back part, and black ones on each side.

The wasps in general feem to fill up a middle rank between the ichneumons and the bees; like the former, they

^{*} Syft. Nat. p. 984.

they are rapacious and carnivorous, and, like the latter, they construct hives, and sometimes feed on the produce of flowers; they devour fruit, butcher meat, and carry on continual hostilities against almost every other species of sly: they are at once the rivals and the enemies of the common bee; many of which annually perish by their attacks.

Almost every person must have seen the establishment made under ground by the common wasp. It is a kind of subterraneous city, which at certain seasons of the year contains many thousands of inhabitants, and is constructed nearly with the same ingenuity and elegance as that of the domestic bee. Like it, it is internally formed with combs, consisting of a number of hexagonal cells, all enveloped under one common covering, like coarse paper, which is constructed with great art. In this particular they excel the common bee, which contents itself with the cover afforded by the hive, or with the trunk of a rotten tree, in their wild state.

Though the wasps generally make choice of some large hole under ground for the construction of their nest, they have nevertheless much labour to undergo in removing protuberances, and carrying away earth, till it is brought to that spherical sigure which suits their purposes. This work completed, they next construct that paper like covering with which the whole hive is lined. The combs in which the cells are lodged next claim their attention: These are ranged horizontally in different stories, sometimes twelve or sisteen above each other, all supported by colonades, between which the whole citizens of this subterraneous commonwealth are seen at times to walk, like men in the streets of a town. The cells of the wasps are

nos

not confiructed with that geometrical skill which has been so often admired in those of the bee; but they are not on that account the less adapted to the purposes they are destined to serve. Each comb has only a single range of cells, with their mouths opening below. They are intended, not for the reception of honey, but for the habitation of the young, which are fed twice or thrice a-day, by the morsels carried in by their parents. For the more commodious reception of their food, each of the larvæ has its head turned downward opposite to the mouth of its cell, ready to receive its meal when offered *.

There are, however, many varieties in the construction of wasp hives, all suited to the views of the different species who inhabit them. Some have only a single row of cells, placed vertically, like those of the bee, and the mouths facing the sun: The reason of this variety seems to be, that some kinds require the heat of the sun to hatch their eggs; an advantage which could not be obtained, were there more rows of cells, or were they placed in a different manner.

As in a hive of bees, fo in that of wafps, there are three different kinds of animals: At certain feafons, it contains only a fingle female, and a number of neuters or mules, who are of no fex; at other times it contains fome hundreds of females, and still a greater number of males. The former are of a fize fo superior to the mules, that one weighs against eight; even the male wasp is not more than half the weight of his female.

The condition of the female wasp differs widely from that of the female bee: The latter, when she leads from Vol. III.

^{*} Reaumur, Tom. VI. Mem. vi.

the parent hive a young colony, and founds a new emepire, is effentially a queen; for the is more implicitly obeyed than the best of monarchs, by the most loyal of his lieges. The female wasp, as her station is more laborious, so her genius seems more enterprising. Unaided by any of her kind, she lays, in the beginning of every season, the foundation of a new edifice, which is destined to be the birth place of many thousands of her species. She constructs the first cells, to which the commits the earliest of her eggs, which in time become mule wasps, the most active and laborious of the whole race; and by these she is assisted in completing the rest of the work.

The male wasps are not so slothful as the male bee; they discharge several duties in the interior parts of the hive; but in the art of building, either the cells or the external covering, they are altogether unskilled. This falls to be executed by the mules, who carry it on with amazing dispatch. They collect together the small sibres of half rotten wood, which they moisten with a glutinous substance, and bake up into that paper of which all their work is formed.

The aerial wasp constructs small nests of about the size of an orange, which it attaches to the branch of a tree. This compact little edifice is exposed to the weather, but is rendered impenetrable to rain, by a number of leaves which are placed round it, exactly resembling an inverted rose. A particular species in the neighbourhood of Cayenne, construct a large oblong box, about twelve or sisteen inches long, of sine parchment, which is also pendent from the branch of a tree; there the combs sabricated of the same substance, are ranged horizontally, in different stories, one above the other; each having a round

round hole in the centre, by which the wasps are enabled to ascend or descend to different flats of their building *.

In the mean time, while these operations are carrying on by the joint labours of the hive, the mother wasp continues to lay till she has produced fifteen or fixteen thoufand neuters or mules, and about five or fix hundred males and females: The commonwealth day by day increases in numbers, and enjoys peace. Towards the month of October, provisions begin to grow scarce, and a new scene ensues: This hitherto amicable tribe seems then fired with mutual rage; and the whole edifice prefents one scene of massacre. The neuters and males tear from their cradles the eggs, the larvæ, and new born infects, with undiffinguishing fury. They next fight one against another. Hopes of the state, folicitude for posterity, or love to their native place, now no longer exist; the whole commonwealth is shaken to the foundation. Rains and frosts ensue; the citizens are seized with disease and languor; and happily for the other insects, and the fruit gardens, almost the whole die. Some few females escape from the disasters of civil war, and the feverity of winter, and again in the ensuing spring become the founders of new empires, which are again to be overturned +.

[·] Reaumur, ubi supra.


[†] Barbut's Gen. Insect. p. 262.

GENUS VIII .- Apis. The Bec.

WE are now come to a tribe of the most useful of in fects, whose history has been the subject of many volumes. Almost every writer on entymology has made the manners and economy of these animals a considerable part of his work; and many authors have treated of them, who have entered into no other department of natural hithory *. Several of thefe, catching that enthusiafm which is natural in contemplating creatures whose instincts are truly wonderful, seem to set no bounds to the eulogies they have lavished upon them. If you believe them, there is hardly any fort of intelligence or moral virtue, which bees do not posses; and the whole of their manners are such as afford us but too just causes of blushing at our own. The celebrated Reaumur, in detailing the economy of these animals, an object to which he applied himself with a perseverance superior to all mankind, has abstracted much of the marvellous from their history; but in return, he has enlarged it with many facts and observations formerly unknown, and which fucceeding experiments have feldom contradicted: The following brief account of these insects, in general, rests upon his authority.

There

Among the most judicious naturalists who have written upon this stripiest, we may rank the following, viz. Mousset, Merian, Aldrevandus, Johnston, Charleton, Ray, Dale, and Reaumur.





There are, at certain feafons of the year, three kinds of bees in every hive; the males, the females, and the bees without fex. The latter every perfon is acquainted with; their number is beyond comparison greater than that of the other two kinds. Nature seems to have destinct them solely for the purpose of labour; and the whole drudgery of the hive lies upon them; hence they have properly been termed Working Bees.

It is only during one or two months of the fummer, when the hive is most crouded, that males are found in it; and even then, they do not amount to a tenth part of the whole: They are of superior fize, and distinguishable by the characters afterwards to be noticed. During the whole course of the season, except a few days, there is only a fingle female to be discovered in the most numerous hive. Her fecundity, however, is fo prodigious, that she is soon capable of multiplying her family to such a degree, that the hive can no longer contain it. To her the whole fwarm, amounting from twenty to forty thousand, owe their birth. Her residence is generally in the interior apartments of the lodging; when she shews herself, she is readily known from her fize, being longer than even the male bees, but inferior in thickness. It is this female whom the ancients dignified with the title of King of the bees, and whom, with the utmost propriety, they might have denominated their Queen. From a number of well attested experiments and observations, it appears that the common bees pay her more than respect. They continually endeavour to be useful to her; and by carreffing her, and offering her honey, they feem to ansicipate her wants. Her life is more precious than any

of the rest; wherever she goes, therefore, she is encircled by a guard, that continually pays court to her *.

It is not without cause that this sedulous attention is paid to the mother bee; she is the foul of all their operations. If a hive is deprived of her, however numerous, it will undertake no labour; and the individuals will hardly give themtelves the trouble of collecting their daily fublishence. A fwarm that was bufy from morning to night constructing cells and collecting wax, immediately upon this accident feems to forget that the flowers contain their food; they scarcely stir from the hive, confruct no new cells, nor even finish fuch as were begun. The moment she is restored, their wonted spirit and activity is refumed by the whole fwarm: Not only do the working bees refume their labours when the queen is restored, but they ply them with an assiduity proportioned to her fertility. Although they contribute nothing themfelves to the generation of the young bees, the Author of Nature feems to have given them the same affection, and to have interested them in their welfare, as much as if they had been their real parents.

The external parts of bees are happily accommodated by their shudure, to the delicate operations they are destined to perform. Each of them is surnished with a trunk or proboscist, commonly solded up, but capable of being extended at pleasure: It is with this instrument that they collect their food; not by pumping or sucking, but by licking it from the nectaria of slowers; vessels that have been but lately discovered by the botanists, but with

* Reaumur, Tome V. Mem. v.

[†] This finall inftrument is faid to confift of no lefs than twenty different parts, perceivable by the microscope. Resumur

with which the bees have been well acquainted from time immemorial *.

The teeth with which these insects are provided, serve another purpose equally important; they are the instruments by which they falhion and give a proper confiftency to the wax. That valuable commodity is not found in a perfect flate upon the flowers, as many have imagined; it is there feattered upon the furface in the form of a fine powder or dutt, and fwept off by the hairy legs of the bee, and deposited in a hollow part of each thigh, prepared for its reception. After being carried in that form into the hive, it is all ate up by the bee; and by the action of the flomach is brought into the state of genuine wax. From the stomach, the working bee brings it back in small parcels to the mouth, like a ruminating animal; and, by chewing it there, fashions it into pieces proper for the construction of the cell, to which it is applied, and afterwards polished as the situation requires.

That fmall fealy spine, commonly termed the sting of the bee, is only the case of two needles or darts, extremely sine, and each dentated towards the point. The wounds made by these slender arms would be little to be apprehended, were their points not impossoned by a small drop of acrid liquor. This liquor, which, when tasted, burns the tongue, is conveyed along a small canal to the case of the sting, on the tip of which it appears in small drops, when the bee intends to make use of her offensive weapons. However disagreeable this instrument may sometimes prove to us, it is absolutely necessary to the bee, surrounded as that animal continually is with many enemies,

^{*} The diligence of the bee is often alluded to in ancient poetry.

Apis manting more moded, grata carpentis thyma, Hor.

enemies, whose hostilities are constantly provoked by the honey and wax.

The sting of the bee is also used for another purpose, the propriety of which we are not so well able to vindica. At a certain time, they are busy from morning to night in no other employment than massicring their associates. The males, it seems, after the semale is secundated, become useless, and even offensive to the rest of the hive. The working bees, who had formerly been their nurses, and hitherto had lived with them in the best understanding, all at once break loose upon them with unrelenting sury, and in two or three days destroy the whole in one general carnage †.

Of the reaf ns alleged by the working bees for this maffacre, we are altogether ignorant; we know not upon what claim their power of life and death over the males is founded, faither than that nature feems to have granted fuch a right, by giving them power to exercise it. Thefe, however, are not the only combats in which the bees are engaged: The working bees of the same hive often quarrel, and challenge each other to battle. Long do they struggle in defence and attack, like two skilful pugilifts, the one endeavouring to find a place in the fealy body of his adversary into which he may thruit his fting, and the other as studiously warding off the blow. The first wound that the puts an end to the engag-ment; and the victorious see walks off, leaving his adversary to expire in the dust. Sometimes three or four attack a fingle bee, without any defign upon his life, but with a view to force him to difgorge his honey; and

as

[†] Virgil was acquainted with this peculiarity. Fucos a presapibus arcent ignavum pecus. Æn. Lib. ii.

as foon as they have succeeded, they lick it up, allowing the former proprietor to walk away.

Battles far more fatal and general are occasioned when a neighbouring swarm, from poverty, or a principle of injustice, invade a hive already occupied. Scarcely have they entered the walls of the city, when a bloody engagement ensues: Those who have the right of possession oppose their invaders with all their forces, and with undaunted courage; not a minute passes that you do not observe a victorious bee dragging to the door of the hive a dead adversary, or one who is yet struggling in all the agonies of death. These engagements do not close but with the day; and before victory declares for either party, they often cost thousands their lives; for very often the one who has stung its opponent leaves its weapon in the wound, an accident which proves statal to itself.

Infects of their own species are far from being the only enemies which the bees have to fear; worms, wasps, hornets, and infects of different kinds, never fail to make their way into the hive, wherever any rent or crevice is left open. When attacked by these robbers, they perish in the unequal combat; and when dead, their bodies are ripped up, in order to extract from them the honey they contain. Apprised of the fatal consequences of admitting them into the hive, the bees carefully fill up every chink and crevice, not with wax, but with the glutinous matter that exsudes from certain trees, a substance still more tenaceous. When a bee enters the hive loaded with a quantity of this stuss, it is met by others, who take small particles of it, and apply it to the sides of the part to be stopped up, till they are entirely closed.

A fwarm of bees, however numerous, as we have alteady feen, all owe their birth to a fingle female. This Vol. III.

queen, which Virgil, and the other writers of antiquity, have charged with all the cares of government, is indeed bufily occupied, but in functions of a different kind; and these are, the production of a valt number of eggs, which the continues to drop, one after another, into the empty cells, during a confiderable part of the fummer. This animal, which is fo amazingly productive, on being opened, has been found to contain upwards of five thousand eggs, all of a fize fufficient to be perceptible. If we make allowance for those that were already dropped, and many more not yet formed fo as to become perceptible, we shall no longer deem it incredible, that this animal thould in one feafon become the mother of fo many thoufands. The most numerous hive is far inferior to the number of spawn that have been taken from the belly of a fmall fish. During the whole time that the female goes from cell to cell depositing her eggs, the is accompanied by the working bees, who tend her with the most officious care. And as the males and females are produced by her of a fuperior fize to the reft, the working bees, as if appriled of the circumstance, construct a few cells larger than the rest; and what is still more remark. alle, the female herfelf knows which of the embryos are to become of her own fex, and accordingly deposits them in cells of a fuitable capacity. The cell which is to contain the future queen bee is of a thructure different from all the rest; and in building it, the labourers abandon the hexagonal shape which is best adapted to the faving of labour, and of wax; nothing, in this instance, feems to coll them too dear: The chadle of their future queen confumes as much as a hundred or a hundred and fitty of the ordinary cells.

From

From the great number of males in each hive to a fingle female, the ancients were induced to believe, that secundation among these insects was not accomplished by any act of copulation, but that it was performed by a vivifying liquor shed upon the ova in the cells: But this opinion was overthrown, as foon as it was observed, that for nine or ten months in the year there was no male in the hive. Swammerdam, who could not observe that in these animals there were any parts of generation +, therefore supposed, that the semale was impregnated folely by the effluvia of the male bees. The observations made by Reaumur have removed those difficulties with which this part of the history of bees seemed to be attended. The female, who is furrounded by fo many males, instead of being fatigued and perfecuted by their importunities, is obliged to carefs and inflame the passions of these drones, who are the most cold and indifferent of all their fex. She mounts upon their backs, and pushes her gallantry to what may be deemed indecency; at length her allurements produce the defired effect, and copulation is performed.

The female thus impregnated proceeds to lay her eggs, which are round and oblong, rather thicker at the one end than the other. Each is deposited in a separate cell, except upon some occasions, when the industry of the working bees has not been sufficient for the sertility of the mother: She then lays two or three in a single cell; the supernumeraries being afterwards carried away and placed by themselves, as soon as new cells are provided. The single egg which remains is saftened to the bottom of the cell by the small end, the only part which seems

3 T 2

to touch it. In a fingle day or two after being placed there, it produces a worm, which the working bees take care to supply with suitable food, confishing of a whitish liquor, which serves it at once for substitutence and a bed; for it lies upon it, folded up like a ring. In the space of fix or seven days more, the whole of its growth is completed; and the bees, who know the time at which it no longer needs to be supplied with food, cease to carry it any more. The last office which they render it, is closing up the mouth of the cell, which is done with wax.

In the mean time, the young animal within, which hitherto was almost entirely inactive, begins to spin filk, and line the infide of its habitation, as a preparative to its entering into the chryfalis flate, in which it for a while remains. Thus, in the space of about three weeks after it was first dropped into the cell, the young bee is ready to make its appearance as a winged animal. The first employment, after it has arrived at that period, is to gnaw off the wax with which the mouth of its cell had been flopped up. On its flut iffuing from the cell, the whole body is wet with the humid fubflance of the cell; but the affictionate bees flock around, and with their trunks affait it in wiping off the moitture. The wings grow dry, and the limbs firm; and in the course of the fame day, it islues forth with the labourers to collect wax and loney, and thereby to make returns to the fociety for the care with which it has been reared.

Thus, after the commencement of fummer, the number of bees daily continues to increase, till, by the month of June, they have become so numerous, that the same hive can no longer contain them. The only measure which

then remains to be adopted, is, to divide; and in an inflant, a large colony prepares to abandon for ever the place of their nativity, and to follow a young queen in forming a new establishment for themselves.

It is not merely the number of bees in a hive that determines them to fwarm; that event depends on the young female, who must not only be sufficiently strong to lead the colony, but must also be impregnated; for upon that circumstance depends the whole hopes of the future progeny: But no sooner do all these predisposing circumstances concur in a hive, than it prepares immediately for colonisation. In the evening before that eventful period, indication of their intentions is given, by an unusual noise and hurry within the hive. All the morning of next day passes without hardly a single bee going to collect honey. Those who are to leave the hive are preparing to take their departure; while those that remain behind defer their operations, till their companions, by their leaving them, have assorted sufficient room.

It is in the great enterprife of colonifation that the influence of the queen bee is most clearly seen: Wherever she alights, there the whole swarm take up their abode, and all cling around her body. If she is removed, they all sly in quest of her: if put into a hive where she has not been placed, they either forfake it or die. Hence the art of those persons is explained, who go about shewing bees, making them alight on different parts of their body, and sollow them for any length of time. They have possession of the queen bee; and by that means can influence all their movements.

When a fwarm thus migrates from the parent hive, it frequently happens that more than one female accompa-

nies it: The prosperity, however, of the new colony requires that there should be but one; one only is accordingly preserved. In a single day, all the rest are put to death; and it is found that the one which is spared is the best entitled to reign, because she is most fertile, or nearest her period of giving a new progeny to the society. The young semales in the old hive, if any remain there, share not a more happy destiny; all the supernumeraries, like those in the swarm, are put to death.

We have already remarked that the bees have many enemies; and in some feasons, we are told, they are liable to satal diseases: but of all their destroyers man is the greatest, from that barbarous avidity which gives rise to the practice of killing every hive that has collected a sufficient quantity of honey*. In the winter scason, cold or hunger are the scourges which often destroy such as our rapacity has spared. In sheltering them from the one of these evils, we often expose them to the other. Too great a degree of cold makes them perish †: If placed in a situation too warm, they are awakened from their torpid state, and devour their food with an appetite which seen brings on a samine. Experience is the only guide that can prevent the bee master from falling into these extremes.

In Egypt and in Greece, where the collection of honey was deemed a trade of importance, the hives were regularly removed to a different part of the country, when the flowers had faded in another. It was for this purpose,

^{*} Reaumur advises the French government to inflict an arbitrary punishment on all who fmeaked bees. Tome V. p. 666.

[†] The ancients were of opinion, that bees benumbed with cold might be reflored by hot affice, Vid, Columbia & Varo.

pole, that boats flored with hives were fet affect on the Nile, gradually bringing the infects within the reach of fresh food as the old was exhausted. A similar method was some time ago practifed in France by a person who had six or seven hundred hives in his possession †.

Such are the outlines of the hittory of the domestic bee, an infect to which we are indebted for two material articles of our enjoyment, wax and honey. The genus of infects to which it belongs is distinguished into firty-five different species, all distering in their manners, as well as their external form. Their common characters, in which they all agree, are, a mouth furnished with jaws and a proboscis, inclining towards the body, and inclosed in a sheath; wings which are extended, and without folds; three stemmata on the head, and stings carried in the other extremity by neuters.

Apis mellifica, the domestic neuter bee. This animal is fo generally known, that no description of its parts is necessary.

Apis mas, the male bee. This species has no string, nor are its feet or rostrum sitted for the collection of wax or honey: it is larger than the common working bee: it hovers upon slowers; and its only office is impregnating the female.

Apis regina, the queen bee. This animal is eafily diflinguished by her great length of body, and the shortness of her wings, which the never uses but when employed in leading out a young swarm.

Apis museocum, the humming bee †. The three former infects, as they breed together, may all be deemed of

the

[†] Reaumur, Tome V. Pref. p. 40. Syft, Nat. Ord. V. Gen. vi i. fp. 46.

the same species: this is larger in size, and more variegated in colour: It constructs its nest in a hasty manner, of piles of moss, collected in the meadows. The vaulted roof proves a security against rain, and the slooring of moss below preserves the nest from damps.

The female bee collect, unwrought wax, and a species of honey, in a few cells coarsely constructed, and there she lays her eggs; the hive gradually increasing, till it contain sifty or sixty bees, which, by the frosts and rains of winter, are almost wholly destroyed. A few impregnated females survive the severity of the season, and build new nests in the spring, which are to be the cradles of new hives. Field mice, hornets, and ants, often plunder from these industrious animals the sixtle vessels of honey which they had laid up for the winter †.

[†] Barbut, p. 268.

GENUS IX .- Formica. The Ant.

THE characters of this remarkable tribe are, a small erect scale between the thorax and abdomen; truncated or broken antennæ, having their sirst articulation longer than the rest; the semales and neuters armed with a sting, which is concealed in the abdomen. The males and semales have wings, while the neuters are apterous *.

In the System of Nature, there are enumerated eighteen different species of the ant; of these, the largest is the hippomyrmex, or horse ant, which is found in the hollow trunks of rotten trees. The head is black, and the thorax ferruginous; the seet are of the same colour, while the abdomen is brown: The erect scale between that part and the thorax, is oval, entire, pointed, and round †. The other animals of this genus most common, are,

The larger and smaller red ants. These are found in woods, or under heaps of dry earth, and are the most laborious and industrious of the whole insect tribe: under these characters they are frequently alluded to by almost every writer of antiquity.

Vol. III. 3 U The

^{*} Systema Naturæ, Ord. V. Gen. ik.

[†] Vide Rai Inf. p. 79.

The ants nest is in fact a small well regulated commonwealth, resembling in many respects that of the bees. Like them, they are continually employed in labour, and have a large proportion of the citizens neuters, whose solecting food for the society. The males and semales have wings, and seem destined for higher enjoyments, as well as more noble occupations. Their operations are all conducted in peace, union, and good understanding; though, if we are to credit the wise man as a naturalist, they have no guide, overseer, or ruler *.

When they have chosen their ground for forming an establishment, which is generally some dry and sunny exposure, all are employed in digging their retreat, to the depth of about a foot or more: They seem to take neither rest nor food till it is completed; and to prevent disorder or consustant, their police assigns to every one his task. While one loosens the earth, another carries it away, and is met by others returning for a new load.

Within this hollow cave, when formed, these insects all live in society, and shelter themselves from the severity of winter. Even after the editice is constructed, there is not a day allotted for rest, so long as the weather will permit them to go abroad. They are continually wandering about in quest of food, and carry all day each his burden into the nest. Should one chance to meet with a booty which is too large for him to remove, he is soon perceived by the rest, who come in legions to gnaw it into smaller pieces, or to assist in rolling along the unwieldy

^{*} Vide Solomon, Prov. chap. vi. 9. Go to the ant thou fluggard, confider her ways, and be wife; which having no guide, overfeer, or ruler, provideth her meat in the fummer, &c.

weildy mass. Corn and feeds form an acceptable food; but as they are also carnivorous, they devour frogs and lizards when delivered over to them, or unable to defend themselves.

Later naturalists have afferted, that all the food collected by the ants during the day, is deposited in the common hall, and devoured every evening, or distributed to the young; and that, in winter, they pass into a torpid and benumbed state, in which they are incapable of taking any food *. This, however, is contradicted by the voice of all antiquity. Horace recommends its example to the miser, as being noted for liberally enjoying during winter, whatever had been amassed by its own industry in the mild season †.

We confess our incapacity to decide this matter; but having been unable, after all our researches, to find out the ant's winter store, we are rather inclined to think that in that season they either use none, or are satisfied with what they can occasionally procure.

The eggs of these insects are deposited pretty early in the spring; and in order to provide the necessary warmth for their young, the old ants are seen carrying them out to expose them to the heat of the sun. The worms go into chrysalids, and are metamorphosed in their own skin. When about to issue forth into new life, it tears this white covering, and comes forth a persect ant. The ants

3 U 2

copulate

Parvula, nam exemplo est, magni formica laboris
Ore trahit quodcunque potest, atque addit acervo
Haud ignara, ac non incauta suturi.
Quæ, simul inversum contristat aquarius annum
Non usquam prorepit, & illis utitur ante,
Quæssitis sapiens.
Lib. I. Sat. i. l. 333

^{*} Earbut, and many other naturalists.

copulate in the air; and the females, who are the fmallest of the race, seldom repair to the common habitation: What is their destiny, whether they die by the severity of the winter, or fall a sacrifice, like the bees, to the merciless sury of the labouring ants, is not ascertained.*

We have already had occasion to remark the fondness of these insects for the saccharine liquor that exsudes from the plant louse: it is in quest of that food that they are frequently seen on plants, in the company of these insects. In Switzerland, however, they are transported to trees for a different purpose; the destruction of caterpillars, and other vermin. A bag filled with ants is sastened to a tree, with a small hole purposely lest open for them to creep out; they spread along the tree, and are prevented from leaving it, by a quantity of pitch with which the stem is covered. Rather than die by samine, they go in pursuit of the caterpillars among the leaves, and devour them.

Barbut's Gen. Insect. p. 280.

GENUS X .- Mutilla.

A FEW only of this tribe are winged; in those that are provided with these instruments, they are resplendent like pearl, and laid horizontally upon their backs. The antennæ are broken, and have the first articulation longer than the rest; the body is covered with down; the thorax is obtuse at the base; and the sting is concealed within the abdomen *.

The common residence of the mutillæ is upon the ground, below moss, especially where there is a hollow space between the moss and the earth. They are gregarious, and when disturbed in their retreats, they make their escape by running very swiftly. The mutillæ, for the greatest part, are inhabitants of foreign climes: that most frequently seen in *Britain* is,

The mutilla Europea. The thorax of this infect is a deep yellow, mingled with black velvety down; the prevailing colour of the abdomen is black, but is adorned with two, and fometimes a greater number of yellowish white bands; the antennæ are black, and bending; the legs of the same colour, and hairy; the cyes are of a beautiful pearl white, shining from a head of a coal black.

^{*} Syft. Nat. p. 966.

ORDER VI .- Depterous Infects.

This order comprehends all the numerous races of flies, tipulæ and gnats, which are characterised by having only two wings: Below each of these there is placed a halter or posser, which is terminated by a knob, and has its base concealed under a little scale *.

* Syft. Nat. Ord. VI.

GENUS I .- Oestrus. The Gad Fly.

This infect has no mouth, nor any visible proboscis: in place of these are three small points. Upon the head, there are three stemmata, or small eyes; the antennæ are setaceous, and commonly short.

Oester bovinus, the ox gad fly. This large species penetrates through the skin of cattle, and deposits its eggs there. It has no mouth, black eyes, a yellow thorax, and pale-coloured feet; the wings are membranous, and interspersed with small black spots *.

Oestrus rangiferus. This species frequents the backs of rein-deer, especially those that are tamed, and deposits its eggs there; it is twice the fize of the former, and claded by the Laplanders curbmat. The wings are without spots, and the body is covered with a yeilowish down.

Oestrus ovis, lodges in the noses of ruminating animals, particularly of the sheep, and is therefore called by the Swedes, noosmalsken I. To this tribe is also added the cestrus hæmorrhoidalis, which makes its way into the anus of the horse, in order to deposit its young §.

In the economy of nature, there is nothing appears more furprifing than that inflinct by which certain animals are directed to deposit their young in the only place where

^{*} Rai Inf. p. 271, & Derham's Physico Theol. l. 8. c. 6.

[†] Fauna Suec. N. 1025. ‡ Reaumur, Tome IV.

[§] Voyage de Gotlande, 277.

where they can find fuitable food, and necessary warmth. The young of certain slies are only capable of subsisting below ground, and of deriving food from a certain slower, the narcissus. The mothers of these deposit their eggs in the roots of that plant. Other slies enter the anus of a horse, because their young are only capable of subsisting in the entrails of that animal, where they remain till they are about to be metamorphosed, and are then ejected with the dung *. Others are provided with a curious instrument, by which they perforate the hides of cattle; and in each hole thus made, they deposit an egg.

Each of these worms, after being hatched, finds itself surrounded with its proper food, among which it grows; and the cell increases in fize at the same time, till it sometimes reaches the bulk of an egg. The wound becomes silled with purulent matter; and the opening from which it runs serves to supply the insect with air. After the worm has grown there to its full size, it needs a place more dry, and of less heat, for its metamorphosis; it then drops from the orisice of the wound, and betakes itself to some hole, or other retreat, where it is transformed.

An inftinct as fingular as any of the former is that by which fome flies are directed to the note of a sheep, a goat, or a deer, to lay its eggs. It is in the frontal sinus of these large animals, that the worms produced by these eggs are destined to search for the only food capable of supporting them. Some of these worms, after being ejected from this strange nidus, become covered with a hard shell, in which their last change is completed. Nature, which provides for every exigence, has surnished

this

^{*} Reaumur, Tome IV. Mem. xii.

this shell, which is of a texture too hard for the animal to break, with a valve at one end, fastened only by a small thread. At the sirst effort for freedom which the young æstrus makes, the door gives way, the prison opens, and the insect slies to those woods and meadows that were frequented by its ancestors *.

* Reaumur, ubi fupra,

Vol. III.

3 X

GENUS II .- Tipula. The Grane Fly.

In this tribe, the mouth is a prolongation of the head; the probofe is is fhort, and bent inwards; the upper jaw being arched: All the different species, which are no less than sifty-eight, have two palpi, which are incurvated, and lenger than the head *. The whole genus is readily distinguished by the enormous length of their legs, by which they are enabled to waik among the grass, as men are affished by shits in passing through marshy ground.

The tipula maxima of Ray is one of the largest seen in this country: The wings are variegated with brown and white; the body and abdomen are of an ash-coloured brown to

Tipula crecata, the faffron crane fly. This elegant species has the thorax and scapuli of black hue; the belly staffron: the wings are marked each with a smooth resplendent spot of a brown colour; the rest of the wings are yellow, and the annuli of the abdomen are distinguished by alternate bands of yellow and black ‡.

The ash coloured tipula is mentioned by Goedart §, I.ewenbock ¶, and Reaumur ||. The whole body is of an

^{*} Syst. Nat. p. 970. † Rai. Insect. p. 72. No. 2. * Actes d'Upsal, 1736, p. 30. § Part II. p. 165.

[¶] Epist. 1693, Dec. 20. | Tome IV.

affi-colour; the wings of a water-colour, having a brown longitudinal bar upon their exterior parts. It is found among garden plants.

The larger tipulæ are known by the name of Sempfireffes; the finall by that of Culiciform. The latter, in fine fummer evenings, flutter about the fides of rivers in legions. The shrill noise which they make with their wings is not very discernible unless they are very near. They are sometimes taken for gnats; but they want their proboscis, and their manners are far more inossensive.

The tipulæ, before they become inhabitants of the air, creep in the form of grubs: Those of the larger kinds dwell in holes of decayed willows, where they change into chryfalids. In that state, they have the power of breathing through fmall curved holes. The larvæ and chryfalids of the fmaller tipulæ are generally found in water; they are extremely various both in colour and form: Some have cylindrical tubes through which they breathe, even while confiderably immerfed in water, by ftretching their points to the furface *. They fwim with great nimbleness, but feldom forfake the hole they have dug in the bank of the river. There are others of these animals that spin a silken coat, that receives part of their long body before their transformation. After these operations are performed, all of them renounce their reptile and aquatic life, having received inflruments of flight from the hands of nature. Their frame is then fo weak, that a touch is enough to cruth them. They are fometimes of a beautiful green, fomecimes coal black; but the most remarkable are those whose fore legs, though extremely long, do not touch the ground, but move in the

air like antennæ. In this their last and most perfect state, the tipulæ being provided with proper organs, apply themselves to the important work of propagation; and having provided for another generation, their short existence draws to a close *.

Barbut, Gen. Inf. p. 294

GENUS III. Musca. The Fly.

This numerous tribe is characterised by that fost and sleshy proboscis which forms the mouth, and by being destitute of palpi. There are about an hundred and thirty different species of slies already known to naturalists, who have divided them into two grand sections; those with sliated and simple antennæ, and those whose antennæ are surnished with a lateral hair or feather *.

Nothing fo clearly demonstrates the imperfection of the science of entymology as the small number of flies that have been hitherto described by naturalists. The System of Nature in its last state of enlargement, was supposed to contain the most accurate and complete enumeration of infects that has yet been published: Mr. Harris, however, in his Expedition of the English infects, describes a greater number of flies indigenous in this country, than the Swediff paturalist had discovered on the whole face of the globe. England is at the fame time far from being the most productive country of these animals. In Spain, they were formerly fo numerous, that the fly-catcher, we are told, was not only a profession, but a number of men were commissioned by government to give chace to these troublesome invaders. In all hot climates they multiply to a degree almost intolerable,

and

^{*} Syft. Nat. Ord. VI. Gen. iii,

and are extremely destructive to butcher meat, and other kinds of provision.

Next to the number of flies, their variety claims our attention. The different species are extremely diversified in their external form, their structure, their organization, their metamorphosis, their manner of propagating their species, and in providing for their posterity. A full explanation of these different parts of their economy, would require a large volume, and would constitute a narrative to many readers not uninteresting. Such an undertaking, however, is incompatible with the limits prescribed to our work.

Some of these insects have trunks instead of a mouth; others have that organ armed with teeth; and many have both a mouth and a trunk. The probose of sies is a machine contrived for pumping the blood from the vessels of large animals, and the nectar from the petula of slowers; and the science of hydraulics has not enabled men to construct machines more complicated, more exactly arranged, or better adapted to these purposes. Some of them possess considerable sirmness and solidity, those especially that are destined to pierce the skins of cattle. In order to see them at work, all that is necessary is to expose a syrup to attract them, and to take up a lens for their inspection.

Each eye of the fly contains in it an assemblage of a great number of small ones, which probably have the effect of multiplying the surrounding objects, and creating representations of them, which the experience of the infect corrects *. From the eye, if you pass along and sur-

respiration. Four small stigmata, for this purpose, are found upon the thorax, and a greater number dispersed over the annuli of the abdomen. The covering of the different parts of the abdomen are scaly, and do not consist all of one piece, but admit of the contraction and dilatation of that part, according to the circumstances of the animal. Each of the feet terminates in a small bunch of seta, resembling a brush.

The interior organization of this genus, when examined with a proper apparatus, presents two pulmonary sacs of a white colour, arranged longitudinally along the body. As the bodies of fome species are diaphanous, the action of the heart may also be discerned, as well as the liquor it contains, continually driven along the great artery leading to it, and returning by the fame course *. During their larva state, the slies in general go through a greater variety of metamorphofes than even the filk or butterfly worms: During their progress from the vermicular to the chryfalis frate, they pass through an intermediate change, unexperienced by other infects. From the shorts ness of their lives, all these vicifitudes must rapidly succeed each other: this circumstance, however, does not prevent many species from constructing a coque for their last metamorphosis, of a curious fabric. The greater part of these coques are of filk, with a mixture of other materials; thefe, indeed, most common with us, ferve themselves with their own skin, which becomes an incrustated covering to protect them during their dormant and aurelia flate.

When

When the different members have acquired fufficient firmness, the fly is ready to burst from its shell in its winged form. The covering of the chrysalis, especially of those who are metamorphosed in their own skin, seem to prove, by their hardness, an essectual bar to that event. About the time, however, of its change, the head of the sly is capable of being swollen and dilated in an unusual manner; and it is by means of this dilatation that the sly bursts its covering, and opens a passage for its escape.

The feod of flies is as various as their different kinds; fome being supported wholly by vegetable substances, while others are carnivorous; and among the latter there are some who only devour slesh in a state of putresaction. Dung, and vegetables in a similar state, are the favourite meals of others. In the history of the æstrus, we have already noticed that instinct by which the semales are directed to deposit their eggs within reach of food proper for each species.

The female flies are all fecundated by copulation; an act in which the most incurious observer must have frequently seen them engaged. The observations of Reasurar seem to prove, that the semales of some kinds take that part in the act of secundation which is usually performed by the male in other animals; and that it is her extremity which penetrates into the body of the male. The far greater part of insects are oviparous; some, however, belonging to this genus bring forth living young: But what appears most unaccountable is, that some apparently of the same species are viviparous, while the rest

lay eggs; a circumstance that thews the futility of our attempts to arrange them. Some of the viviparous flies possess a degree of fecundity that must appear altogether incredible to those who have not been versant in the study of insects. Some of them have been found to contain in their body no less than twenty thousand living animals at one time *.

Musca domestica, the house fly. The thorax of this species is brown, having upon it sour occult dark lines; the abdomen is of an orange brown, with a few spots of black. This fly seldom makes its appearance before the month of July +.

Musica vespa similis, the wasp sly, entirely resembles the animal whose name it bears; the head is lemon coloured; the antennæ are brown; the thorax and abdomen are black, the latter marked with transverse bands of yellow.

Musca pellucens, the transparent fly. The thorax is black, partially covered with a few brown hairs, and having the point sometimes yellow: The abdomen has its under part black, the upper white, and is transparent both above and below ‡.

Reaumur, Tome IV. Mem. x. + Harris's Inf. p. 144.

[#] Barbut's Gen. Inf. p. 301.

GENUS IV .- Tabanus. The Horse Fly, or Flogg.

The infects of this genus are the great tormentors of horfes and black cattle; their probofels is shaped like a sting; and with it this fly sucks out the blood of larger animals, which is its favourite food. Notwithstanding its diminutive fize, and apparent infignificancy, it engages with the bull, and at once enjoys the honours and the struits of victory: For in spite of all his efforts, it gluts itself with his blood. Cattle, either when intended for fattening or for milk, are greatly harrassed and injured by these insects in the hot months; and whoever would prevent their mischief, would certainly be entitled to those honours and rewards that are due to the discoverers of useful improvements in rural economy.

Tabanus bovinus, the ox fly. In this species the head is grey; but the eyes, which occupy the greater part of it, are brown. The thorax is of a grey colour, and the abdomen yellow.

Tabanus niger, the dark brown. This species varies considerably in colour; some describe it as approaching to black; others grey, and some dark brown. The wings are grey; the feet brown; and the segments of the abdomen are black, with a clay coloured edging upon each. This insect is a great termenter of the rein deer in Lapland.

[#] Fauna Swecica, Ord. VI. Gen. vii. † Regne Animale, p. 198.

GENUS V .- Culex. The Gnat.

The infects of this tribe are extremely numerous, and univerfally known, by the uneafiness they occasion. The Swedish naturalist only enumerates seven species; but there are some places in America where a number of non-descripts might certainly be sound. The genus is discriminated by the mouth, which consists of a slexible sheath, inclosing bristles pointed like stings. The antennæ of the males are filisorm, those of the semales feathered*.

In their larva flate, these animals are aquatic: During the greater part of summer, all stagnated waters are full of their small worms, hanging with their heads downwards, while their hinder parts reach the surface of the water. In this state, the stigmata, or organs of respiration, are situated near the anus; and hence the reason why these parts are frequently seen within reach of the atmosphere. It is in this state, too, the gnat is provided with small sins on its sides, to enable it to go in quest of food.

After having remained in the flate of larva for near twenty days, there infects are transformed into chryfalids, in which all the limbs of the winged gnat are diffinguishable through the diaphanous robe with which they are then shrouded. After remaining three or four days 3 Y 2 wrapped.

wrapped up in this spiral manner, they become gnats, and ascend into a new element. No sooner does the chrysalis reach the surface of the water, than the insect with its head bursts the shell, which then serves it for a boat, of which the wings are the fails. If in this critical moment a croeze arise, it proves to these pigmy sailors a dreadful hurricane; for it oversets the little bark, and the insect not being yet disengaged from it, suffers a fatal shipwreck. If, however, the weather prove calm, the gnat makes a more prosperous voyage: Having time to dry his wings before leaving the boat, he is enabled to mount into the air; where, contemptible as he may seem, he soon becomes the inveterate tormentor of the lords of the creation *.

Cules cinercus †. This species is the most numerous and common; it is called by the Swedes Mygg ‡, a name it has also received in some parts of Britain. The body is of an oblong shape, and the colour is cinereous. It is this species which, towards evening, is so troublesome to man and other animals. Its efforts are far more feeble and temporary than those of the musquito in North America, where the inhabitants are obliged to have their beds close hung with a thin cloth, called Musquito Gurtains, to prevent their intrusion.

Reaumur, Tome IV. † Fauna Swecica. † Vide Flora Lapponica, p. 363, & Blankner, p. 171.

GENUS VI .- Empis.

The empis is distinguished by that horny substance which forms the proboscis; it is bivalved, inclining downwards under the breast, and is longer than the thorax; the valves are horizontal *. The antennæ of these insects are composed of three articulations; the first is long, the second short and globular, and the third much larger at its base than in the middle; from whence it again grows larger, and is finally terminated by a long sharp point †:

The empis, in its winged state, is seen frequenting slowers; but its manners and residence in its larva and chrysalis state are but impersectly known to the naturalist.

Empis livida. This species is of a leaden colour; the thorax is gibbous, of a pale green, and marked longitudinally with three black lines: The upper part of the abdomen is dusky brown: and the legs, which are long and hairy, are of a deeper shade of the same colour.

⁹ Lin. Syft. Ord. VI. Gen. vi. 4 Barbut, p. 311.

GENUS VII .- Conops.

THERE are thirteen different kinds of this tribe enumerated by naturalists, who have distinguished them by a striking character drawn from the rostrum, which is porrected, and jointed like the knee. These insects are universally spread; and from their air and general appearance are commonly taken for certain kinds of wasps. They come later in the season than the horse slies, but are much greater tormentors of the horses, were they equally numerous.

Conops vefpa. The antennæ of this infect terminate in a broad articulation, like the mouth of a fpoon; and which, when closely examined, appears very hairy. The feet and poifers are dun coloured; the thorax is variegated with black and reddiff casts of the same colour, which also prevails in the abdomen. These infects are of a large size, and many of them adorned with very vivid and resplendent colours.

[†] Vide Barbut, p. 210, 311.

GENUS VIII .- Afilis.

This tribe is much more flender in the body than the preceding; many of them are superior in length; and they all glow with the most brilliant tints of green, purple, yellow, and brown. The generic characters are, a horny rostrum, porrected, strait, and bivalved †.

These animals are generally found in swampy meadow ground, where, like the preceding tribe, they prove very troublesome to cattle. They are provided with a sting, which inslicts a very painful wound, if you incautiously endeavour to lay hold of them. The proboscis, though sharp enough to perforate the thickest skin, is nevertheless perforated, and supplies the place of a pump, in sucking out the blood of animals.

Inufca crabroniformis ‡, the ferruginous afilis. The thorax is gibbous; the abdomen oval, flender, and tapering §; the three first segments of the abdomen are black, the rest yellow; the feet and wings are of an iron colour. The graziers imagine, and perhaps with justice, that of all the insect tribe, this is the most noxious to black cattle; the very noise of their wings sets them arunning; and before they stop, they generally make their escape into a river, if any is near ¶.

⁺ Syft. Nat. p. 1006.

[†] Vide Mouffet, p. 46.

[§] Actes d'Upfal, 1736, p. 29.

Moffnagel, t. 16.

GENUS IX .- Bombilius. The Buzz Fly.

THE British infects of this genus, in the month of April, are feen hovering about the garden flowers, generally in damp fituations. Even when in the act of fipping the nectarous dew from plants, they do not fettle, but keep constantly buzzing upon the wing. They have no sting, and may be handled with impunity.

The characters of this genus are, a roftrum very long, fetaceous, and porrected, formed by two horizontal valves; antennæ which are bent, fetaceous, and of a conic form; three stemmata, and two spread wings.

Bombilius medius †. The probofcis of this infect is black, and flender, fometimes divided in two at the extremity, which extends nearly three fourths of the animal's length: the legs are of the fame colour, flender, and long in proportion to the fize of the infect. The body is fhort and truncated; the ground colour black, but covered externally with dun coloured tufty down; the wings are unufually long, dark, and dotted with brown ‡.

GENUS X .- Hippolofca. The Horfe Leech.

In the horse leeches, the rostrum is of a cylindrical shape, bivalved, and nodding: the feet are armed with claws; and it is by means of these instruments that they adhere to cattle and horses, in spite of every effort to rub them off. They abound in New Forest, Hampshire, and render it a very uneasy pasture for cows. The horse leech is covered with a crustaceous shell, which renders it difficult to be killed, and gives it the air of some species of the bugs.

Hippolofca pedibus pidactitis, the horse leech with four talons on each foot. This species is but too generally known, from its hostilities against every kind of cattle. It is so tenacious of life, that it will live for a considerable while after the head is cut off. The colour of this hippolosca is wholly grey *. There is a species of these animals with fix talons on each foot, which is sound in swallows ness; it is much smaller in size, and has the extremity of the abdomen marginated f.

^{*} Ricinus volans of M. Frisch, p. 43. † Reaumur, Tome V.


CHAPTER VIII.

SECTION I.

ORDER VII .- Apterous Insects.

A MULTIFARIOUS group of animals are comprehended under this order, which are all characterifed by the privation of wings in both fexes. All the varieties of crabs and lobsters are introduced by *Linnœus* into this class of the animal kingdom; and that with great propriety, because they possess the antennæ, those organs which are peculiarly characteristic of the insect tribes.





GENUS I .- Lepisma.

THE infects of this genus are diffinguished by possessing fix feet formed for running; a mouth furnished with four pulpi; two of which are setaceous, and two capitated. The body is covered with imbricated scales; and the tail is terminated by bristly appendices †.

The refidence of the lepifma is commonly in gardens and damp houses. In the former, they retreat under mouldy boards or flag-stones; and in the latter, they are feen about window sashes, and old boxes that have remained for some time in a humid situation. When their haunts are disturbed, they make off with so great velocity, that they are but seldom caught; and when that happens, they are often destroyed, being, from their softness, easily crushed.

Lepisma faccharina ‡. The colour of this infect is a shade of lead, brightening into silver: Its most conspicuous member is the tail, which consists of three long threads, slender and sine; but if viewed with a microscope, are covered with tusts of hair. The antennæ of this lepisma are long and siliform; in some, they are nearly equal to the length of the body: The seet are six in number, presenting a short and truncated appearance at their insertion: They are received into a groove in the body

3 Z 2 of

of the animal, where each has its origin, covered with an oval feale. Towards the tail there are fix other appendices, much florter, which may be termed false feet, as they seem to give no assistance to the progressive motion of the animal *. There are only three species of this genus enumerated in the System of Nature.

^{*} Barbut's Gen. Insect. p. 322.

GENUS II .- Podura. The Springtail.

THERE are above a dozen of different kinds of the springtail; they derive their English name from their habit of bounding from the ground; an exertion which is performed by means of the forked elastic tail which bends under the body, and acts like a spring. The antennæ of the springtails are long and setaceous; their six seet are formed for running; and their eyes are composed each of eight facets †.

Podura villosa, the rough springtail, is a British insect, and the largest of the tribe known to us. The ground-colour is dark brown, mixed with a shade of yellow, and variegated all over with streaks of black. The abdomen of this insect alone is smooth, the head being very much covered with hairs, which are so deciduous, that they stick to the singers on handling the animal; the antennæ, in the subject we saw, consisted of sour articulations, and were nearly as long as the body.

This lepisma is generally found below stones; but there are others of the genus that seem purely aquatic, and able to walk and skip upon the surface with apparent ease. It is this species that we may observe in the mornings assembling in pretty numerous tribes upon the banks of stagnated pools. Other species reside among rotten bank, or decayed leaves; and in Sweden there is a species which

runs

runs upon the snow, but perishes there as soon as it begins to thaw +.

Podura atra terrestris alba t. This species frequents melon beds and hot-houses; where, on watering the ground, numbers are feen jumping into the air like atoms: They are the smallest of the poduræ, and of a white colour.

† De Geer, Actes d'Upfal, 1740, p. 54. ‡ Fauna Swee. Gen. iv.

GENUS III .- Termes.

THE discriminating characters of these insects are drawn from the mouth, which is armed with two jaws; from the setaceous antennæ, and the six seet, which are formed for walking †.

There are only three species of these small infects; to one of which Linneus has given the name of Pulsatoria, from its being supposed to imitate the ticking of a watch, by striking its head against the wainscot in the dead of night. This noise, so alarming to some superstitious minds, has been called the death watch, and is probably occasioned by the plinus one of the colcopterous insects.

The termes pulfatorius varies in its colour according to its refidence: When met with in houses, it is commonly of a whitish colour; if found in the fields, as is sometimes the case, it tends more to a brown colour, and is somewhat hairy ‡.

Syft. Nat. p. 1015.

[‡] Barbut's Gen. Insect. p. 326.

GENUS IV .- Pediculus. The Loufe's

THE characters of this unfeemly tribe are, fix feet formed for walking; a mouth containing an exferted fling; antennæ as long as the thorax; and an abdomen deprefed, and formed of different lobes or fegments +.

It is remarkable that almost every animal and vegetable is frequented by its particular louse, although only about fifty different species of this tribe have yet found their way into the systems of natural history. Even man, whose fancied prerogatives lead him to boast of being the lord of the creation, has often the mortification of being trampled upon by the most despicable of all animals.

Pediculus bumanus, the human loufe. Of this infect there are two varieties; the one inhabits the head, and the other the body. No medical impuries have yet found out why fome families of the func tribe choose these different abodes. Infants and children are most infested with the former of these kinds, and failors, foldiers, and old men are most liable to be troubled with the latter. Cleanliness is the best antilote against these hideous intruders; and their nausons society seems a very proper punishment instituted by nature on its neglect. The pthirish, or loufy diffeste, though very little known in latter times, was not unfrequent among the ancients. Herod, Antiochus, Callishenss, and Sylla, with many others, died

of this diforder. It is uncertain whether this is to be afcribed to their inattention to cleanliness, their scarcity of linen, or their ignorance of the use of mercury. However this may be, there is no person in our climate attacked by these animals, except such as by sloth and nastiness invite their company.

We will spare our readers the trouble of perusing a more minute description of this motly tribe: Such, however, as are not satisfied with their personal knowledge, we refer for farther particulars to Rhedi*, Swammerdam†, and Bonani‡, where they may enjoy an intellectual feast, of which sew will envy them.

* Exper. t. 18. † Quart. 169. ‡ Micogr. f. 55.

Vol. III. 4 A

GENUS V .- Pulex. The Flea.

I'ms genus confils of two species, which are characterifed by fix feet formed for leaping; filiform antonna; a rostrum bent inwards, setaceous, and concealing a sting: The abdomen is laterally compressed, and covered with imbricated lamina.*.

This tribe, like the preceding, is blood-thirfly, and fattens at the expence of the human race. It is faid to prefer the more delicate skin of women; but feeds neither upon epileptic persons, upon the dead, nor the dying. Other animals, as well as man, suffer by its intrusions. It neitles in the fur of dogs, cats, and rats; and the nests of the river swallow swarm with them †.

Pules irritans, the bed flea. In this animal, the antennæ are longer than the thorax; the body of a dark brown colour, and of an oval form; it is apterous, walks but little; but by the eladicity of its legs, can fpring above two hundred times the length of its own body. From microscopic observations which have been made upon this animal, it has been found to be oviparous. The eggs are deposited on blankets, or among the down of animals; they are round and smooth; and in four or five days, give birth to a very minute worm, which feeds upon greafy fur. After having crept about in this larva

Syft. Nat. p. 1021. + Syft. Nat. du Regne Animale, p. 232.

flate for a few days, it fpins a filken cod, in which it lies dormant for about a fortnight; after which it burfts from its confinement, endowed with powers to diffurb the peace of an emperor.

Preparations of mercury are effectual in destroying the slea, and most other noxious vermin. In some places, bugs, sleas, and other abominable intects, are held in superstitious veneration. At Surat, there is an hospital endowed for their preservation; where every night they prevail upon some poor wretch to suffer himself to be preyed upon for their sustenance. They sasten him naked upon a bed, where he is deprived of the means of self desence; and the samished pensioners riot in his blood during the whole night *. Another pious soundation of a similar nature, but less ludicrous, is crected in Turkey, for decayed or infirm dogs.

. * Barbut's Genera, p. 332.



GENUS VI .- Acarus. The Tick.

THE ticks have all eight feet; two cycs, placed one on each fide of the head, and two articulated tentacula, in the form of feet *. The tribe contains thirty five different species, many of which have been inaccurately placed in the genus of lice.

The ticks are all produced from eggs; the females besing oviparous. Many of the species of this genus are carnivorous, and prey upon man, as well as other living creatures. When, however, this food is not to be procured, they are capable of deriving their subsistence from plants and vegetables. The dysentry and the itch, two insupportable diseases, are both occasioned by some of the smaller acari infinuating themselves into the human body. In the latter of these complaints, they dig under the skin, and occasion that raging pruriency with which the unhappy patient is afflicted †.

Acarus longicornis is found below stones and the bark of trees. The abdomen and legs are of a red colour; and the whole infect fomewhat resembles the shape of a pear. The antennæ resemble those of the scorpion, the articulations being proportionably smaller and more slender; their extremities terminate in a forceps formed by two threads ‡.

Some

Syft. Nat. p. 1022. † Acarus Scabici, Actes d'Upsal, 1736. p. 37.

Some of the acari are well known to frequent cheefe; others of them attach themselves to the legs and bodies of bees, spiders, and other insects. The pediculus pubis of Rbedi, the cow and sheep louse, seem all to belong to this tribe: The former is engendered, or at least multiplied, by uncleanliness upon the pudenda, where it is more vexatious than the louse *.

The monoculus, or arborefeent water flea, may also be added to the tribe of acari †. This infect, when viewed with the assistance of the microscope, appears to have but one eye; for, on account of the smallness of the head, the two seem to be joined together. When you view them with a lens, however, both appear, and are of a reticulated structure. Its trunk then also is seen, small, sharp, and transparent.

This infect is of a blood red colour; and its numbers, that float on the furface of flagnated water, are fometimes fo great, that they give it the appearance of blood. Swammerdam mentions an inflance in which the people of Leyden were greatly alarmed by this colour of the water, deeming it portentous of some great calamity.

^{*} Rhedi Expos. t. 18. + Swammerdam, Quart. p. 66.


GENUS VII .- Phalangium.

THE infects of this family bear a strong resemblence to the crab: They have eight feet, and four eyes; two on the summit of the head, and two on the sides. The antenne are fixed to the forespart of the head, and have the appearance of feet; the abdomen is round *..

Some have imagined that all those threads with which the grass and stubbles in autumn are covered, are the production of the phalangia. So numerous are these threads, that in the course of one day the whole surface of the ground is covered with them. Their uses to the animal are probably to ensure its prey, and to affish it in travelling through the air. Other naturalists have supposed these waving threads to be spun by a species of tick called the autumnal weaver.

Phalangium opilio, the shepherd weaver. The body of this species is round; above, the colour is a dusky brown; below, a dirty white. The legs are uncommonly long, and slender: when caught by one of them, the owner parts with it to save his body, and makes off without any apparent uneaffiness: This is also the case both with the crab and lebster, whose loss is quickly repaired by the growth of a new limb. No experiments have yet ascertained whether the power of reproducing their limbs may not also belong to the insects of this genus.

^{*} Syft. Nat. p. 1027.





GENUS VIII .- Aranea. The Spider.

The infects of this family may be considered as a kind of Arguses, and possessing eight eyes, and distinguishable into families by their arrangement and position. They have eight seet, and a mouth armed with crotchets. The instruments for spinning are shaped like nipples, and placed in the anus.

But the circumstance by which the spiders are most distinguished, are the two articulated pulpi, which are headed by the genitals of the male in that fex †. As the spiders prey upon each other, except during the time of their amours, they dare not venture within reach of each other, but with the utmost caution. They are then seen stretching out their legs, shaking their webs, and tempering with each other, by a slight touch with the extremity of the feet; then, in a fright, dropping hastily down their threads, and returning in a few moments to make a fresh trial by feeling. When once both parties are well affored of the fex hey have to deal with, the approaches of their feet to seel become more frequent, considence takes place, and the instant of amorous dalliance ensure ‡.

We cannot, fays L, onnet, fufficiently admire how careful these au mals are, not to give themselves up blindly to a passion, or venture upon an imprudent it p, which might prove faul to them. After the male and semale spider

⁺ Syft. Nut. p. 1030. + Barbut; p. 339.

fpider have thus met, it is afferted by the most accurate observers, that the extremity of the claws of the former opens, by a kind of spring, and lets out a white substance, which he applies beneath the abdomen of the semale, and thus accomplishes the great purpose of nature †.

In this country, where all the infect tribes are kept under by the coldness of the climate, or by human affiduity, the spiders are small in fize, and comparatively few in number: Here, they are an inossensive, as well as a humble tribe; but in the warmer regions of Africa and America, they are hideous and terrible. There, the spiders, as well as all the insect tribe, acquire a size and vigour, which to the inhabitants of the north of Europe must appear incredible. The abdomen of the Martinico spider is as large as a hen's egg, and covered all over with hair: Its web is strong; and its bite is dangerous. Happy it is for us, that we are placed at such a distance from these animals, that we can study their history without dreading their resentment.

The eyes are placed all around the head; they are immoveable, and destitute of eyelids, but are sortified with a transparent horny substance, which at once protects and assists their vision. A number of eyes so unusual among the rest of animated beings, seems to have been deemed necessary to an infect destined to procure its support by the most watchful attention. The mouth of the spider is equally sitted for his occupation: It consists of two serrated pincers, which spring from the forehead, and terminate like the claws of a cat. A little below the point of the claw there is a small hole, through which the ani-

mai

mal emits a poison; which, though harmless to us, is capable of instantly destroying its prey.

But all these arms of the spider, formidable as they are, would prove insufficient for producing him subsidence, did he not add stratagem to force. Every one is acquainted with his invention of spreading a net for the sty, an insect which, without this artifice, must for ever have escaped him. The position of the spider's web is generally well chosen; being placed either in the corners of rooms, the sides of windows, or among the branches of trees, where slies are most frequent and plenty.

For the confiruction of this delicate engine, Nature has fupplied these animals with a large portion of a glutinous substance, which she has granted them the power of spinning into a thread so delicate, that ninety of them twisted together are only equal in thickness to that of the common silk worm. Every thread of the common spider's web, though scarcely visible by the naked eye, is composed of a vast number of slender silaments joined together. The instruments of this delicate operation are sive dugs or teats, with small orisices, capable of being contracted or dilated, according to the sineness of the instended thread.

After the net has been thus formed, the next care of its owner is to provide a proper retreat, from which he may furvey the whole, and observe the success of his own artifice. Some threads being in contact with his body, whenever the web is touched from without, he instantly feels the motion, and accordingly prepares either for attack or desence. If the insect invading his territories prove a fly, he immediately fallies forth, and devours the devoted prey with all that ferocity which distinguishes Vol. III.

the lion or the tiger. If, however, the affault be committed by an enemy stronger than himself, the spider keeps close concealed in his fortress; and it is not till the danger is over, that he ventures forth to repair the damages which his property has sustained.

By fuch accidents, as well as the force of the wind, the labours of many days are often infantly defroyed a and either the conftruction of a new web, or a repair of the old, becomes absolutely necessary. Here the spider has again recourse to that glatinous substance with which he was originally provided. The life of this animal feems chequered with misfortunes. The time feldom fails to arrive, when the receives of gluten, by repeated draining, become entirely exhausted; and the poor spider is left to all the chances and vicissitudes of want, which often bring him to an untimely end *.

The natural longevity, however, of the spider, is greater than seems to have been granted to the generality of the infect tribes. The length of his life is not exactly ascertained; but it probably endures for several years. The semale, it is said, does not begin to lay her eggs till she has completed her second year; and even then, her brood is not so numerous as when she has attained her full maturity. When that period arrives, she has been known to produce near a thousand eggs in a single season. The eggs being thus deposited, the animal prepares a silven bag for their reception, till they are hatched. Thus parted up, the glues them to her body, by means of the single adhesive matter with which the web is spun; and

113

^{*} Goldsmith's Nat. Hist. Vol. VII.

in this manner they are carried by the parent, like and ther body attached to her own.

With all this tenderness for its young, there is no animal more ferocious, or more powerfully armed for de-Aruction than the spider: It can destroy infects far above its fize; and its indifcriminate rapacity does not even fpare those of its own kind. Reaumur, when he endeav nued to turn the labours of different infects that fain fills to the advantage of man, was foiled in all his attempts to domesticate the fpider. He provided them with a habitation, and with every kind of food, flies, and blood in the ends of unripe feathers, a morfel of which they are particularly fond: but all was in vain; their malignant nature rendered them unlit for fociety, and turnut them from their food to destroy each other. By perseverance, however, he obtained as much of the filk intended for the covering of their eggs, as enabled him to manufacture a pair of gloves of these curious materials.

Aranea hortensis. This species frequents the garden trees, among the branches of which it spins its web, and stations itself in the centre. The body is of a brownish colour; the eyes very small, and of a violet purplet. With the silk of this spider M. Bon of Languedoc contrived to manufacture a pair of silk stockings, of a sine grey colour, and in strength and elegance nearly equal to those of common silk.

Aranea domestica. This species inhabits windows and unfrequented apartments; it is oblong, and of a brown colour; the abdomen black. This animal has four ca-

† Lifter, Aran. 83.

vities upon the buck, by which it may easily be distinguished. Between the nails of the feet there issues a clammy liquor, by which it is enabled to climb up glass and other smooth surfaces. The web of this species is constructed of much finer threads. It is said to possess fix muscular nipples, each having a thousand orisices for letting out the threads. If this be true, each silament of the spider's web must consist of fix thousand threads, which, when united, are scarcely visible to the naked eye.

To this tribe is commonly referred the tarantula, the the largest spider known in Europe; but rendered still more remarkable by the extraordinary qualities which credulity has ascribed to it. It is three quarters of an inch in length, the body covered all over with down, generally of an olive dusky brown. The external parts resemble those of the rest of the genus: It has eight eyes and as many legs; the front armed with serrated pincers.

The bite of the tarantula is faid, but without foundation, to be attended with fatal confequences. After violent pains and fickness, the unhappy victim is seized with an involuntary laughter; dances without intermission, exhibiting all the symptoms of infanity. At the return of the season in which he was bit, the patient's madness begins again to appear; and these troublesome relapses, after having recurred for several years, at last terminate in death.

Music and dancing, as the siction proceeds, is the only remedy for this dreadful malady. The musician begins to play a tune famous for the cure, slow at first, but increasing

[†] Fauna Swecica. † Vide Reaumur, Barbut, and Goldsmith.

creating in quickness as the patient seems affected. After continuing this ridiculous, but violent exercise, till a profuse perspiration is brought on, the imaginary venom is forced from the system, and the cure is completed. Such were the reputed effects of the bite of the tarantula, till travellers of discernment traced the whole siction to the artistice of the peasants of Apulia, who, for a small sum had been in the habit of allowing themselves to be bitten, and then counterseited madness, to associate two dering multitude.

GENUS IX .- Scorpio. The Scorpion.

WE are now to contemplate the manners of an animal, the most hideous, and perhaps the most formidable of the infect tribes. There are fix different species of the fortion, diffinguished by their fize and colour, and the countries where they refide. Happily for Britain, none are found there, but a species no bigger than a louse; and even that is extremely rare +. In Italy, Spain, and the fouth of France, they are frequently to be met with three inches in length, and are confidered as the greatest pells that torment mankind. But the fize and malignity of the feorpions of Europe may be deemed trifling, when compared with that of the African monsters that are diflinguished by that name. Along the Gold Coast they are fometimes found larger than a lobster, and their sting is inevitably fatal t. From the language of Scripture, too, we find that in the east thefe animals have long been formidable to mankind. In Batavia, they fometimes grow twelve inches in length; and in removing furniture, behind which they skulk, there is the utmost danger of being stung.

The genus of fcorpions is characterifed by eight eyes, three of which are placed on each fide of the thorax, and

two

[†] Barbut's Gen. Inf. p. 351.

two almost contiguous upon the back. Besides eight feet, they have two palpi, each armed at the extremity with a forceps. The tail is lengthened out by different articulations, the last terminating in a sharp and crooked sling.

Scorpio Afer. This terrible and venomous animal is an inhabitant of Africa: Its colour is a deep gloffy brown; the fegments of which the body is composed, each terminate in a band of bright yellow; while the two eyes sparkle in the back like diamonds. Its appearance bears a strong resemblance to the looster. Most of the travellers who have visited Africa represent this animal not only as possonous, but exceedingly irascible; and the effects of its resentment, unless speedily relieved, become stated.

It is to M. de Manperiuis that the public is indebted for the first accurate and philosophical account of these animals. According to him, they are all viviparous; the body of the pregnant semale exhibiting, when opened, between forty and fifty living young. Each of these is separated from the rest by a thin membrane, while all are united by a common slament. In order to ascertain the venom of their sting, that philosopher bred a great number together, and let them loose upon dogs and other animals; and he sound that sometimes the sting was so empoisoned as to swell the whole body of the wounded animal; and in that case it brought on violent retching, convulsions, and death. At other times, whether from the difference of food, or some other cause, he found their sting harmless.


The

The common food of the feorpion are worms, flies, and spiders. The Italian scorpion, when inclesed with the latter animal in a glass bottle, exhibits a dreadful combat, which terminates in favour of the scorpion. After having stung his adversary to death, he cuts off his limbs with his claws, and then sucks the juice from his body. No observations nor experiments of naturalists have yet enabled them to assign the cause why the sting of the scorpion, which at one time is satal, proves harmless at another; nor has the medical art yet sound out any certain antidote against its malignity. The esseacy of the oil of scorpions, and some other remedies that are used, is perhaps rather to be ascribed to want of activity, or the total absence of the poisonous matter.

Fphemef. Dec. II. 1687. Observ. 224.



Plate XIII.



GENUS X .- Cancer. The Grab.

After the example of Linneus, we place the crab among the infects, because it is possessed of some of the invariable characters of this class of beings. All the eighty-seven different species of crabs are distinguished in possessing two members performing the office of hands, and terminating each in a forceps. They have generally eight seet, and two moveable eyes, projecting from the head, or placed upon a pedicle or stalk. The two palpi are armed with claws; the tail is articulated, and unarmed.

In general, the crabs are aquatic animals, whose residence is in the sea: Some, however, are known to frequent rivers; and in America, there are some species wholly terrestrial; these are commonly about four inches long, and seed upon leaves. The liquid substance found in their shell cures the blisters that are raised on the skin by the milky substance that exsudes from the mancellina tree. The savages apply the sat, when melted down, as a specific against the rheumatism to

Cancer Bernardus, the St. Bernard or hermit crab. This species is small in size, and semicrustaceous. As it has no scales to defend it, it takes up its dwelling in the empty shells found lying upon the shore. Of these it tries various kinds, till at length one is pitched upon,

Vol. III. 4 C fitted

⁺ Barbut's Gen, Inf. p. 358.

fitted to its shape and fize: the increase of its bulk, however, soon forces it to exchange this for a different and more capacious habitation; again, therefore, it sallies forth in quest of a new mansion. Should it meet with another crab engaged in the same pursuit, they frequently are seen quarrelling for the same shell. This battle, pro aris et focis, often proves obstinate, till the vanquished sneaks off; and the empty apartment becomes at last the prize of the victor. Of the house thus obtained, they are so tenacious, that nothing but heat will disengage them from their retreat. When caught, they pinch forcibly, and utter a faint cry.

Cancer curfor. This animal has obtained its name from the nimbleness with which it runs. It is amphibious, leaving the sea towards noon, and parading upon the thore; from which, however, it speedily makes its retreat on the approach of danger †.

The common lobster belongs to this genus, the species of which are far more numerous than naturalists have yet allowed. This animal is so well known, that a description of it can hardly be necessary. It is sound all around the shores of Britain, frequenting particularly the stony ground, where it is caught by a creel or willowwork of a particular construction, that admits of its entering, but precludes any return. The bait by which it is enticed into this prison is fish guts.

Every lobster is supposed to be felf impregnated, and to exhibit an example of an hermaphrodite of the most perfect kind. The ovary where the spawn is first produced, lies towards the tail, and is filled with that red substance.

[†] Barbut, abi fupra.

fubstance, each particle of which is the embryo of a future animal. When they are first excluded from the belly, they are placed under the tail, where the parent animal protects them from danger, till they acquire limbs and animation. In that state, they drop off into the water, when yet very small, and betake themselves to the crevice of a rock for shelter, till they are enabled by greater strength to search for their prey. When full grown, their voracity is so merciles and indiscriminate, that they mutually devour each other.

The lobsters, like all other infects, possess antennæ, and like thefe animals change their external covering once a-year: To them, this is a painful and tedious operation. The lobster continues growing, while its crust continues unalterably the fame: hence it foon becomes too large for its habitation, and is necessitated to get free. During their transformation, all their vigour and activity ceases; and they no sooner feel its approach, than they betake themselves to some retired situation, where they may be fafe from the attacks of their enemies. The claws, and even the stomach, are said to be changed in this convultive effort of nature. Before casting the shell, the animal throws itself upon its back, strikes its claws against each other, and every limb seems to tremble; its feelers are agitated, and its whole body is in violent motion. It then swells itself in an unusual manner, and at last the shell is seen beginning to divide at its junctures, particularly at the belly, where it was least firmly united +.

4 C 2

In

572

In this painful manner does the lobster attain its liberty; but so feeble is its condition, that it remains for hours motionless, from the satigue of the operation. Many die in this violent criss of their existence: Such as survive keep close to their dens for several days, shunning danger with all the timidity of worms, whose softness and imbecility they then acquire.

GENUS XI. Oniscus.

This is a diminutive, but harmless tribe of animals: They are distinguished by having fourteen seet, setac our antennæ, and the body of an oblong form. Linnæus has enumerated sisteen species, some of which are terrestrial, and others aquatic t.

Onifeus aquaticus. This insect is found in wells and stagnated pools ‡. The body is composed of seven articulations, besides the head and tail; the last is much larger than any of the rest. The shape of the animal is stat and round: From each side there spring seven seet, each growing longer as you approach the tail.

The sea onisei are larger than those of the fresh water, having ten instead of seven segments. Their motions in the water are rapid; for besides the seet, they are affished by lateral threads, which push them forward like the oars of a boat. Among them, copulation endures for several days; when the male seizes his semale with his two fore seet, and drags her along with him wherever he directs his course; and nothing can induce him to leave his semale till this operose function is discharged. A few days after impregnation, the semale discovers a dissended belly; on the seventh, the young issue alive from

her

[†] Systema Nat. p. 1059. † Rai Ins. p. 43,

A Regne Animale, p. 250, fp. 4.

her womb, and fwim around with alertness and anima-

All this while, the male, it is faid, continues fixed upon his female, not for the purpose of secundation, but of affishing her to cast her slough; a good office; in which he is not successful till he has exerted the whole of his strength.

On her first appearance after leaving her coat, the female is entirely white; her slough, of a deep ash-colour, is left sloating upon the surface, and so entire, that you would imagine it to be the real animal. This transformation being completed, the male leaves his female, and unassisted undergoes a similar change.

Onifcus domestica. This infect inhabits damp houses and walls, and must have been seen by every person. The upper part of the body is dark brown, the belly grey. Two varieties of this animal are found in the fields, having each, like it, the body divided into ten segments, exclusive of the head and tail. The latter part terminates in two appendices *. The land onisci are all supposed to be oviparous, while the aquatic produce living young. For some time after being excluded from the shell, the sormer are often of a pale red colour. The domestic oniscus is gathered for medicinal purposes †.

^{*} Charleton Exercit. 54. † Vide Barbut, p. 365.

GENUS XII .- Scolopendra.

This tribe has in the Latin language obtained the name of mille-pes, from the extraordinary number of feet with which nature has provided fome of its species; one kind of the scolopendra having no less than an hundred on each fide. In general, the feet of these insects are as numerous as the fegments into which their bodies are divided. They have the antennæ fetaceous, articulated palpi, and the body depressed or slattened +. Of this hideous race there are eleven different species, distinguished by their form, fize, and colour. Some live beneath the bark of decayed trees, or are found below stones and garden boxes; others inhabit the fresh and salt water, and are all remarkable for their quick progressive motion.

Scolopendra forficata. The feet of this species are fifteen on each fide; the last longer than the rest, and turning backwards, form a forked tail. The body is of a dun colour, fmooth, and composed of nine scaly segments, without reckoning the head 1.

The marine scolopendra || in form somewhat resembles the leech; it builds those small edifices of a brittle and porous texture which are feen upon the shore at low water. These masses are composed of a number of small funnels.

[†] Syst. Nat. p. 1062.

i Barbat's Gen. Inf. p. 367.

Vide Rai Inf. p. 44.

funnels, each having the aperture closed up with a covering of fand, to protect the inhabitant from danger.

In the East Indies and America, there are scolopendres from four to fix inches long, and as thick as the finger: When of that bulk, they appear truly hideous, and their sting is reckoned as dangerous as that of the scorpion. Some of these animals are said to be without eyes, and to direct their course by means of two seelers that extend beyond the head. It has been afferted, too, that when cut into pieces, each segment, like those of the polypi, is capable of reproducing an entire animal.

^{*} Barbut, ubi fuora.

GENUS XIII .- Tulus.

This tribe fo nearly resembles the former, that in the Fauna Swecica they are both included under one genus. The feet of the tuli are still more numerous than those of the scolopendra, being, on each side, twice as many as the segments of the body. The antennæ are moniliform, the palpi articulated, and the body of a semicylindrical form.


Tulus fabulofus*. This fpecies has no less than two hundred and forty feet, although its length is not more than an inch. It is of an ash-colour, and smooth; the body composed of fixty segments, each giving off two pair of small white feet. The antennæ of this infect are short, and consist of sive rings. When the animal is touched, it wraps itself up into a round ball, the feet all turned inward.

Millepeda orientalis f. This is the largest of all the tuli, being as long and as thick as a man's singer: The head is small, and obtuse, the antennæ clavated, and the eyes black; the body is of a pale colour, and divided in forty-three segments, each having a pair of seet on every side, the whole amounting to an hundred and seet venty-two ‡.

* Rai Inf. p. 47. † Seba, I. p. 131. & Regne Anim. p. 253. sp. 3.

THE END.

Printed by 3 Mundell and Son. \$





SMITHSONIAN INSTITUTION LIBRARIES
3 9088 01506 5311